

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 365 034 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
26.11.2003 Bulletin 2003/48

(51) Int Cl.⁷: **C12Q 1/68, B01L 3/00,
A61K 48/00, G06F 17/00**

(21) Application number: **03010447.5**

(22) Date of filing: **09.05.2003**

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PT RO SE SI SK TR**
Designated Extension States:
AL LT LV MK

(71) Applicant: **Bayer Aktiengesellschaft
51368 Leverkusen (DE)**

(72) Inventors:
• **Wirtz, Ralph, Dr.
50674 Köln (DE)**
• **Munnes, Marc, Dr.
40699 Erkrath (DE)**
• **Kallabis, Harald, Dr.
50161 Köln (DE)**

(30) Priority: **21.05.2002 EP 02010291
13.02.2003 EP 03003112**

(54) **Methods and compositions for the prediction, diagnosis, prognosis, prevention and treatment of malignant neoplasia**

(57) The invention provides compositions, methods and uses, for the prediction., diagnosis, prognosis, prevention and treatment of malignant neoplasia and breast cancer in particular. Genes that are differentially

expressed in breast tissue of breast cancer patients versus those of normal people are disclosed.

EP 1 365 034 A2

Description

TECHNICAL FIELD OF THE INVENTION

[0001] The invention relates to methods and compositions for the prediction, diagnosis, prognosis, prevention and treatment of neoplastic disease. Neoplastic disease is often caused by chromosomal rearrangements which lead to over- or underexpression of the rearranged genes. The invention discloses genes which are overexpressed in neoplastic tissue and are useful as diagnostic markers and targets for treatment. Methods are disclosed for predicting, diagnosing and prognosing as well as preventing and treating neoplastic disease.

BACKGROUND OF THE INVENTION

[0002] Chromosomal aberrations (amplifications, deletions, inversions, insertions, translocations and/or viral integrations) are of importance for the development of cancer and neoplastic lesions, as they account for deregulations of the respective regions. Amplifications of genomic regions have been described, in which genes of importance for growth characteristics, differentiation, invasiveness or resistance to therapeutic intervention are located. One of those regions with chromosomal aberrations is the region carrying the HER-2/neu gene which is amplified in breast cancer patients. In approximately 25% of breast cancer patients the HER-2/neu gene is overexpressed due to gene amplification. HER-2/neu overexpression correlates with a poor prognosis (relapse, overall survival, sensitivity to therapeutics). The importance of HER-2/neu for the prognosis of the disease progression has been described [Gusterson et al., 1992, (1)]. Gene specific antibodies raised against HER-2/neu (Herceptin™) have been generated to treat the respective cancer patients. However, only about 50% of the patients benefit from the antibody treatment with Herceptin™, which is most often combined with chemotherapeutic regimen. The discrepancy of HER-2/neu positive tumors (overexpressing HER-2/neu to similar extent) with regard to responsiveness to therapeutic intervention suggest, that there might be additional factors or genes being involved in growth and apoptotic characteristics of the respective tumor tissues. There seems to be no monocausal relationship between overexpression of the growth factor receptor HER-2/neu and therapy outcome. In line with this the measurement of commonly used tumor markers such as estrogen receptor, progesterone receptor, p53 and Ki-67 do provide only very limited information on clinical outcome of specific therapeutic decisions. Therefore there is a great need for a more detailed diagnostic and prognostic classification of tumors to enable improved therapy decisions and prediction of survival of the patients. The present invention addresses the need for additional markers by providing genes, which expression is deregulated in tumors and correlates with clinical outcome. One focus is the deregulation of genes present in specific chromosomal regions and their interaction in disease development and drug responsiveness.

[0003] HER-2/neu and other markers for neoplastic disease are commonly assayed with diagnostic methods such as immunohistochemistry (IHC) (e.g. HercepTest™ from DAKO Inc.) and Fluorescence-In-Situ-Hybridization (FISH) (e.g. quantitative measurement of the HER-2/neu and Topoisomerase II alpha with a fluorescence-in-situ-Hybridization kit from VYSIS). Additionally HER-2/neu can be assayed by detecting HER-2/neu fragments in serum with an ELISA test (BAYER Corp.) or a with a quantitative PCR kit which compares the amount of HER-2/neu gene with the amount of a non-amplified control gene in order to detect HER-2/neu gene amplifications (ROCHE). These methods, however, exhibit multiple disadvantages with regard to sensitivity, specificity, technical and personnel efforts, costs, time consumption, inter-lab reproducibility. These methods are also restricted with regard to measurement of multiple parameters within one patient sample ("multiplexing"). Usually only about 3 to 4 parameters (e.g. genes or gene products) can be detected per tissue slide. Therefore, there is a need to develop a fast and simple test to measure simultaneously multiple parameters in one sample. The present invention addresses the need for a fast and simple high-resolution method, that is able to detect multiple diagnostic and prognostic markers simultaneously.

SUMMARY OF THE INVENTION

[0004] The present invention is based on discovery that chromosomal alterations in cancer tissues can lead to changes in the expression of genes that are encoded by the altered chromosomal regions. Exemplary 43 human genes have been identified that are co-amplified in neoplastic lesions from breast cancer tissue resulting in altered expression of several of these genes (Tables 1 to 4). These 43 genes are differentially expressed in breast cancer states, relative to their expression in normal, or non-breast cancer states. The present invention relates to derivatives, fragments, analogues and homologues of these genes and uses or methods of using of the same.

[0005] The present invention further relates to novel preventive, predictive, diagnostic, prognostic and therapeutic compositions and uses for malignant neoplasia and breast cancer in particular. Especially membrane bound marker gene products containing extracellular domains can be a particularly useful target for treatment methods as well as diagnostic and clinical monitoring methods.

[0006] It is a discovery of the present invention that several of these genes are characterized in that their gene products functionally interact in signaling cascades or by directly or indirectly influencing each other. This interaction is important for the normal physiology of certain non-neoplastic tissues (e.g. brain or neurogenic tissue). The deregulation of these genes in neoplastic lesions where they are normally exhibit of different level of activity or are not active, however, results in pathophysiology and affects the characteristics of the disease-associated tissue.

[0007] The present invention further relates to methods for detecting these deregulations in malignant neoplasia on DNA and mRNA level.

[0008] The present invention further relates to a method for the detection of chromosomal alterations characterized in that the relative abundance of individual mRNAs, encoded by genes, located in altered chromosomal regions is detected.

[0009] The present invention further relates to a method for the detection of the flanking breakpoints of named chromosomal alterations by measurement of DNA copy number by quantitative PCR or DNA-Arrays and DNA sequencing.

[0010] A method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of DNA sequences flanking named genomic breakpoint or are located within such.

[0011] The present invention further relates to a method for the detection of chromosomal alterations characterized in that the copy number of one or more genomic nucleic acid sequences located within an altered chromosomal region (s) is detected by quantitative PCR techniques (e.g. TaqMan™, Lightcycler™ and Cycler™).

[0012] The present invention further relates to a method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least 2 markers whereby the markers are genes and fragments thereof or genomic nucleic acid sequences that are located on one chromosomal region which is altered in malignant neoplasia and breast cancer in particular.

[0013] The present invention also discloses a method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least 2 markers whereby the markers are located on one or more chromosomal region(s) which is/are altered in malignant neoplasia; and the markers interact as (i) receptor and ligand or (ii) members of the same signal transduction pathway or (iii) members of synergistic signal transduction pathways or (iv) members of antagonistic signal transduction pathways or (v) transcription factor and transcription factor binding site.

[0014] Also disclosed is a method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least one marker whereby the marker is a VNTR, SNP, RFLP or STS which is located on one chromosomal region which is altered in malignant neoplasia due to amplification and the marker is detected in (a) a cancerous and (b) a non cancerous tissue or biological sample from the same individual. A preferred embodiment is the detection of at least one VNTR marker of Table 6 or at least on SNP marker of Table 4 or combinations thereof. Even more preferred can the detection, quantification and sizing of such polymorphic markers be achieved by methods of (a) for the comparative measurement of amount and size by PCR amplification and subsequent capillary electrophoresis, (b) for sequence determination and allelic discrimination by gel electrophoresis (e.g. SSCP, DGGE), real time kinetic PCR, direct DNA sequencing, pyro-sequencing, mass-specific allelic discrimination or resequencing by DNA array technologies, (c) for the determination of specific restriction patterns and subsequent electrophoretic separation and (d) for allelic discrimination by allele specific PCR (e.g. ASO). An even more favorable detection of a heterozygous VNTR, SNP, RFLP or STS is done in a multiplex fashion, utilizing a variety of labeled primers (e.g. fluorescent, radioactive, bioactive) and a suitable capillary electrophoresis (CE) detection system.

[0015] In another embodiment the expression of these genes can be detected with DNA-arrays as described in WO9727317 and US6379895.

[0016] In a further embodiment the expression of these genes can be detected with bead based direct fluorescent readout techniques such as described in WO9714028 and WO9952708.

[0017] In one embodiment, the invention pertains to a method of determining the phenotype of a cell or tissue, comprising detecting the differential expression, relative to a normal or untreated cell, of at least one polynucleotide comprising SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19 or 21 to 26 or 53 to 75, wherein the polynucleotide is differentially expressed by at least about 1.5 fold, at least about 2 fold or at least about 3 fold.

[0018] In a further aspect the invention pertains to a method of determining the phenotype of a cell or tissue, comprising detecting the differential expression, relative to a normal or untreated cell, of at least one polynucleotide which hybridizes under stringent conditions to one of the polynucleotides of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19 or 21 to 26 or 53 to 75 and encodes a polypeptide exhibiting the same biological function as given in Table 2 or 3 for the respective polynucleotide, wherein the polynucleotide is differentially expressed by at least about 1.5 fold, at least about 2 fold or at least about 3 fold.

[0019] In another embodiment of the invention a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19 or 21 to 26 and 53 to 75 or encoding one of the polypeptides with SEQ ID NO: 28 to 32, 34, 35, 37 to 42, 44, 45 or 47 to 52 or 76 to 98 can be used to identify cells or tissue in individuals which exhibit a phenotype predisposed to breast cancer or a diseased phenotype, thereby (a) predicting whether an individual is at risk for the development, or (b) diagnosing whether an individual is having, or (c) prognosing the progression or the

outcome of the treatment malignant neoplasia and breast cancer in particular.

[0020] In yet another embodiment the invention provides a method for identifying genomic regions which are altered on the chromosomal level and encode genes that are linked by function and are differentially expressed in malignant neoplasia and breast cancer in particular.

[0021] In yet another embodiment the invention provides the genomic regions 17q12, 3p21 and 12q13 for use in prediction, diagnosis and prognosis as well as prevention and treatment of malignant neoplasia and breast cancer. In particular not only the intragenic regions, but also intergenic regions, pseudogenes or non-transcribed genes of said chromosomal regions can be used for diagnostic, predictive, prognostic and preventive and therapeutic compositions and methods.

[0022] In yet another embodiment the invention provides methods of screening for agents which regulate the activity of a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75. A test compound is contacted with a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75. Binding of the test compound to the polypeptide is detected. A test compound which binds to the polypeptide is thereby identified as a potential therapeutic agent for the treatment of malignant neoplasia and more particularly breast cancer.

[0023] In even another embodiment the invention provides another method of screening for agents which regulate the activity of a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75. A test compound is contacted with a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75. A biological activity mediated by the polypeptide is detected. A test compound which decreases the biological activity is thereby identified as a potential therapeutic agent for decreasing the activity of the polypeptide encoded by a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 in malignant neoplasia and breast cancer in particular. A test compound which increases the biological activity is thereby identified as a potential therapeutic agent for increasing the activity of the polypeptide encoded by a polypeptide selected from one of the polypeptides with SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 in malignant neoplasia and breast cancer in particular.

[0024] In another embodiment the invention provides a method of screening for agents which regulate the activity of a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75. A test compound is contacted with a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75. Binding of the test compound to the polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 is detected. A test compound which binds to the polynucleotide is thereby identified as a potential therapeutic agent for regulating the activity of a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 in malignant neoplasia and breast cancer in particular.

[0025] The invention thus provides polypeptides selected from one of the polypeptides with SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 which can be used to identify compounds which may act, for example, as regulators or modulators such as agonists and antagonists, partial agonists, inverse agonists, activators, co-activators and inhibitors of the polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75. Accordingly, the invention provides reagents and methods for regulating a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 in malignant neoplasia and more particularly breast cancer. The regulation can be an up-or-down regulation. Reagents that modulate the expression, stability or amount of a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or the activity of the polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 can be a protein, a peptide, a peptidomimetic, a nucleic acid, a nucleic acid analogue (e.g. peptide nucleic acid, locked nucleic acid) or a small molecule. Methods that modulate the expression, stability or amount of a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or the activity of the polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 can be gene replacement therapies, antisense, ribozyme and triplex nucleic acid approaches.

[0026] In one embodiment of the invention provides antibodies which specifically bind to a full-length or partial polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 for use in prediction, prevention, diagnosis, prognosis and

treatment of malignant neoplasia and breast cancer in particular.

[0027] Yet another embodiment of the invention is the use of a reagent which specifically binds to a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 in the preparation of a medicament for the treatment of malignant neoplasia and breast cancer in particular.

[0028] Still another embodiment is the use of a reagent that modulates the activity or stability of a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or the expression, amount or stability of a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 in the preparation of a medicament for the treatment of malignant neoplasia and breast cancer in particular.

[0029] Still another embodiment of the invention is a pharmaceutical composition which includes a reagent which specifically binds to a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75, and a pharmaceutically acceptable carrier.

[0030] Yet another embodiment of the invention is a pharmaceutical composition including a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or encoding a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98.

[0031] In one embodiment, a reagent which alters the level of expression in a cell of a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or encoding a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98, or a sequence complementary thereto, is identified by providing a cell, treating the cell with a test reagent, determining the level of expression in the cell of a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or encoding a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or a sequence complementary thereto, and comparing the level of expression of the polynucleotide in the treated cell with the level of expression of the polynucleotide in an untreated cell, wherein a change in the level of expression of the polynucleotide in the treated cell relative to the level of expression of the polynucleotide in the untreated cell is indicative of an agent which alters the level of expression of the polynucleotide in a cell.

[0032] The invention further provides a pharmaceutical composition comprising a reagent identified by this method.

[0033] Another embodiment of the invention is a pharmaceutical composition which includes a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or which is encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75.

[0034] A further embodiment of the invention is a pharmaceutical composition comprising a polynucleotide including a sequence which hybridizes under stringent conditions to a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 and encoding a polypeptide exhibiting the same biological function as given for the respective polynucleotide in Table 2 or 3, or encoding a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98. Pharmaceutical compositions, useful in the present invention may further include fusion proteins comprising a polypeptide comprising a polynucleotide selected from SEQ ID NO: 27 to 52 and 76 to 98, or a fragment thereof, antibodies, or antibody fragments

BRIEF DESCRIPTION OF THE DRAWINGS

[0035]

Fig. 1 shows a sketch of the chromosome 17 with G-banding pattern and cytogenetic positions. In the blow out at the lower part of the figure a detailed view of the chromosomal area of the long arm of chromosome 17 (17q 12-21.1) is provided. Each vertical rectangle depicted in medium gray, represents a gene as labeled below or above the individual position. The order of genes depicted in this graph has been deduced from experiments questioning the amplification an over expression and from public available data (e.g. UCSC, NCBI or Ensembl).

Fig. 2 shows the same region as depicted before in Fig. 1 and a cluster representation of the individual expression values measured by DNA-chip hybridization. The gene representing squares are indicated by a dotted line. In the upper part of the cluster representation 4 tumor cell lines, of which two harbor a known HER-2/neu over expression (SKBR3 and AU565), are depicted with their individual expression profiles. Not only the HER-2/neu gene shows a clear over expression but as provided by this invention several other genes with in the surrounding. In the middle part of the cluster representation expression data obtained from immune histochemically characterized tumor samples are presented. Two of the depicted probes show a significant over

expression of genes marked by the white rectangles. For additional information and comparison expression profiles of several non diseased human tissues (RNAs obtained from Clontech Inc.) are provided. Closest relation to the expression profile of HER-2/neu positive tumors displays human brain and neural tissue.

Fig. 3 provides data from DNA amplification measurements by qPCR (e.g. TaqMan). Data indicates that in several analyzed breast cancer cell lines harbor amplification of genes which were located in the previously described region (ARCHEON). Data were displayed for each gene on the x-axis and 40-Ct at the y-axis. Data were normalized to the expression level of GAPDH as seen in the first group of columns.

Fig. 4 represents a graphical overview on the amplified regions and provides information on the length of the individual amplification and over expression in the analyzed tumor cell lines. The length of the amplification and the composition of genes has a significant impact on the nature of the cancer cell and on the responsiveness on certain drugs, as described elsewhere.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

[0036] "Differential expression", as used herein, refers to both quantitative as well as qualitative differences in the genes' expression patterns depending on differential development and/or tumor growth. Differentially expressed genes may represent "marker genes," and/or "target genes". The expression pattern of a differentially expressed gene disclosed herein may be utilized as part of a prognostic or diagnostic breast cancer evaluation. Alternatively, a differentially expressed gene disclosed herein may be used in methods for identifying reagents and compounds and uses of these reagents and compounds for the treatment of breast cancer as well as methods of treatment.

[0037] "Biological activity" or "bioactivity" or "activity" or "biological function", which are used interchangeably, herein mean an effector or antigenic function that is directly or indirectly performed by a polypeptide (whether in its native or denatured conformation), or by any fragment thereof *in vivo* or *in vitro*. Biological activities include but are not limited to binding to polypeptides, binding to other proteins or molecules, enzymatic activity, signal transduction, activity as a DNA binding protein, as a transcription regulator, ability to bind damaged DNA, etc. A bioactivity can be modulated by directly affecting the subject polypeptide. Alternatively, a bioactivity can be altered by modulating the level of the polypeptide, such as by modulating expression of the corresponding gene.

[0038] The term "marker" or "biomarker" refers a biological molecule, e.g., a nucleic acid, peptide, hormone, etc., whose presence or concentration can be detected and correlated with a known condition, such as a disease state.

[0039] "Marker gene," as used herein, refers to a differentially expressed gene which expression pattern may be utilized as part of predictive, prognostic or diagnostic malignant neoplasia or breast cancer evaluation, or which, alternatively, may be used in methods for identifying compounds useful for the treatment or prevention of malignant neoplasia and breast cancer in particular. A marker gene may also have the characteristics of a target gene.

[0040] "Target gene", as used herein, refers to a differentially expressed gene involved in breast cancer in a manner by which modulation of the level of target gene expression or of target gene product activity may act to ameliorate symptoms of malignant neoplasia and breast cancer in particular. A target gene may also have the characteristics of a marker gene.

[0041] The term "biological sample", as used herein, refers to a sample obtained from an organism or from components (e.g., cells) of an organism. The sample may be of any biological tissue or fluid. Frequently the sample will be a "clinical sample" which is a sample derived from a patient. Such samples include, but are not limited to, sputum, blood, blood cells (e.g., white cells), tissue or fine needle biopsy samples, cell-containing bodyfluids, free floating nucleic acids, urine, peritoneal fluid, and pleural fluid, or cells therefrom. Biological samples may also include sections of tissues such as frozen sections taken for histological purposes.

[0042] By "array" or "matrix" is meant an arrangement of addressable locations or "addresses" on a device. The locations can be arranged in two dimensional arrays, three dimensional arrays, or other matrix formats. The number of locations can range from several to at least hundreds of thousands. Most importantly, each location represents a totally independent reaction site. Arrays include but are not limited to nucleic acid arrays, protein arrays and antibody arrays. A "nucleic acid array" refers to an array containing nucleic acid probes, such as oligonucleotides, polynucleotides or larger portions of genes. The nucleic acid on the array is preferably single stranded. Arrays wherein the probes are oligonucleotides are referred to as "oligonucleotide arrays" or "oligonucleotide chips." A "microarray," herein also refers to a "biochip" or "biological chip", an array of regions having a density of discrete regions of at least about 100/cm², and preferably at least about 1000/cm². The regions in a microarray have typical dimensions, e.g., diameters, in the range of between about 10-250 µm, and are separated from other regions in the array by about the same distance. A "protein array" refers to an array containing polypeptide probes or protein probes which can be in native form or

denatured. An "antibody array" refers to an array containing antibodies which include but are not limited to monoclonal antibodies (e.g. from a mouse), chimeric antibodies, humanized antibodies or phage antibodies and single chain antibodies as well as fragments from antibodies.

[0043] The term "agonist", as used herein, is meant to refer to an agent that mimics or upregulates (e.g., potentiates or supplements) the bioactivity of a protein. An agonist can be a wild-type protein or derivative thereof having at least one bioactivity of the wild-type protein. An agonist can also be a compound that upregulates expression of a gene or which increases at least one bioactivity of a protein. An agonist can also be a compound which increases the interaction of a polypeptide with another molecule, e.g., a target peptide or nucleic acid.

[0044] The term "antagonist" as used herein is meant to refer to an agent that downregulates (e.g., suppresses or inhibits) at least one bioactivity of a protein. An antagonist can be a compound which inhibits or decreases the interaction between a protein and another molecule, e.g., a target peptide, a ligand or an enzyme substrate. An antagonist can also be a compound that downregulates expression of a gene or which reduces the amount of expressed protein present.

[0045] "Small molecule" as used herein, is meant to refer to a composition, which has a molecular weight of less than about 5 kD and most preferably less than about 4 kD. Small molecules can be nucleic acids, peptides, polypeptides, peptidomimetics, carbohydrates, lipids or other organic (carbon-containing) or inorganic molecules. Many pharmaceutical companies have extensive libraries of chemical and/or biological mixtures, often fungal, bacterial, or algal extracts, which can be screened with any of the assays of the invention to identify compounds that modulate a bioactivity.

[0046] The terms "modulated" or "modulation" or "regulated" or "regulation" and "differentially regulated" as used herein refer to both upregulation (i.e., activation or stimulation (e.g., by agonizing or potentiating) and down regulation (i.e., inhibition or suppression (e.g., by antagonizing, decreasing or inhibiting)).

[0047] "Transcriptional regulatory unit" refers to DNA sequences, such as initiation signals, enhancers, and promoters, which induce or control transcription of protein coding sequences with which they are operably linked. In preferred embodiments, transcription of one of the genes is under the control of a promoter sequence (or other transcriptional regulatory sequence) which controls the expression of the recombinant gene in a cell-type in which expression is intended. It will also be understood that the recombinant gene can be under the control of transcriptional regulatory sequences which are the same or which are different from those sequences which control transcription of the naturally occurring forms of the polypeptide.

[0048] The term "derivative" refers to the chemical modification of a polypeptide sequence, or a polynucleotide sequence. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

[0049] The term "nucleotide analog" refers to oligomers or polymers being at least in one feature different from naturally occurring nucleotides, oligonucleotides or polynucleotides, but exhibiting functional features of the respective naturally occurring nucleotides (e.g. base pairing, hybridization, coding information) and that can be used for said compositions. The nucleotide analogs can consist of non-naturally occurring bases or polymer backbones, examples of which are LNAs, PNAs and Morpholinos. The nucleotide analog has at least one molecule different from its naturally occurring counterpart or equivalent.

[0050] "BREAST CANCER GENES" or "BREAST CANCER GENE" as used herein refers to the polynucleotides of SEQ ID NO: 1 to 26 and 53 to 75, as well as derivatives, fragments, analogs and homologues thereof, the polypeptides encoded thereby, the polypeptides of SEQ ID NO: 27 to 52 and 76 to 98 as well as derivatives, fragments, analogs and homologues thereof and the corresponding genomic transcription units which can be derived or identified with standard techniques well known in the art using the information disclosed in Tables 1 to 5 and Figures 1 to 4. The GenBank, Locuslink ID and the UniGene accession numbers of the polynucleotide sequences of the SEQ ID NO: 1 to 26 and 53 to 75 and the polypeptides of the SEQ ID NO: 27 to 52 and 76 to 98 are shown in Table 1, the gene description, gene function and subcellular localization is given in Tables 2 and 3.

[0051] The term "chromosomal region" as used herein refers to a consecutive DNA stretch on a chromosome which can be defined by cytogenetic or other genetic markers such as e.g. restriction length polymorphisms (RFLPs), single nucleotide polymorphisms (SNPs), expressed sequence tags (ESTs), sequence tagged sites (STSs), micro-satellites, variable number of tandem repeats (VNTRs) and genes. Typically a chromosomal region consists of up to 2 Megabases (MB), up to 4 MB, up to 6 MB, up to 8 MB, up to 10 MB, up to 20 MB or even more MB.

[0052] The term "altered chromosomal region" or "aberrant chromosomal region" refers to a structural change of the chromosomal composition and DNA sequence, which can occur by the following events: amplifications, deletions, inversions, insertions, translocations and/or viral integrations. A trisomy, where a given cell harbors more than two copies of a chromosome, is within the meaning of the term "amplification" of a chromosome or chromosomal region.

[0053] The present invention provides polynucleotide sequences and proteins encoded thereby, as well as probes

derived from the polynucleotide sequences, antibodies directed to the encoded proteins, and predictive, preventive, diagnostic, prognostic and therapeutic uses for individuals which are at risk for or which have malignant neoplasia and breast cancer in particular. The sequences disclosure herein have been found to be differentially expressed in samples from breast cancer.

[0054] The present invention is based on the identification of 43 genes that are differentially regulated (up- or down-regulated) in tumor biopsies of patients with clinical evidence of breast cancer. The identification of 43 human genes which were not known to be differentially regulated in breast cancer states and their significance for the disease is described in the working examples herein. The characterization of the co-expression of these genes provides newly identified roles in breast cancer. The gene names, the database accession numbers (GenBank and UniGene) as well as the putative or known functions of the encoded proteins and their subcellular localization are given in Tables 1 to 4. The primer sequences used for the gene amplification are shown in Table 5.

[0055] In either situation, detecting expression of these genes in excess or in with lower level as compared to normal expression provides the basis for the diagnosis of malignant neoplasia and breast cancer. Furthermore, in testing the efficacy of compounds during clinical trials, a decrease in the level of the expression of these genes corresponds to a return from a disease condition to a normal state, and thereby indicates a positive effect of the compound.

[0056] Another aspect of the present invention is based on the observation that neighboring genes within defined genomic regions functionally interact and influence each others function directly or indirectly. A genomic region encoding functionally interacting genes that are co-amplified and co-expressed in neoplastic lesions has been defined as an "ARCHEON". (ARCHEON = Altered Region of Changed Chromosomal Expression Observed in Neoplasms). Chromosomal alterations often affect more than one gene. This is true for amplifications, duplications, insertions, integrations, inversions, translocations, and deletions. These changes can have influence on the expression level of single or multiple genes. Most commonly in the field of cancer diagnostics and treatment the changes of expression levels have been investigated for single, putative relevant target genes such as MLV12 (5p14), NRASL3 (6p12), EGFR (7p12), c-myc (8q23), Cyclin D1 (11q13), IGF1R (15q25), HER-2/neu (17q12), PCNA (20q12). However, the altered expression level and interaction of multiple (i.e. more than two) genes within one genomic region with each other has not been addressed. Genes of an ARCHEON form gene clusters with tissue specific expression patterns. The mode of interaction of individual genes within such a gene cluster suspected to represent an ARCHEON can be either protein-protein or protein-nucleic acid interaction, which may be illustrated but not limited by the following examples: ARCHEON gene interaction may be in the same signal transduction pathway, may be receptor to ligand binding, receptor kinase and SH2 or SH3 binding, transcription factor to promoter binding, nuclear hormone receptor to transcription factor binding, phosphogroup donation (e.g. kinases) and acceptance (e.g. phosphoprotein), mRNA stabilizing protein binding and transcriptional processes. The individual activity and specificity of a pair genes and or the proteins encoded thereby or of a group of such in a higher order, may be readily deduced from literature, published or deposited within public databases by the skilled person. However in the context of an ARCHEON the interaction of members being part of an ARCHEON will potentiate, exaggerate or reduce their singular functions. This interaction is of importance in defined normal tissues in which they are normally co-expressed. Therefore, these clusters have been commonly conserved during evolution. The aberrant expression of members of these ARCHEON in neoplastic lesions, however, (especially within tissues in which they are normally not expressed) has influence on tumor characteristics such as growth, invasiveness and drug responsiveness. Due to the interaction of these neighboring genes it is of importance to determine the members of the ARCHEON which are involved in the deregulation events. In this regard amplification and deletion events in neoplastic lesions are of special interest.

[0057] The invention relates to a method for the detection of chromosomal alterations by (a) determining the relative mRNA abundance of individual mRNA species or (b) determining the copy number of one or more chromosomal region (s) by quantitative PCR. In one embodiment information on the genomic organization and spatial regulation of chromosomal regions is assessed by bioinformatic analysis of the sequence information of the human genome (UCSC, NCBI) and then combined with RNA expression data from GeneChip™ DNA-Arrays (Affymetrix) and/or quantitative PCR (TaqMan) from RNA-samples or genomic DNA.

[0058] In a further embodiment the functional relationship of genes located on a chromosomal region which is altered (amplified or deleted) is established. The altered chromosomal region is defined as an ARCHEON if genes located on that region functionally interact.

[0059] The 17q 12 locus was investigated as one model system, harboring the HER-2/neu gene. By establishing a high-resolution assay to detect amplification events in neighboring genes, 43 genes that are commonly co-amplified in breast cancer cell lines and patient samples were identified. By gene array technologies and immunological methods their co-overexpression in tumor samples was demonstrated. Surprisingly, by clustering tissue samples with HER-2/neu positive Tumor samples, it was found that the expression pattern of this larger genomic region (consisting of 43 genes) is very similar to control brain tissue. HER-2/neu negative breast tumor tissue did not show a similar expression pattern. Indeed, some of the genes within these cluster are important for neural development (HER-2/neu, THRA) in mouse model systems or are described to be expressed in neural cells (NeuroD2). Moreover, by searching similar

gene combinations in the human and rodent genome additional homologous chromosomal regions on chromosome 3p21 and 12q13 harboring several isoforms of the respective genes (see below) were found. There was a strong evidence for multiple interactions between the 43 candidate genes, as being part of identical pathways (HER-2, neu, GRB7, CrkRS, CDC6), influencing the expression of each other (HER-2/neu, THRA, RARA), interacting with each other (PPARGBP, THRA, RARA, NR1D1 or HER-2/neu, GRB7) or expressed in defined tissues (CACNB1, PPARGBP, etc.). Interestingly, the genomic regions of the ARCHEONs that were identified are amplified in acquired Tamoxifen resistance of HER-2/neu negative cells (MCF7), which are normally sensitive to Tamoxifen treatment [Achuthan et al., 2001,(2)].

[0060] Moreover, altered responsiveness to treatment due to the alterations of the genes within these ARCHEONs was observed. Surprisingly, genes within the ARCHEONs are of importance even in the absence of HER-2/neu homologues. Some of the genes within the ARCHEONs, do not only serve as marker genes for prognostic purposes, but have already been known as targets for therapeutic intervention. For example TOP2 alpha is a target of anthracyclins. THRA and RARA can be targeted by hormones and hormone analogs (e.g. T3, rT3, RA). Due to their high affinity binding sites and available screening assays (reporter assays based on their transcriptional potential) the hormone receptors which are shown to be linked to neoplastic pathophysiology for the first time herein are ideal targets for drug screening and treatment of malignant neoplasia and breast cancer in particular. In this regard it is essential to know which members of the ARCHEON are altered in the neoplastic lesions. Particularly it is important to know the nature, number and extent to which the ARCHEON genes are amplified or deleted. The ARCHEONs are flanked by similar, endogenous retroviruses (e.g. HERV-K= "human endogenous retrovirus"), some of which are activated in breast cancer. These viruses may have also been involved in the evolutionary duplication of the ARCHEONs.

[0061] The analysis of the 17q12 region proved data obtained by IHC and identified several additional genes being co-amplified with the HER-2/neu gene. Comparative Analysis of RNA-based quantitative RT-PCR (TaqMan) with DNA-based qPCR from tumor cell lines identified the same amplified region. Genes at the 17q11.2-21. region are offered by way of illustration not by way of limitation. A graphical display of the described chromosomal region is provided in Figure 1.

Biological relevance of the genes which are part of the 17q12 ARCHEON

MLN50

[0062] By differential screening of cDNAs from breast cancer-derived metastatic axillary lymph nodes, TRAF4 and 3 other novel genes (MLN51, MLN62, MLN64) were identified that are overexpressed in breast cancer [Tomasetto et al., 1995, (3)]. One gene, which they designated MLN50, was mapped to 17q11.2-q21.3 by radioactive in situ hybridization. In breast cancer cell lines, overexpression of the 4 kb MLN50 mRNA was correlated with amplification of the gene and with amplification and overexpression of ERBB2, which maps to the same region. The authors suggested that the 2 genes belong to the same amplicon. Amplification of chromosomal region 17q11-q21 is one of the most common events occurring in human breast cancers. They reported that the predicted 261-amino acid MLN50 protein contains an N-terminal LIM domain and a C-terminal SH3 domain. They renamed the protein LASP1, for 'LIM and SH3 protein.' Northern blot analysis revealed that LASP1 mRNA was expressed at a basal level in all normal tissues examined and overexpressed in 8% of primary breast cancers. In most of these cancers, LASP1 and ERBB2 were simultaneously overexpressed.

MLLT6

[0063] The MLLT6 (AF17) gene encodes a protein of 1,093 amino acids, containing a leucine-zipper dimerization motif located 3-prime of the fusion point and a cysteine-rich domain at the end terminus. AF17 was found to contain stretches of amino acids previously associated with domains involved in transcriptional repression or activation.

[0064] Chromosome translocations involving band 1 q23 are associated with approximately 10% of patients with acute lymphoblastic leukemia (ALL) and more than 5% of patients with acute myeloid leukemia (AML). The gene at 11q23 involved in the translocations is variously designated ALL1, HRX, MLL, and TAX1. The partner gene in one of the rarer translocations, t(11;17)(q23;q21), designated MLLT6 on 17q12.

ZNFI44 (MEL18)

[0065] MEL18 cDNA encodes a novel cys-rich zinc finger motif. The gene is expressed strongly in most tumor cell lines, but its normal tissue expression was limited to cells of neural origin and was especially abundant in fetal neural cells. It belongs to a RING-finger motif family which includes BMI1. The MEL18/BMI1 gene family represents a mammalian homolog of the Drosophila 'polycomb' gene group, thereby belonging to a memory mechanism involved in

maintaining the the expression pattern of key regulatory factors such as Hox genes. Bmi1, Mel18 and M33 genes, as representative examples of mouse Pc-G genes. Common phenotypes observed in knockout mice mutant for each of these genes indicate an important role for Pc-G genes not only in regulation of Hox gene expression and axial skeleton development but also in control of proliferation and survival of haematopoietic cell lineages. This is in line with the observed proliferative deregulation observed in lymphoblastic leukemia. The MEL18 gene is conserved among vertebrates. Its mRNA is expressed at high levels in placenta, lung, and kidney, and at lower levels in liver, pancreas, and skeletal muscle. Interestingly, cervical and lumbo-sacral-HOX gene expression is altered in several primary breast cancers with respect to normal breast tissue with the HoxB gene cluster being present on 17q distal to the 17q12 locus. Moreover, delay of differentiation with persistent nests of proliferating cells was found in endothelial cells cocultured with HOXB7-transduced SkBr3 cells, which exhibit a 17q12 amplification. Tumorigenicity of these cells has been evaluated in vivo. Xenograft in athymic nude mice showed that SkBr3/HOXB7 cells developed tumors with an increased number of blood vessels, either irradiated or not, whereas parental SkBr3 cells did not show any tumor take unless mice were sublethally irradiated. As part of this invention, we have found MEL18 to be overexpressed specifically in tumors bearing Her-2/neu gene amplification, which can be critical for Hox expression.

PHOSPHATIDYLINOSITOL-4-PHOSPHATE 5-KINASE, TYPE II, BETA; PIP5K2B

[0066] Phosphoinositide kinases play central roles in signal transduction. Phosphatidylinositol-4-phosphate 5-kinases (PIP5Ks) phosphorylate phosphatidylinositol 4-phosphate, giving rise to phosphatidylinositol 4,5-bisphosphate. The PIP5K enzymes exist as multiple isoforms that have various immunoreactivities, kinetic properties, and molecular masses. They are unique in that they possess almost no homology to the kinase motifs present in other phosphatidylinositol, protein, and lipid kinases. By screening a human fetal brain cDNA library with the PIP5K2B EST the full length gene could be isolated. The deduced 416-amino acid protein is 78% identical to PIP5K2A. Using SDS-PAGE, the authors estimated that bacterially expressed PIP5K2B has a molecular mass of 47 kD. Northern blot analysis detected a 6.3-kb PIP5K2B transcript which was abundantly expressed in several human tissues. PIP5K2B interacts specifically with the juxtamembrane region of the p55 TNF receptor (TNFR1) and PIP5K2B activity is increased in mammalian cells by treatment with TNF-alpha. A modeled complex with membrane-bound substrate and ATP shows how a phosphoinositide kinase can phosphorylate its substrate in situ at the membrane interface. The substrate-binding site is open on 1 side, consistent with dual specificity for phosphatidylinositol 3- and 5-phosphates. Although the amino acid sequence of PIP5K2A does not show homology to known kinases, recombinant PIP5K2A exhibited kinase activity. PIP5K2A contains a putative Src homology 3 (SH3) domain-binding sequence. Overexpression of mouse PIP5K1B in COS7 cells induced an increase in short actin fibers and a decrease in actin stress fibers.

TEM7

[0067] Using serial analysis of gene expression (SAGE) a partial cDNAs corresponding to several tumor endothelial markers (TEMs) that displayed elevated expression during tumor angiogenesis could be identified. Among the genes identified was TEM7. Using database searches and 5-prime RACE the entire TEM7 coding region, which encodes a 500-amino acid type I transmembrane protein, has been described. The extracellular region of TEM7 contains a plexin-like domain and has weak homology to the ECM protein nidogen. The function of these domains, which are usually found in secreted and extracellular matrix molecules, is unknown. Nidogen itself belongs to the entactin protein family and helps to determine pathways of migrating axons by switching from circumferential to longitudinal migration. Entactin is involved in cell migration, as it promotes trophoblast outgrowth through a mechanism mediated by the RGD recognition site, and plays an important role during invasion of the endometrial basement membrane at implantation. As entactin promotes thymocyte adhesion but affects thymocyte migration only marginally, it is suggested that entactin may play a role in thymocyte localization during T cell development.

[0068] In situ hybridization analysis of human colorectal cancer demonstrated that TEM7 was expressed clearly in the endothelial cells of the tumor stroma but not in the endothelial cells of normal colonic tissue. Using in situ hybridization to assay expression in various normal adult mouse tissues, they observed that TEM7 was largely undetectable in mouse tissues or tumors, but was abundantly expressed in mouse brain.

ZNFN1A3

[0069] By screening a B-cell cDNA library with a mouse Aiolos N-terminal cDNA probe, a cDNA encoding human Aiolos, or ZNFN1A3, was obtained. The deduced 509-amino acid protein, which is 86% identical to its mouse counterpart, has 4 DNA-binding zinc fingers in its N terminus and 2 zinc fingers that mediate protein dimerization in its C terminus. These domains are 100% and 96% homologous to the corresponding domains in the mouse protein, respectively. Northern blot analysis revealed strong expression of a major 11.0- and a minor 4.4-kb ZNFN1A3 transcript in

peripheral blood leukocytes, spleen, and thymus, with lower expression in liver, small intestine, and lung.

[0070] Ikaros (ZNFN1A1), a hemopoietic zinc finger DNA-binding protein, is a central regulator of lymphoid differentiation and is implicated in leukemogenesis. The execution of normal function of Ikaros requires sequence-specific DNA binding, transactivation, and dimerization domains. Mice with a mutation in a related zinc finger protein, Aiolos, are prone to B-cell lymphoma. In chemically induced murine lymphomas allelic losses on markers surrounding the Znf1a1 gene were detected in 27% of the tumors analyzed. Moreover specific Ikaros expression was in primary mouse hormone-producing anterior pituitary cells and substantial for Fibroblast growth factor receptor 4 (FGFR4) expression, which itself is implicated in a multitude of endocrine cell hormonal and proliferative properties with FGFR4 being differentially expressed in normal and neoplastic pituitary. Moreover Ikaros binds to chromatin remodelling complexes containing SWI/SNF proteins, which antagonize Polycomb function. Interestingly at the telomeric end of the disclosed ARCHEON the SWI/SNF complex member SMARCE1 (= SWI/SNF-related, matrix-associated, actin-dependent regulators of chromatin) is located and part of the described amplification. Due to the related binding specificities of Ikaros and Palindrom Binding Protein (PBP) it is suggestive, that ZNFN1A3 is able to regulate the Her-2/neu enhancer.

PPP1R1B

[0071] Midbrain dopaminergic neurons play a critical role in multiple brain functions, and abnormal signaling through dopaminergic pathways has been implicated in several major neurologic and psychiatric disorders. One well-studied target for the actions of dopamine is DARPP32. In the densely dopamine- and glutamate-innervated rat caudate-putamen, DARPP32 is expressed in medium-sized spiny neurons that also express dopamine D1 receptors. The function of DARPP32 seems to be regulated by receptor stimulation. Both dopaminergic and glutamatergic (NMDA) receptor stimulation regulate the extent of DARPP32 phosphorylation, but in opposite directions.

[0072] The human DARPP32 was isolated from a striatal cDNA library. The 204-amino acid DARPP32 protein shares 88% and 85% sequence identity, respectively, with bovine and rat DARPP32 proteins. The DARPP32 sequence is particularly conserved through the N terminus, which represents the active portion of the protein. Northern blot analysis demonstrated that the 2.1-kb DARPP32 mRNA is more highly expressed in human caudate than in cortex. In situ hybridization to postmortem human brain showed a low level of DARPP32 expression in all neocortical layers, with the strongest hybridization in the superficial layers. CDK5 phosphorylated DARPP32 in vitro and in intact brain cells. Phospho-thr75 DARPP32 inhibits PKA in vitro by a competitive mechanism. Decreasing phospho-thr75 DARPP32 in striatal cells either by a CDK5-specific inhibitor or by using genetically altered mice resulted in increased dopamine-induced phosphorylation of PKA substrates and augmented peak voltage-gated calcium currents. Thus, DARPP32 is a bifunctional signal transduction molecule which, by distinct mechanisms, controls a serine/threonine kinase and a serine/threonine phosphatase.

[0073] DARPP32 and t-DARPP are overexpressed in gastric cancers. It's suggested that overexpression of these 2 proteins in gastric cancers may provide an important survival advantage to neoplastic cells. It could be demonstrated that Darpp32 is an obligate intermediate in progesterone-facilitated sexual receptivity in female rats and mice. The facilitative effect of progesterone on sexual receptivity in female rats was blocked by antisense oligonucleotides to Darpp32. Homozygous mice carrying a null mutation for the Darpp32 gene exhibited minimal levels of progesterone-facilitated sexual receptivity when compared to their wildtype littermates, and progesterone significantly increased hypothalamic cAMP levels and cAMP-dependent protein kinase activity.

CACNB1

[0074] In 1991 a cDNA clone encoding a protein with high homology to the beta subunit of the rabbit skeletal muscle dihydropyridine-sensitive calcium channel from a rat brain cDNA library [Pragnell et al., 1991, (4)]. This rat brain beta-subunit cDNA hybridized to a 3.4-kb message that was expressed in high levels in the cerebral hemispheres and hippocampus and much lower levels in cerebellum. The open reading frame encodes 597 amino acids with a predicted mass of 65,679 Da which is 82% homologous with the skeletal muscle beta subunit. The corresponding human beta-subunit gene was localized to chromosome 17 by analysis of somatic cell hybrids. The authors suggested that the encoded brain beta subunit, which has a primary structure highly similar to its isoform in skeletal muscle, may have a comparable role as an integral regulatory component of a neuronal calcium channel.

RPL19

[0075] The ribosome is the only organelle conserved between prokaryotes and eukaryotes. In eukaryotes, this organelle consists of a 60S large subunit and a 40S small subunit. The mammalian ribosome contains 4 species of RNA and approximately 80 different ribosomal proteins, most of which appear to be present in equimolar amounts. In mammalian cells, ribosomal proteins can account for up to 15% of the total cellular protein, and the expression of the different

ribosomal protein genes, which can account for up to 7 to 9% of the total cellular mRNAs, is coordinately regulated to meet the cell's varying requirements for protein synthesis. The mammalian ribosomal protein genes are members of multigene families, most of which are composed of multiple processed pseudogenes and a single functional intron-containing gene. The presence of multiple pseudogenes hampered the isolation and study of the functional ribosomal protein genes. By study of somatic cell hybrids, it has been elucidated that DNA sequences complementary to 6 mammalian ribosomal protein cDNAs could be assigned to chromosomes 5, 8, and 17. Ten fragments mapped to 3 chromosomes [Nakamichi et al., 1986, (5)]. These are probably a mixture of functional (expressed) genes and pseudogenes. One that maps to 5q23-q33 rescues Chinese hamster emeline-resistance mutations in interspecies hybrids and is therefore the transcriptionally active RPS14 gene. In 1989 a PCR-based strategy for the detection of intron-containing genes in the presence of multiple pseudogenes was described. This technique was used to identify the intron-containing PCR products of 7 human ribosomal protein genes and to map their chromosomal locations by hybridization to human/rodent somatic cell hybrids [Feo et al., 1992, (6)]. All 7 ribosomal protein genes were found to be on different chromosomes: RPL19 on 17p12-q11, RPL30 on 8; RPL35A on 18; RPL36A on 14; RPS6 on 9pter-p13; RPS11 on 19cen-qter; and RPS17 on 11pter-p13. These are also different sites from the chromosomal location of previously mapped ribosomal protein genes S14 on chromosome 5, S4 on Xq and Yp, and RP117A on 9q3-q34. By fluorescence in situ hybridization the position of the RPL19 gene was mapped to 17q11 [Davies et al., 1989, (7)].

PPARBP, PBP, CRSP1, CRSP200, TRIP2, TRAP220, RB18A, DRIP230

[0076] The thyroid hormone receptors (TRs) are hormone-dependent transcription factors that regulate expression of a variety of specific target genes. They must specifically interact with a number of proteins as they progress from their initial translation and nuclear translocation to heterodimerization with retinoid X receptors (RXRs), functional interactions with other transcription factors and the basic transcriptional apparatus, and eventually, degradation. To help elucidate the mechanisms that underlie the transcriptional effects and other potential functions of TRs, the yeast interaction trap, a version of the yeast 2-hybrid system, was used to identify proteins that specifically interact with the ligand-binding domain of rat TR-beta-1 (TRHB) [Lee et al., 1995, (8)]. The authors isolated HeLa cell cDNAs encoding several different TR-interacting proteins (TRIPs), including TRIP2. TRIP2 interacted with rat TRb only in the presence of thyroid hormone. It showed a ligand-independent interaction with RXR-alpha, but did not interact with the glucocorticoid receptor (NR3C1) under any condition. By immunoscreening a human B-lymphoma cell cDNA expression library with the anti-p53 monoclonal antibody Pab1801, PPARBP was identified, which was called RB18A for "recognized by Pab1801 monoclonal antibody" [Drane et al., 1997, (9)]. The predicted 1,566-amino acid RB 18A protein contains several potential nuclear localization signals, 13 potential N-glycosylation sites, and a high number of potential phosphorylation sites. Despite sharing common antigenic determinants with p53, RB18A does not show significant nucleotide or amino acid sequence similarity with p53. Whereas the calculated molecular mass of RB18A is 166 kD, the apparent mass of recombinant RB18A was 205 kD by SDS-PAGE analysis. The authors demonstrated that RB18A shares functional properties with p53, including DNA binding, p53 binding, and self-oligomerization. Furthermore, RB18A was able to activate the sequence-specific binding of p53 to DNA, which was induced through an unstable interaction between both proteins. Northern blot analysis of human tissues detected an 8.5-kb RB18A transcript in all tissues examined except kidney, with highest expression in heart. Moreover mouse Pparbp, which was called Pbp for "Ppar-binding protein," as a protein that interacts with the Ppar-gamma (PPARG) ligand-binding domain in a yeast 2-hybrid system was identified [Zhu et al., 1997, (10)]. The authors found that Pbp also binds to PPAR-alpha (PPARA), RAR-alpha (RARA), RXR, and TR-beta-1 in vitro. The binding of Pbp to these receptors increased in the presence of specific ligands. Deletion of the last 12 amino acids from the C terminus of PPAR-gamma resulted in the abolition of interaction between Pbp and PPAR-gamma. Pbp modestly increased the transcriptional activity of PPAR-gamma, and a truncated form of Pbp acted as a dominant-negative repressor, suggesting that Pbp is a genuine transcriptional co-activator for PPAR. The predicted 1,560-amino acid Pbp protein contains 2 LXXLL motifs, which are considered necessary and sufficient for the binding of several co-activators to nuclear receptors. Northern blot analysis detected Pbp expression in all mouse tissues examined, with higher levels in liver, kidney, lung, and testis. In situ hybridization showed that Pbp is expressed during mouse ontogeny, suggesting a possible role for Pbp in cellular proliferation and differentiation. In adult mouse, in situ hybridization detected Pbp expression in liver, bronchial epithelium in the lung, intestinal mucosa, kidney cortex, thymic cortex, splenic follicles, and seminiferous epithelium in testis. Later on PPARBP was identified, which was called TRAP220, from an immunopurified TR-alpha (THRA)-TRAP complex [Yuan et al., 1998, (11)]. The authors cloned Jurkat cell cDNAs encoding TRAP220. The predicted 1,581-amino acid TRAP220 protein contains LXXLL domains, which are found in other nuclear receptor-interacting proteins. TRAP220 is nearly identical to RB18A, with these proteins differing primarily by an extended N terminus on TRAP220. In the absence of TR-alpha, TRAP220 appears to reside in a single complex with other TRAPs. TRAP220 showed a direct ligand-dependent interaction with TR-alpha, which was mediated through the C terminus of TR-alpha and, at least in part, the LXXLL domains of TRAP220. TRAP220 also interacted with other nuclear receptors, including vitamin D receptor,

RARA, RXRA, PPARA, PPARG, and estrogen receptor-alpha (ESR1; 133430), in a ligand-dependent manner. TRAP220 moderately stimulated human TR-alpha-mediated transcription in transfected cells, whereas a fragment containing the LXXLL motifs acted as a dominant-negative inhibitor of nuclear receptor-mediated transcription both in transfected cells and in cell-free transcription systems. Further studies indicated that TRAP220 plays a major role in anchoring other TRAPs to TR-alpha during the function of the TR-alpha-TRAP complex and that TRAP220 may be a global co-activator for the nuclear receptor superfamily. PBP, a nuclear receptor co-activator, interacts with estrogen receptor-alpha (ESR1) in the absence of estrogen. This interaction was enhanced in the presence of estrogen, but was reduced in the presence of the anti-estrogen Tamoxifen. Transfection of PBP into cultured cells resulted in enhancement of estrogen-dependent transcription, indicating that PBP serves as a co-activator in estrogen receptor signaling. To examine whether overexpression of PBP plays a role in breast cancer because of its co-activator function in estrogen receptor signaling, the levels of PBP expression in breast tumors was determined [Zhu et al., 1999, (12)]. High levels of PBP expression were detected in approximately 50% of primary breast cancers and breast cancer cell lines by ribonuclease protection analysis, in situ hybridization, and immunoperoxidase staining. By using FISH, the authors mapped the PBP gene to 17q12, a region that is amplified in some breast cancers. They found PBP gene amplification in approximately 24% (6 of 25) of breast tumors and approximately 30% (2 of 6) of breast cancer cell lines, implying that PBP gene overexpression can occur independent of gene amplification. They determined that the PBP gene comprises 17 exons that together span more than 37 kb. Their findings, in particular PBP gene amplification, suggested that PBP, by its ability to function as an estrogen receptor-alpha co-activator, may play a role in mammary epithelial differentiation and in breast carcinogenesis.

NEUROD2

[0077] Basic helix-loop-helix (bHLH) proteins are transcription factors involved in determining cell type during development. In 1995 a bHLH protein was described, termed NeuroD (for 'neurogenic differentiation'), that functions during neurogenesis [Lee et al., 1995, (13)]. The human NEUROD gene maps to chromosome 2q32. The cloning and characterization of 2 additional NEUROD genes, NEUROD2 and NEUROD3 was described in 1996 [McCormick et al., 1996, (14)]. Sequences for the mouse and human homologues were presented. NEUROD2 shows a high degree of homology to the bHLH region of NEUROD, whereas NEUROD3 is more distantly related. The authors found that mouse neuroD2 was initially expressed at embryonic day 11, with persistent expression in the adult nervous system. Similar to neuroD, neuroD2 appears to mediate neuronal differentiation. The human NEUROD2 was mapped to 17q12 by fluorescence in situ hybridization and the mouse homologue to chromosome 11 [Tamimi et al., 1997, (15)].

TELETHONIN

[0078] Telethonin is a sarcomeric protein of 19 kD found exclusively in striated and cardiac muscle. It appears to be localized to the Z disc of adult skeletal muscle and cultured myocytes. Telethonin is a substrate of titin, which acts as a molecular 'ruler' for the assembly of the sarcomere by providing spatially defined binding sites for other sarcomeric proteins. After activation by phosphorylation and calcium/calmodulin binding, titin phosphorylates the C-terminal domain of telethonin in early differentiating myocytes. The telethonin gene has been mapped to 17q12, adjacent to the phenylethanolamine N-methyltransferase gene [Valle et al., 1997, (16)].

PENT, PNMT

[0079] Phenylethanolamine N-methyltransferase catalyzes the synthesis of epinephrine from norepinephrine, the last step of catecholamine biosynthesis. The cDNA clone was first isolated in 1998 for bovine adrenal medulla PNMT using mixed oligodeoxyribonucleotide probes whose synthesis was based on the partial amino acid sequence of tryptic peptides from the bovine enzyme [Kaneda et al., 1988, (17)]. Using a bovine cDNA as a probe, the authors screened a human pheochromocytoma cDNA library and isolated a cDNA clone with an insert of about 1.0 kb, which contained a complete coding region of the enzyme. Northern blot analysis of human pheochromocytoma polyadenylated RNA using this cDNA insert as the probe demonstrated a single RNA species of about 1,000 nucleotides, suggesting that this clone is a full-length cDNA. The nucleotide sequence showed that human PNMT has 282 amino acid residues with a predicted molecular weight of 30,853, including the initial methionine. The amino acid sequence was 88% homologous to that of bovine enzyme. The PNMT gene was found to consist of 3 exons and 2 introns spanning about 2,100 basepairs. It was demonstrated that in transgenic mice the gene is expressed in adrenal medulla and retina. A hybrid gene consisting of 2 kb of the PNMT 5'-prime-flanking region fused to the simian virus 40 early region also resulted in tumor antigen mRNA expression in adrenal glands and eyes; furthermore, immunocytochemistry showed that the tumor antigen was localized in nuclei of adrenal medullary cells and cells of the inner nuclear cell layer of the retina, both prominent sites of epinephrine synthesis. The results indicate that the enhancer(s) for appropriate expres-

sion of the gene in these cell types are in the 2-kb 5-prime-flanking region of the gene.

[0080] Kaneda et al., 1988 (17), assigned the human PNMT gene to chromosome 17 by Southern blot analysis of DNA from mouse-human somatic cell hybrids. In 1992 the localization was narrowed down to 17q21-q22 by linkage analysis using RFLPs related to the PNMT gene and several 17q DNA markers [Hoehe et al., 1992, (18)]. The findings are of interest in light of the description of a genetic locus associated with blood pressure regulation in the stroke-prone spontaneously hypertensive rat (SHR-SP) on rat chromosome 10 in a conserved linkage synteny group corresponding to human chromosome 17q22-q24. See essential hypertension.

MGC9753

[0081] This gene maps on chromosome 17, at 17q12 according to RefSeq. It is expressed at very high level. It is defined by cDNA clones and produces, by alternative splicing, 7 different transcripts can be obtained (SEQ ID NO:60 to 66 and 83 to 89, Table 1), altogether encoding 7 different protein isoforms. Of specific interest is the putatively secreted isoform g, encoded by a mRNA of 2.55 kb. It's premessenger covers 16.94 kb on the genome. It has a very long 3' UTR. The protein (226 aa, MW 24.6 kDa, pI 8.5) contains no Pfam motif. The MGC9753 gene produces, by alternative splicing, 7 types of transcripts, predicted to encode 7 distinct proteins. It contains 13 confirmed introns, 10 of which are alternative. Comparison to the genome sequence shows that 11 introns follow the consensual [gt-ag] rule, 1 is atypical with good support [tg_cg]. The six most abundant isoforms are designated by a) to i) and code for proteins as follows:

a) This mRNA is 3.03 kb long, its premessenger covers 16.95 kb on the genome. It has a very long 3' UTR. The protein (190 aa, MW 21.5 kDa, pI 7.2) contains no Pfam motif. It is predicted to localise in the endoplasmic reticulum.

c) This mRNA is 1.17 kb long, its premessenger covers 16.93 kb on the genome. It may be incomplete at the N terminus. The protein (368 aa, MW 41.5 kDa, pI 7.3) contains no Pfam motif.

d) This mRNA is 3.17 kb long, its premessenger covers 16.94 kb on the genome. It has a very long 3' UTR and 5p UTR. The protein (190 aa, MW 21.5 kDa, pI 7.2) contains no Pfam motif. It is predicted to localise in the endoplasmic reticulum.

g) This mRNA is 2.55 kb long, its premessenger covers 16.94 kb on the genome. It has a very long 3' UTR. The protein (226 aa, MW 24.6 kDa, pI 8.5) contains no Pfam motif. It is predicted to be secreted.

h) This mRNA is 2.68 kb long, its premessenger covers 16.94 kb on the genome. It has a very long 3' UTR. The protein (320 aa, MW 36.5 kDa, pI 6.8) contains no Pfam motif. It is predicted to localise in the endoplasmic reticulum.

i) This mRNA is 2.34 kb long, its premessenger covers 16.94 kb on the genome. It may be incomplete at the N terminus. It has a very long 3' UTR. The protein (217 aa, MW 24.4 kDa, pI 5.9) contains no Pfam motif.

[0082] The MGC9753 gene may be homologue to the CAB2 gene located on chromosome 17q12. The CAB2, a human homologue of the yeast COS16 required for the repair of DNA double-strand breaks was cloned. Autofluorescence analysis of cells transfected with its GFP fusion protein demonstrated that CAB2 translocates into vesicles, suggesting that overexpression of CAB2 may decrease intercellular Mn-(2+) by accumulating it in the vesicles, in the same way as yeast.

Her-2/neu, ERBB2, NGL, TKR1

[0083] The oncogene originally called NEU was derived from rat neuro/glioblastoma cell lines. It encodes a tumor antigen, p185, which is serologically related to EGFR, the epidermal growth factor receptor. EGFR maps to chromosome 7. In 1985 it was found, that the human homologue, which they designated NGL (to avoid confusion with neuraminidase, which is also symbolized NEU), maps to 17q12-q22 by in situ hybridization and to 17q21-qter in somatic cell hybrids [Yang-Feng et al., 1985, (19)]. Thus, the SRO is 17q21-q22. Moreover, in 1985 a potential cell surface receptor of the tyrosine kinase gene family was identified and characterized by cloning the gene [Coussens et al., 1985, (20)]. Its primary sequence is very similar to that of the human epidermal growth factor receptor. Because of the seemingly close relationship to the human EGF receptor, the authors called the gene HER2. By Southern blot analysis of somatic cell hybrid DNA and by in situ hybridization, the gene was assigned to 17q21-q22. This chromosomal location of the gene is coincident with the NEU oncogene, which suggests that the 2 genes may in fact be the same; indeed, sequencing indicates that they are identical. In 1988 a correlation between overexpression of NEU protein and the

large-cell, comedo growth type of ductal carcinoma was found [van de Vijver et al., 1988, (21)]. The authors found no correlation, however, with lymph-node status or tumor recurrence. The role of HER2/NEU in breast and ovarian cancer was described in 1989, which together account for one-third of all cancers in women and approximately one-quarter of cancer-related deaths in females [Slamon et al., 1989, (22)].

[0084] An ERBB-related gene that is distinct from the ERBB gene, called ERBB1 was found in 1985. ERBB2 was not amplified in vulva carcinoma cells with EGFR amplification and did not react with EGF receptor mRNA. About 30-fold amplification of ERBB2 was observed in a human adenocarcinoma of the salivary gland. By chromosome sorting combined with velocity sedimentation and Southern hybridization, the ERBB2 gene was assigned to chromosome 17 [Fukushige et al., 1986, (23)]. By hybridization to sorted chromosomes and to metaphase spreads with a genomic probe, they mapped the ERBB2 locus to 17q21. This is the chromosome 17 breakpoint in acute promyelocytic leukemia (APL). Furthermore, they observed amplification and elevated expression of the ERBB2 gene in a gastric cancer cell line. Antibodies against a synthetic peptide corresponding to 14 amino acid residues at the COOH-terminus of a protein deduced from the ERBB2 nucleotide sequence were raised in 1986. With these antibodies, the ERBB2 gene product from adenocarcinoma cells was precipitated and demonstrated to be a 185-kD glycoprotein with tyrosine kinase activity. A cDNA probe for ERBB2 and by in situ hybridization to APL cells with a 15;17 chromosome translocation located the gene to the proximal side of the breakpoint [Kaneko et al., 1987, (24)]. The authors suggested that both the gene and the breakpoint are located in band 17q21.1 and, further, that the ERBB2 gene is involved in the development of leukemia. In 1987 experiments indicated that NEU and HER2 are both the same as ERBB2 [Di Fiore et al., 1987, (25)]. The authors demonstrated that overexpression alone can convert the gene for a normal growth factor receptor, namely, ERBB2, into an oncogene. The ERBB2 to 17q11-q21 by in situ hybridization [Popescu et al., 1989, (26)]. By in situ hybridization to chromosomes derived from fibroblasts carrying a constitutional translocation between 15 and 17, they showed that the ERBB2 gene was relocated to the derivative chromosome 15; the gene can thus be localized to 17q12-q21.32. By family linkage studies using multiple DNA markers in the 17q12-q21 region the ERBB2 gene was placed on the genetic map of the region.

[0085] Interleukin-6 is a cytokine that was initially recognized as a regulator of immune and inflammatory responses, but also regulates the growth of many tumor cells, including prostate cancer. Overexpression of ERBB2 and ERBB3 has been implicated in the neoplastic transformation of prostate cancer. Treatment of a prostate cancer cell line with IL6 induced tyrosine phosphorylation of ERBB2 and ERBB3, but not ERBB1/EGFR. The ERBB2 forms a complex with the gp130 subunit of the IL6 receptor in an IL6-dependent manner. This association was important because the inhibition of ERBB2 activity resulted in abrogation of IL6-induced MAPK activation. Thus, ERBB2 is a critical component of IL6 signaling through the MAP kinase pathway [Qiu et al., 1998, (27)]. These findings showed how a cytokine receptor can diversify its signaling pathways by engaging with a growth factor receptor kinase.

[0086] Overexpression of ERBB2 confers Taxol resistance in breast cancers. Overexpression of ERBB2 inhibits Taxol-induced apoptosis [Yu et al., 1998, (28)]. Taxol activates CDC2 kinase in MDA-MB-435 breast cancer cells, leading to cell cycle arrest at the G2/M phase and, subsequently, apoptosis. A chemical inhibitor of CDC2 and a dominant-negative mutant of CDC2 blocked Taxol-induced apoptosis in these cells. Overexpression of ERBB2 in MDA-MB-435 cells by transfection transcriptionally upregulates CDKN1A which associates with CDC2, inhibits Taxol-mediated CDC2 activation, delays cell entrance to G2/M phase, and thereby inhibits Taxol-induced apoptosis. In CDKN1A antisense-transfected MDA-MB-435 cells or in p21-/- MEF cells, ERBB2 was unable to inhibit Taxol-induced apoptosis. Therefore, CDKN1A participates in the regulation of a G2/M checkpoint that contributes to resistance to Taxol-induced apoptosis in ERBB2-overexpressing breast cancer cells.

[0087] A secreted protein of approximately 68 kD was described, designated herstatin, as the product of an alternative ERBB2 transcript that retains intron 8 [Doherty et al., 1999, (29)]. This alternative transcript specifies 340 residues identical to subdomains I and II from the extracellular domain of p185ERBB2, followed by a unique C-terminal sequence of 79 amino acids encoded by intron 8. The recombinant product of the alternative transcript specifically bound to ERBB2-transfected cells and was chemically crosslinked to p185ERBB2, whereas the intron-encoded sequence alone also bound with high affinity to transfected cells and associated with p185 solubilized from cell extracts. The herstatin mRNA was expressed in normal human fetal kidney and liver, but was at reduced levels relative to p185ERBB2 mRNA in carcinoma cells that contained an amplified ERBB2 gene. Herstatin appears to be an inhibitor of p185ERBB2, because it disrupts dimers, reduces tyrosine phosphorylation of p185, and inhibits the anchorage-independent growth of transformed cells that overexpress ERBB2. The HER2 gene is amplified and HER2 is overexpressed in 25 to 30% of breast cancers, increasing the aggressiveness of the tumor. Finally, it was found that a recombinant monoclonal antibody against HER2 increased the clinical benefit of first-line chemotherapy in metastatic breast cancer that overexpresses HER2 [Slamon et al., 2001, (30)].

GRB7

[0088] Growth factor receptor tyrosine kinases (GF-RTKs) are involved in activating the cell cycle. Several substrates

of GF-RTKs contain Src-homology 2 (SH2) and SH3 domains. SH2 domain-containing proteins are a diverse group of molecules important in tyrosine kinase signaling. Using the CORT (cloning of receptor targets) method to screen a high expression mouse library, the gene for murine Grb7, which encodes a protein of 535 amino acids, was isolated [Margolis et al., 1992, (31)]. GRB7 is homologous to ras-GAP (ras-GTPase-activating protein). It contains an SH2 domain and is highly expressed in liver and kidney. This gene defines the GRB7 family, whose members include the mouse gene Grb 10 and the human gene GRB14.

[0089] A putative GRB7 signal transduction molecule and a GRB7V novel splice variant from an invasive human esophageal carcinoma was isolated [Tanaka et al., 1998, (32)]. Although both GRB7 isoforms shared homology with the Mig-10 cell migration gene of *Caenorhabditis elegans*, the GRB7V isoform lacked 88 basepairs in the C terminus; the resultant frameshift led to substitution of an SH2 domain with a short hydrophobic sequence. The wildtype GRB7 protein, but not the GRB7V isoform, was rapidly tyrosyl phosphorylated in response to EGF stimulation in esophageal carcinoma cells. Analysis of human esophageal tumor tissues and regional lymph nodes with metastases revealed that GRB7V was expressed in 40% of GRB7-positive esophageal carcinomas. GRB7V expression was enhanced after metastatic spread to lymph nodes as compared to the original tumor tissues. Transfection of an antisense GRB7 RNA expression construct lowered endogenous GRB7 protein levels and suppressed the invasive phenotype exhibited by esophageal carcinoma cells. These findings suggested that GRB7 isoforms are involved in cell invasion and metastatic progression of human esophageal carcinomas. By sequence analysis, The GRB7 gene was mapped to chromosome 17q21-q22, near the topoisomerase-2 gene [Dong et al., 1997, (33)]. GRB-7 is amplified in concert with HER2 in several breast cancer cell lines and that GRB-7 is overexpressed in both cell lines and breast tumors. GRB-7, through its SH2 domain, binds tightly to HER2 such that a large fraction of the tyrosine phosphorylated HER2 in SKBR-3 cells is bound to GRB-7 [Stein et al., 1994, (34)].

GCSF, CSF3

[0090] Granulocyte colony-stimulating factor (or colony stimulating factor-3) specifically stimulates the proliferation and differentiation of the progenitor cells for granulocytes. The partial amino acid sequence of purified GCSF protein was determined, and by using oligonucleotides as probes, several GCSF cDNA clones were isolated from a human squamous carcinoma cell line cDNA library [Nagata et al., 1986, (35)]. Cloning of human GCSF cDNA shows that a single gene codes for a 177- or 180-amino acid mature protein of molecular weight 19,600. The authors found that the GCSF gene has 4 introns and that 2 different polypeptides are synthesized from the same gene by differential splicing of mRNA. The 2 polypeptides differ by the presence or absence of 3 amino acids. Expression studies indicate that both have authentic GCSF activity. A stimulatory activity from a glioblastoma multiform cell line being biologically and biochemically indistinguishable from GCSF produced by a bladder cell line was found in 1987. By somatic cell hybridization and in situ chromosomal hybridization, the GCSF gene was mapped to 17q11 in the region of the breakpoint in the 15;17 translocation characteristic of acute promyelocytic leukemia [Le Beau et al., 1987, (36)]. Further studies indicated that the gene is proximal to the said breakpoint and that it remains on the rearranged chromosome 17. Southern blot analysis using both conventional and pulsed field gel electrophoresis showed no rearranged restriction fragments. By use of a full-length cDNA clone as a hybridization probe in human-mouse somatic cell hybrids and in flow-sorted human chromosomes, the gene for GCSF was mapped to 17q21-q22 lateron

THRA, THRA1, ERBA, EAR7, ERBA2, ERBA3

[0091] Both human and mouse DNA have been demonstrated to have two distantly related classes of ERBA genes and that in the human genome multiple copies of one of the classes exist [Jansson et al., 1983, (37)]. A cDNA was isolated derived from rat brain messenger RNA on the basis of homology to the human thyroid receptor gene [Thompson et al., 1987, (38)]. Expression of this cDNA produced a high-affinity binding protein for thyroid hormones. Messenger RNA from this gene was expressed in tissue-specific fashion, with highest levels in the central nervous system and no expression in the liver. An increasing body of evidence indicated the presence of multiple thyroid hormone receptors. The authors suggested that there may be as many as 5 different but related loci. Many of the clinical and physiologic studies suggested the existence of multiple receptors. For example, patients had been identified with familial thyroid hormone resistance in which peripheral response to thyroid hormones is lost or diminished while neuronal functions are maintained. Thyroidologists recognize a form of cretinism in which the nervous system is severely affected and another form in which the peripheral functions of thyroid hormone are more dramatically affected.

[0092] The cDNA encoding a specific form of thyroid hormone receptor expressed in human liver, kidney, placenta, and brain was isolated [Nakai et al., 1988, (39)]. Identical clones were found in human placenta. The cDNA encodes a protein of 490 amino acids and molecular mass of 54,824. Designated thyroid hormone receptor type alpha-2 (THRA2), this protein is represented by mRNAs of different size in liver and kidney, which may represent tissue-specific processing of the primary transcript.

[0093] The THRA gene contains 10 exons spanning 27 kb of DNA. The last 2 exons of the gene are alternatively spliced. A 5-kb THRA1 mRNA encodes a predicted 410-amino acid protein; a 2.7-kb THRA2 mRNA encodes a 490-amino acid protein. A third isoform, TR-alpha-3, is derived by alternative splicing. The proximal 39 amino acids of the TH-alpha-2 specific sequences are deleted in TR-alpha-3. A second gene, THRB on chromosome 3, encodes 2 isoforms of TR-beta by alternative splicing. In 1989 the structure and function of the EAR1 and EAR7 genes was elucidated, both located on 17q21 [Miyajima et al., 1989, (40)]. The authors determined that one of the exons in the EAR7 coding sequence overlaps an exon of EAR1, and that the 2 genes are transcribed from opposite DNA strands. In addition, the EAR7 mRNA generates 2 alternatively spliced isoforms, referred to as EAR71 and EAR72, of which the EAR71 protein is the human counterpart of the chicken c-erbA protein.

[0094] The thyroid hormone receptors, beta, alpha-1, and alpha-2 3 mRNAs are expressed in all tissues examined and the relative amounts of the three mRNAs were roughly parallel. None of the 3 mRNAs was abundant in liver, which is the major thyroid hormone-responsive organ. This led to the assumption that another thyroid hormone receptor may be present in liver. It was found that ERBA, which potentiates ERBB, has an amino acid sequence different from that of other known oncogene products and related to those of the carbonic anhydrases [Debuire et al., 1984, (41)]. ERBA potentiates ERBB by blocking differentiation of erythroblasts at an immature stage. Carbonic anhydrases participate in the transport of carbon dioxide in erythrocytes. In 1986 it was shown that the ERBA protein is a high-affinity receptor for thyroid hormone. The cDNA sequence indicates a relationship to steroid-hormone receptors, and binding studies indicate that it is a receptor for thyroid hormones. It is located in the nucleus, where it binds to DNA and activates transcription.

[0095] Maternal thyroid hormone is transferred to the fetus early in pregnancy and is postulated to regulate brain development. The ontogeny of TR isoforms and related splice variants in 9 first-trimester fetal brains by semi-quantitative RT-PCR analysis has been investigated. Expression of the TR-beta-1, TR-alpha-1, and TR-alpha-2 isoforms was detected from 8.1 weeks' gestation. An additional truncated species was detected with the TR-alpha-2 primer set, consistent with the TR-alpha-3 splice variant described in the rat. All TR-alpha-derived transcripts were coordinately expressed and increased approximately 8-fold between 8.1 and 13.9 weeks' gestation. A more complex ontogenic pattern was observed for TR-beta-1, suggestive of a nadir between 8.4 and 12.0 weeks' gestation. The authors concluded that these findings point to an important role for the TR-alpha-1 isoform in mediating maternal thyroid hormone action during first-trimester fetal brain development.

[0096] The identification of the several types of thyroid hormone receptor may explain the normal variation in thyroid hormone responsiveness of various organs and the selective tissue abnormalities found in the thyroid hormone resistance syndromes. Members of sibships, who were resistant to thyroid hormone action, had retarded growth, congenital deafness, and abnormal bones, but had normal intellect and sexual maturation, as well as augmented cardiovascular activity. In this family abnormal T3 nuclear receptors in blood cells and fibroblasts have been demonstrated. The availability of cDNAs encoding the various thyroid hormone receptors was considered useful in determining the underlying genetic defect in this family.

[0097] The ERBA oncogene has been assigned to chromosome 17. The ERBA locus remains on chromosome 17 in the t(15;17) translocation of acute promyelocytic leukemia (APL). The thymidine kinase locus is probably translocated to chromosome 15; study of leukemia with t(17;21) and apparently identical breakpoint showed that TK was on 21q+. By in situ hybridization of a cloned DNA probe of c-erb-A to meiotic pachytene spreads obtained from uncultured spermatocytes it has been concluded that ERBA is situated at 17q21.33-17q22, in the same region as the break that generated the t(15;17) seen in APL. Because most of the grains were seen in 17q22, they suggested that ERBA is probably in the proximal region of 17q22 or at the junction between 17q22 and 17q21.33. By in situ hybridization it has been demonstrated, that that ERBA remains at 17q11-q12 in APL, whereas TP53, at 17q21-q22, is translocated to chromosome 15. Thus, ERBA must be at 17q11.2 just proximal to the breakpoint in the APL translocation and just distal to it in the constitutional translocation.

[0098] The aberrant THRA expression in nonfunctioning pituitary tumors has been hypothesized to reflect mutations in the receptor coding and regulatory sequences. They screened THRA mRNA and THRB response elements and ligand-binding domains for sequence anomalies. Screening THRA mRNA from 23 tumors by RNase mismatch and sequencing candidate fragments identified 1 silent and 3 missense mutations, 2 in the common THRA region and 1 that was specific for the alpha-2 isoform. No THRB response element differences were detected in 14 nonfunctioning tumors, and no THRB ligand-binding domain differences were detected in 23 nonfunctioning tumors. Therefore it has been suggested that the novel thyroid receptor mutations may be of functional significance in terms of thyroid receptor action, and further definition of their functional properties may provide insight into the role of thyroid receptors in growth control in pituitary cells.

RAR-alpha

[0099] A cDNA encoding a protein that binds retinoic acid with high affinity has been cloned [Petkovich et al., 1987,

(42)]. The protein was found to be homologous to the receptors for steroid hormones, thyroid hormones, and vitamin D3, and appeared to be a retinoic acid-inducible transacting enhancer factor. Thus, the molecular mechanisms of the effect of vitamin A on embryonic development, differentiation and tumor cell growth may be similar to those described for other members of this nuclear receptor family. In general, the DNA-binding domain is most highly conserved, both within and between the 2 groups of receptors (steroid and thyroid); Using a cDNA probe, the RAR- α gene has been mapped to 17q21 by in situ hybridization [Mattei et al., 1988, (43)]. Evidence has been presented for the existence of 2 retinoic acid receptors, RAR- α and RAR- β , mapping to chromosome 17q21.1 and 3p24, respectively. The α and β forms of RAR were found to be more homologous to the 2 closely related thyroid hormone receptors α and β , located on 17q11.2 and 3p25-p21, respectively, than to any other members of the nuclear receptor family. These observations suggest that the thyroid hormone and retinoic acid receptors evolved by gene, and possibly chromosome, duplications from a common ancestor, which itself diverged rather early in evolution from the common ancestor of the steroid receptor group of the family. They noted that the counterparts of the human RARA and RARB genes are present in both the mouse and chicken. The involvement of RARA at the APL breakpoint may explain why the use of retinoic acid as a therapeutic differentiation agent in the treatment of acute myeloid leukemias is limited to APL. Almost all patients with APL have a chromosomal translocation t(15;17)(q22;q21). Molecular studies reveal that the translocation results in a chimeric gene through fusion between the PML gene on chromosome 15 and the RARA gene on chromosome 17. A hormone-dependent interaction of the nuclear receptors RARA and RXRA with CLOCK and MOP4 has been presented.

20 CDC18 L, CDC 6

[0100] In yeasts, Cdc6 (*Saccharomyces cerevisiae*) and Cdc18 (*Schizosaccharomyces pombe*) associate with the origin recognition complex (ORC) proteins to render cells competent for DNA replication. Thus, Cdc6 has a critical regulatory role in the initiation of DNA replication in yeast. cDNAs encoding *Xenopus* and human homologues of yeast CDC6 have been isolated [Williams et al., 1997, (44)]. They designated the human and *Xenopus* proteins p62(cdc6). Independently, in a yeast 2-hybrid assay using PCNA as bait, cDNAs encoding the human CDC6/Cdc18 homologue have been isolated [Saha et al., 1998, (45)]. These authors reported that the predicted 560-amino acid human protein shares approximately 33% sequence identity with the 2 yeast proteins. On Western blots of HeLa cell extracts, human CDC6/Cdc18 migrates as a 66-kD protein. Although Northern blots indicated that CDC6/Cdc18 mRNA levels peak at the onset of S phase and diminish at the onset of mitosis in HeLa cells, the authors found that total CDC6/Cdc18 protein level is unchanged throughout the cell cycle. Immunofluorescent analysis of epitope-tagged protein revealed that human CDC6/Cdc18 is nuclear in G1 and cytoplasmic in S-phase cells, suggesting that DNA replication may be regulated by either the translocation of this protein between the nucleus and cytoplasm or by selective degradation of the protein in the nucleus. Immunoprecipitation studies showed that human CDC6/Cdc18 associates in vivo with cyclin A, CDK2, and ORC1. The association of cyclin-CDK2 with CDC6/Cdc18 was specifically inhibited by a factor present in mitotic cell extracts. Therefore it has been suggested that if the interaction between CDC6/Cdc18 with the S phase-promoting factor cyclin-CDK2 is essential for the initiation of DNA replication, the mitotic inhibitor of this interaction could prevent a premature interaction until the appropriate time in G1 Cdc6 is expressed selectively in proliferating but not quiescent mammalian cells, both in culture and within tissues in intact animals [Yan et al., 1998, (46)]. During the transition from a growth-arrested to a proliferative state, transcription of mammalian Cdc6 is regulated by E2F proteins, as revealed by a functional analysis of the human Cdc6 promoter and by the ability of exogenously expressed E2F proteins to stimulate the endogenous Cdc6 gene. Immunodepletion of Cdc6 by microinjection of anti-Cdc6 antibody blocked initiation of DNA replication in a human tumor cell line. The authors concluded that expression of human Cdc6 is regulated in response to mitogenic signals through transcriptional control mechanisms involving E2F proteins, and that Cdc6 is required for initiation of DNA replication in mammalian cells.

[0101] Using a yeast 2-hybrid system, co-purification of recombinant proteins, and immunoprecipitation, it has been demonstrated later that an N-terminal segment of CDC6 binds specifically to PR48, a regulatory subunit of protein phosphatase 2A (PP2A). The authors hypothesized that dephosphorylation of CDC6 by PP2A, mediated by a specific interaction with PR48 or a related B-double prime protein, is a regulatory event controlling initiation of DNA replication in mammalian cells. By analysis of somatic cell hybrids and by fluorescence in situ hybridization the human p62(cdc6) gene has been to 17q21.3.

TOP2A, TOP2

[0102] DNA topoisomerases are enzymes that control and alter the topologic states of DNA in both prokaryotes and eukaryotes. Topoisomerase II from eukaryotic cells catalyzes the relaxation of supercoiled DNA molecules, catenation, decatenation, knotting, and unknotting of circular DNA. It appears likely that the reaction catalyzed by topoisomerase II involves the crossing-over of 2 DNA segments. It has been estimated that there are about 100,000 molecules of

topoisomerase II per HeLa cell nucleus, constituting about 0.1% of the nuclear extract. Since several of the abnormal characteristics of ataxia-telangiectasia appear to be due to defects in DNA processing, screening for these enzyme activities in 5 AT cell lines has been performed [Singh et al., 1988, (47)]. In comparison to controls, the level of DNA topoisomerase II, determined by unknottting of P4 phage DNA, was reduced substantially in 4 of these cell lines and to a lesser extent in the fifth. DNA topoisomerase I, assayed by relaxation of supercoil DNA, was found to be present at normal levels.

[0103] The entire coding sequence of the human TOP2 gene has been determined [Tsai-Pflugfelder et al., 1988, (48)].

[0104] In addition human cDNAs that had been isolated by screening a cDNA library derived from a mechlorethamine-resistant Burkitt lymphoma cell line (Raji-HN2) with a *Drosophila* Topo II cDNA had been sequenced [Chung et al., 1989, (49)]. The authors identified 2 classes of sequence representing 2 TOP2 isoenzymes, which have been named TOP2A and TOP2B. The sequence of 1 of the TOP2A cDNAs is identical to that of an internal fragment of the TOP2 cDNA isolated by Tsai-Pflugfelder et al., 1988 (48). Southern blot analysis indicated that the TOP2A and TOP2B cDNAs are derived from distinct genes. Northern blot analysis using a TOP2A-specific probe detected a 6.5-kb transcript in the human cell line U937. Antibodies against a TOP2A peptide recognized a 170-kD protein in U937 cell lysates. Therefore it was concluded that their data provide genetic and immunochemical evidence for 2 TOP2 isozymes. The complete structures of the TOP2A and TOP2B genes has been reported [Lang et al., 1998, (50)]. The TOP2A gene spans approximately 30 kb and contains 35 exons.

[0105] Tsai-Pflugfelder et al., 1988 (48) showed that the human enzyme is encoded by a single-copy gene which they mapped to 17q21-q22 by a combination of *in situ* hybridization of a cloned fragment to metaphase chromosomes and by Southern hybridization analysis with a panel of mouse-human hybrid cell lines. The assignment to chromosome 17 has been confirmed by the study of somatic cell hybrids. Because of co-amplification in an adenocarcinoma cell line, it was concluded that the TOP2A and ERBB2 genes may be closely linked on chromosome 17 [Keith et al., 1992, (51)]. Using probes that detected RFLPs at both the TOP2A and TOP2B loci, the demonstrated heterozygosity at a frequency of 0.17 and 0.37 for the alpha and beta loci, respectively. The mouse homologue was mapped to chromosome 11 [Kingsmore et al., 1993, (52)]. The structure and function of type II DNA topoisomerases has been reviewed [Watt et al., 1994, (53)]. DNA topoisomerase II- α is associated with the pol II holoenzyme and is a required component of chromatin-dependent co-activation. Specific inhibitors of topoisomerase II blocked transcription on chromatin templates, but did not affect transcription on naked templates. Addition of purified topoisomerase II- α reconstituted chromatin-dependent activation activity in reactions with core pol II. Therefore the transcription on chromatin templates seems to result in the accumulation of superhelical tension, making the relaxation activity of topoisomerase II essential for productive RNA synthesis on nucleosomal DNA.

IGFBP4

[0106] Six structurally distinct insulin-like growth factor binding proteins have been isolated and their cDNAs cloned: IGFBP1, IGFBP2, IGFBP3, IGFBP4, IGFBP5 and IGFBP6. The proteins display strong sequence homologies, suggesting that they are encoded by a closely related family of genes. The IGFBPs contain 3 structurally distinct domains each comprising approximately one-third of the molecule. The N-terminal domain 1 and the C-terminal domain 3 of the 6 human IGFBPs show moderate to high levels of sequence identity including 12 and 6 invariant cysteine residues in domains 1 and 3, respectively (IGFBP6 contains 10 cysteine residues in domain 1), and are thought to be the IGF binding domains. Domain 2 is defined primarily by a lack of sequence identity among the 6 IGFBPs and by a lack of cysteine residues, though it does contain 2 cysteines in IGFBP4. Domain 3 is homologous to the thyroglobulin type I repeat unit. Recombinant human insulin-like growth factor binding proteins 4, 5, and 6 have been characterized by their expression in yeast as fusion proteins with ubiquitin [Kiefer et al., 1992, (54)]. Results of the study suggested to the authors that the primary effect of the 3 proteins is the attenuation of IGF activity and suggested that they contribute to the control of IGF-mediated cell growth and metabolism.

[0107] Based on peptide sequences of a purified insulin-like growth factor-binding protein (IGFBP) rat IGFBP4 has been cloned by using PCR [Shimasaki et al., 1990, (55)]. They used the rat cDNA to clone the human ortholog from a liver cDNA library. Human IGFBP4 encodes a 258-amino acid polypeptide, which includes a 21-amino acid signal sequence. The protein is very hydrophilic, which may facilitate its ability as a carrier protein for the IGFs in blood. Northern blot analysis of rat tissues revealed expression in all tissues examined, with highest expression in liver. It was stated that IGFBP4 acts as an inhibitor of IGF-induced bone cell proliferation. The genomic region containing the IGFBP gene. The gene consists of 4 exons spanning approximately 15 kb of genomic DNA has been examined [Zazzi et al., 1998, (56)]. The upstream region of the gene contains a TATA box and a cAMP-responsive promoter.

[0108] By *in situ* hybridization, the IGFBP4 gene was mapped to 17q12-q21 [Bajajica et al., 1992, (57)]. Because the hereditary breast-ovarian cancer gene BRCA1 had been mapped to the same region, it has been investigated whether IGFBP4 is a candidate gene by linkage analysis of 22 BRCA1 families; the finding of genetic recombination suggested that it is not the BRCA1 gene [Tonin et al., 1993, (58)].

EB1 1, CCR7, CMKBR7

[0109] Using PCR with degenerate oligonucleotides, a lymphoid-specific member of the G protein-coupled receptor family has been identified and mapped mapped to 17q12-q21.2 by analysis of human/mouse somatic cell hybrid DNAs and fluorescence in situ hybridization. It has been shown that this receptor had been independently identified as the Epstein-Barr-induced cDNA (symbol EB11) [Birkenbach et al., 1993, (59)]. EB11 is expressed in normal lymphoid tissues and in several B- and T-lymphocyte cell lines. While the function and the ligand for EB11 remains unknown, its sequence and gene structure suggest that it is related to receptors that recognize chemoattractants, such as interleukin-8, RANTES, C5a, and fMet-Leu-Phe. Like the chemoattractant receptors, EB11 contains intervening sequences near its 5-prime end; however, EB11 is unique in that both of its introns interrupt the coding region of the first extracellular domain. Mouse Eb11 cDNA has been isolated and found to encode a protein with 86% identity to the human homologue.

[0110] Subsets of murine CD4⁺ T cells localize to different areas of the spleen after adoptive transfer. Naive and T helper-1 (TH1) cells, which express CCR7, home to the periarteriolar lymphoid sheath, whereas activated TH2 cells, which lack CCR7, form rings at the periphery of the T-cell zones near B-cell follicles. It has been found that retroviral transduction of TH2 cells with CCR7 forced them to localize in a TH1-like pattern and inhibited their participation in B-cell help in vivo but not in vitro. Apparently differential expression of chemokine receptors results in unique cellular migration patterns that are important for effective immune responses.

[0111] CCR7 expression divides human memory T cells into 2 functionally distinct subsets. CCR7-memory cells express receptors for migration to inflamed tissues and display immediate effector function. In contrast, CCR7⁺ memory cells express lymph node homing receptors and lack immediate effector function, but efficiently stimulate dendritic cells and differentiate into CCR7⁺ effector cells upon secondary stimulation. The CCR7⁺ and CCR7⁻ T cells, named central memory (T-CM) and effector memory (T-EM), differentiate in a step-wise fashion from naive T cells, persist for years after immunization, and allow a division of labor in the memory response.

[0112] CCR7 expression in memory CD8⁺ T lymphocyte responses to HIV and to cytomegalovirus (CMV) tetramers has been evaluated. Most memory T lymphocytes express CD45RO, but a fraction express instead the CD45RA marker. Flow cytometric analyses of marker expression and cell division identified 4 subsets of HIV- and CMV-specific CD8⁺ T cells, representing a lineage differentiation pattern: CD45RA⁺CCR7⁺ (double-positive); CD45RA⁺CCR7⁻; CD45RA⁺CCR7⁻ (double-negative); CD45RA⁺CCR7⁻. The capacity for cell division, as measured by 5-(and 6-)carboxyfluorescein diacetate, succinimidyl ester, and intracellular staining for the Ki67 nuclear antigen, is largely confined to the CCR7⁺ subsets and occurred more rapidly in cells that are also CD45RA⁺. Although the double-negative cells did not divide or expand after stimulation, they did revert to positivity for either CD45RA or CCR7 or both. The CD45RA⁺CCR7⁻ cells, considered to be terminally differentiated, fail to divide, but do produce interferon-gamma and express high levels of perforin. The representation of subsets specific for CMV and for HIV is distinct. Approximately 70% of HIV-specific CD8⁺ memory T cells are double-negative or preterminally differentiated compared to 40% of CMV-specific cells. Approximately 50% of the CMV-specific CD8⁺ memory T cells are terminally differentiated compared to fewer than 10% of the HIV-specific cells. It has been proposed that terminally differentiated CMV-specific cells are poised to rapidly intervene, while double-positive precursor cells remain for expansion and replenishment of the effector cell pool. Furthermore, high-dose antigen tolerance and the depletion of HIV-specific CD4⁺ helper T-cell activity may keep the HIV-specific memory CD8⁺ T cells at the double-negative stage, unable to differentiate to the terminal effector state. B lymphocytes recirculate between B cell-rich compartments (follicles or B zones) in secondary lymphoid organs, surveying for antigen. After antigen binding, B cells move to the boundary of B and T zones to interact with T-helper cells. Furthermore it has been demonstrated that antigen-engaged B cells have increased expression of CCR7, the receptor for the T-zone chemokines CCL19 (also known as ELC) and CCL21, and that they exhibit increased responsiveness to both chemoattractants. In mice lacking lymphoid CCL19 and CCL21 chemokines, or with B cells that lack CCR7, antigen engagement fails to cause movement to the T zone. Using retroviral-mediated gene transfer, the authors demonstrated that increased expression of CCR7 is sufficient to direct B cells to the T zone. Reciprocally, overexpression of CXCR5, the receptor for the B-zone chemokine CXCL13, is sufficient to overcome antigen-induced B-cell movement to the T zone. This points toward a mechanism of B-cell relocalization in response to antigen, and established that cell position in vivo can be determined by the balance of responsiveness to chemoattractants made in separate but adjacent zones.

BAF57, SMARCE 1

[0113] The SWI/SNF complex in *S. cerevisiae* and *Drosophila* is thought to facilitate transcriptional activation of specific genes by antagonizing chromatin-mediated transcriptional repression. The complex contains an ATP-dependent nucleosome disruption activity that can lead to enhanced binding of transcription factors. The BRG1/brm-associated factors, or BAF, complex in mammals is functionally related to SWI/SNF and consists of 9 to 12 subunits, some of which are homologous to SWI/SNF subunits. A 57-kD BAF subunit, BAF57, is present in higher eukaryotes, but not in

yeast. Partial coding sequence has been obtained from purified BAF57 from extracts of a human cell line [Wang et al., 1998, (60)]. Based on the peptide sequences, they identified cDNAs encoding BAF57. The predicted 411-amino acid protein contains an HMG domain adjacent to a kinesin-like region. Both recombinant BAF57 and the whole BAF complex bind 4-way junction (4WJ) DNA, which is thought to mimic the topology of DNA as it enters or exits the nucleosome. The BAF57 DNA-binding activity has characteristics similar to those of other HMG proteins. It was found that complexes with mutations in the BAF57 HMG domain retain their DNA-binding and nucleosome-disruption activities. They suggested that the mechanism by which mammalian SWI/SNF-like complexes interact with chromatin may involve recognition of higher-order chromatin structure by 2 or more DNA-binding domains. RNase protection studies and Western blot analysis revealed that BAF57 is expressed ubiquitously. Several lines of evidence point toward the involvement of SWI/SNF factors in cancer development [Klochendler-Yeivin et al., 2002, (61)]. Moreover, SWI/SNF related genes are assigned to chromosomal regions that are frequently involved in somatic rearrangements in human cancers [Ring et al., 1998, (62)]. In this respect it is interesting that some of the SWI/SNF family members (i.e. SMARCC1, SMARCC2, SMARCD1 and SMARCD22 are neighboring 3 of the eucaryotic ARCHEONs we have identified (i.e. 3p21-p24, 12q13-q14 and 17q respectively) and which are part of the present invention. In this invention we could also map SMARCE1/BAF57 to the 17q12 region by PCR karyotyping.

KRT 10, K10

[0114] Keratin 10 is an intermediate filament (IF) chain which belongs to the acidic type I family and is expressed in terminally differentiated epidermal cells. Epithelial cells almost always co-express pairs of type I and type II keratins, and the pairs that are co-expressed are highly characteristic of a given epithelial tissue. For example, in human epidermis, 3 different pairs of keratins are expressed: keratins 5 (type II) and 14 (type I), characteristic of basal or proliferative cells; keratins 1 (type II) and 10 (type I), characteristic of suprabasal terminally differentiating cells; and keratins 6 (type II) and 16 (type I) (and keratin 17 [type I]), characteristic of cells induced to hyper-proliferate by disease or injury, and epithelial cells grown in cell culture. The nucleotide sequence of a 1,700 bp cDNA encoding human epidermal keratin 10 (56.5 kD) [Darmon et al., 1987, (63)] has been published as well as the complete amino acid sequence of human keratin 10 [Zhou et al., 1988, (64)]. Polymorphism of the KRT10 gene, restricted to insertions and deletions of the glycine-rich quasipeptide repeats that form the glycine-loop motif in the C-terminal domain, have been extensively described [Korge et al., 1992, (65)].

[0115] By use of specific cDNA clones in conjunction with somatic cell hybrid analysis and in situ hybridization, KRT10 gene has been mapped to 17q12-q21 in a region proximal to the breakpoint at 17q21 that is involved in a t(17;21)(q21;q22) translocation associated with a form of acute leukemia. KRT10 appeared to be telomeric to 3 other loci that map in the same region: CSF3, ERBA1, and HER2 [Lessin et al., 1988, (66)]. NGFR and HOX2 are distal to K9. It has been demonstrated that the KRT10, KRT13, and KRT15 genes are located in the same large pulsed field gel electrophoresis fragment [Romano et al., 1991, (67)]. A correlation of assignments of the 3 genes makes 17q21-q22 the likely location of the cluster. Transgenic mice expressing a mutant keratin 10 gene have the phenotype of epidermolytic hyperkeratosis, thus suggesting that a genetic basis for the human disorder resides in mutations in genes encoding suprabasal keratins KRT1 or KRT10 [Fuchs et al 1992, (68)]. The authors also showed that stimulation of basal cell proliferation can result from a defect in suprabasal cells and that distortion of nuclear shape or alterations in cytokinesis can occur when an intermediate filament network is perturbed. In a family with keratosis palmaris et plantaris without blistering either spontaneously or in response to mild mechanical or thermal stress and with no involvement of the skin and parts of the body other than the palms and soles, a tight linkage to an insertion-deletion polymorphism in the C-terminal coding region of the KRT10 gene (maximum lod score = 8.36 at theta = 0.00) was found [Rogaev et al., 1993, (69)]. It is noteworthy that it was a rare, high molecular weight allele of the KRT10 polymorphism that segregated with the disorder. The allele was observed once in 96 independent chromosomes from unaffected Caucasians. The KRT10 polymorphism arose from the insertion/deletion of imperfect (CCG)_n repeats within the coding region and gave rise to a variable glycine loop motif in the C-terminal tail of the keratin 10 protein. It is possible that there was a pathogenic role for the expansion of the imperfect trinucleotide repeat.

KRT12, K12

[0116] Keratins are a group of water-insoluble proteins that form 10 nm intermediate filaments in epithelial cells. Approximately 30 different keratin molecules have been identified. They can be divided into acidic and basic-neutral subfamilies according to their relative charges, immunoreactivity, and sequence homologies to types I and II wool keratins, respectively. In vivo, a basic keratin usually is co-expressed and 'paired' with a particular acidic keratin to form a heterodimer. The expression of various keratin pairs is tissue specific, differentiation dependent, and developmentally regulated. The presence of specific keratin pairs is essential for the maintenance of the integrity of epithelium. For example, mutations in human K14/K5 pair and the K10/K1 pair underlie the skin diseases, epidermolysis bullosa

simplex and epidermolytic hyperkeratosis, respectively. Expression of the K3 and K12 keratin pair have been found in the cornea of a wide number of species, including human, mouse, and chicken, and is regarded as a marker for corneal-type epithelial differentiation. The murine Krt12 (Krt1.12) gene and demonstrated that its expression is corneal epithelial cell specific, differentiation dependent, and developmentally regulated [Liu et al., 1993, (70)]. The corneal-specific nature of keratin 12 gene expression signifies keratin 12 plays a unique role in maintaining normal corneal epithelial function. Nevertheless, the exact function of keratin 12 remains unknown and no hereditary human corneal epithelial disorder has been linked directly to the mutation in the keratin 12 gene. As part of a study of the expression profile of human corneal epithelial cells, a cDNA with an open reading frame highly homologous to the cornea-specific mouse keratin 12 gene has been isolated [Nishida et al., 1996, (71)]. To elucidate the function of keratin 12 knockout mice lacking the Krt1.12 gene have been created by gene targeting techniques. The heterozygous mice appeared normal. Homozygous mice developed normally and suffered mild corneal epithelial erosion. The corneal epithelia were fragile and could be removed by gentle rubbing of the eyes or brushing. The corneal epithelium of the homozygotes did not express keratin 12 as judged by immunohistochemistry, Western immunoblot analysis with epitope-specific anti-keratin 12 antibodies, Northern hybridization, and in situ hybridization with an antisense keratin 12 riboprobe. The KRT12 gene has been mapped to 17q by study of radiation hybrids and localized it to the type I keratin cluster in the interval between D17S800 and D17S930 (17q12-q21) [Nishida et al., 1997, (72)]. The authors presented the exon-intron boundary structure of the KRT12 gene and mapped the gene to 17q12 by fluorescence in situ hybridization. The gene contains 7 introns, defining 8 exons that cover the coding sequence. Together the exons and introns span approximately 6 kb of genomic DNA.

[0117] Meesmann corneal dystrophy is an autosomal dominant disorder causing fragility of the anterior corneal epithelium, where the cornea-specific keratins K3 and K12 are expressed. Dominant-negative mutations in these keratins might be the cause of Meesmann corneal dystrophy. Indeed, linkage of the disorder to the K12 locus in Meesmann's original German kindred [Meesmann and Wilke, 1939, (73)] with $Z(\max) = 7.53$ at $\theta = 0.0$ has been found. In 2 pedigrees from Northern Ireland, they found that the disorder co-segregated with K12 in one pedigree and K3 in the other. Heterozygous missense mutations in K3 or in K12 (R135T, V143L...) in each family have been identified. All these mutations occurred in highly conserved keratin helix boundary motifs, where dominant mutations in other keratins have been found to compromise cytoskeletal function severely, leading to keratinocyte fragility.

[0118] The regions of the human KRT12 gene have been sequenced to enable mutation detection for all exons using genomic DNA as a template [Corden et al., 2000, (74)]. The authors found that the human genomic sequence spans 5,919 bp and consists of 8 exons. A microsatellite dinucleotide repeat was identified within intron 3, which was highly polymorphic and which they developed for use in genotype analysis. In addition, 2 mutations in the helix initiation motif of K12 were found in families with Meesmann corneal dystrophy. In an American kindred, a missense M129T mutation was found in the KRT12 gene. They stated that a total of 8 mutations in the KRT12 gene had been reported.

Genetic interactions within ARCHEONS

[0119] Genes involved in genomic alterations (amplifications, insertions, translocations, deletions, etc.) exhibit changes in their expression pattern. Of particular interest are gene amplifications, which account for gene copy numbers >2 per cell or deletions accounting for gene copy numbers <2 per cell. Gene copy number and gene expression of the respective genes do not necessarily correlate. Transcriptional overexpression needs an intact transcriptional context, as determined by regulatory regions at the chromosomal locus (promoter, enhancer and silencer), and sufficient amounts of transcriptional regulators being present in effective combinations. This is especially true for genomic regions, which expression is tightly regulated in specific tissues or during specific developmental stages. ARCHEONS are specified by gene clusters of more than two genes being directly neighbored or in chromosomal order, interspersed by a maximum of 10, preferably 7, more preferably 5 or at least 1 gene. The interspersed genes are also co-amplified but do not directly interact with the ARCHEON. Such an ARCHEON may spread over a chromosomal region of a maximum of 20, more preferably 10 or at least 6 Megabases. The nature of an ARCHEON is characterized by the simultaneous amplification and/or deletion and the correlating expression (i.e. upregulation or downregulation respectively) of the encompassed genes in a specific tissue, cell type, cellular or developmental state or time point. Such ARCHEONS are commonly conserved during evolution, as they play critical roles during cellular development. In case of these ARCHEONS whole gene clusters are overexpressed upon amplification as they harbor self-regulatory feedback loops, which stabilize gene expression and/or biological effector function even in abnormal biological settings, or are regulated by very similar transcription factor combinations, reflecting their simultaneous function in specific tissues at certain developmental stages. Therefore, the gene copy numbers correlates with the expression level especially for genes in gene clusters functioning as ARCHEONS. In case of abnormal gene expressions in neoplastic lesions it is of great importance to know whether the self-regulatory feedback loops have been conserved as they determine the biological activity of the ARCHEON gene members.

[0120] The intensive interaction between genes in ARCHEONS is described for the 17q12 ARCHEON (Fig. 1) by

way of illustration not by limitation. In one embodiment the presence or absence of alterations of genes within distinct genomic regions are correlated with each other, as exemplified for breast cancer cell lines (Fig.3 and Fig. 4). This confers to the discovery of the present invention, that multiple interactions of said gene products of defined chromosomal localizations happen, that according to their respective alterations in abnormal tissue have predictive, diagnostic, prognostic and/or preventive and therapeutic value. These interactions are mediated directly or indirectly, due to the fact that the respective genes are part of interconnected or independent signaling networks or regulate cellular behavior (differentiation status, proliferative and/or apoptotic capacity, invasiveness, drug responsiveness, immune modulatory activities) in a synergistic, antagonistic or independent fashion. The order of functionally important genes within the ARCHEONS has been conserved during evolution (e.g. the ARCHEON on human chromosome 17q12 is present on mouse chromosome 11). Moreover, it has been found that the 17q12 ARCHEON is also present on human chromosome 3p21 and 12q13, both of which are also involved in amplification events and in tumor development. Most probably these homologous ARCHEONS were formed by duplications and rearrangements during vertebrate evolution. Homologous ARCHEONS consist of homologous genes and/or isoforms of specific gene families (e.g. RARA or RARB or RARG, THRA or THRB, TOP2A or TOP2B, RABSA or RABSB, BAF170 or BAF 155, BAF60A or BAF60B, WNT5A or WNT5B, IGFBP4 or IGFBP6). Moreover these regions are flanked by homologous chromosomal gene clusters (e.g. CACN, SCYA, HOX, Keratins). These ARCHEONS have diverged during evolution to fulfill their respective functions in distinct tissues (e.g. the 17q12 ARCHEON has one of its main functions in the central nervous system). Due to their tissue specific function extensive regulatory loops control the expression of the members of each ARCHEON. During tumor development these regulations become critical for the characteristics of the abnormal tissues with respect to differentiation, proliferation, drug responsiveness, invasiveness. It has been found that the co-amplification of genes within ARCHEONS can lead to co-expression of the respective gene products. Some of said genes also exhibit additional mutations or specific patterns of polymorphisms, which are substantial for the oncogenic capacities of these ARCHEONS. It is one of the critical features of such amplicons, which members of the ARCHEON have been conserved during tumor formation (e.g. during amplification and deletion events), thereby defining these genes as diagnostic marker genes. Moreover, the expression of the certain genes within the ARCHEON can be influenced by other members of the ARCHEON, thereby defining the regulatory and regulated genes as target genes for therapeutic intervention. It was also observed, that the expression of certain members of the ARCHEON is sensitive to drug treatment (e.g. TOPO2 alpha, RARA, THRA, HER-2) which defines these genes as "marker genes". Moreover several other genes are suitable for therapeutic intervention by antibodies (CACNB1, EB1), ligands (CACNB1) or drugs like e.g. kinase inhibitors (CrkRS, CDC6). The following examples of interactions between members of ARCHEONS are offered by way of illustration, not by way of limitation.

[0121] EB1/CCR7 is lymphoid-specific member of the G protein-coupled receptor family. EB1 recognizes chemotactants, such as interleukin-8, SCYAs, Rantes, C5a, and fMet-Leu-Phe. The capacity for cell division is largely confined to the CCR7⁺ subsets in lymphocytes. Double-negative cells did not divide or expand after stimulation. CCR7⁺ cells, considered to be terminally differentiated, fail to divide, but do produce interferon-gamma and express high levels of perforin. EB1 is induced by viral activities such as the Epstein-Barr-Virus. Therefore, EB1 is associated with transformation events in lymphocytes. A functional role of EB1 during tumor formation in non-lymphoid tissues has been investigated in this invention. Interestingly, also ERBA and ERBB, located in the same genomic region, are associated with lymphocyte transformation. Moreover, ligands of the receptor (i.e. SCYA5/Rantes) are in genomic proximity on 17q. Abnormal expression of both of these factors in lymphoid and non-lymphoid tissues establishes an aut-regulatory feedback loop, inducing signaling events within the respective cells. Expression of lymphoid factors has effect on immune cells and modulates cellular behavior. This is of particular interest with regard to abnormal breast tissue being infiltrated by lymphocytes. In line with this, another immunomodulatory and proliferation factor is located nearby on 17q12. Granulocyte colony-stimulating factor (GCSF3) specifically stimulates the proliferation and differentiation of the progenitor cells for granulocytes. A stimulatory activity from a glioblastoma multiforme cell line being biologically and biochemically indistinguishable from GCSF produced by a bladder cell line has also been found. Colony-stimulating factors not only affects immune cells, but also induce cellular responses of non-immune cells, indicating possible involvement in tumor development upon abnormal expression. In addition several other genes of the 17q12 ARCHEON are involved in proliferation, survival, differentiation of immune cells and/or lymphoblastic leukemia, such as MLLT6, ZNF144 and ZNFN1A3, again demonstrating the related functions of the gene products in interconnected key processes within specific cell types. Aberrant expression of more than one of these genes in non-immune cells constitutes signalling activities, that contribute to the oncogenic activities that derive solely from overexpression of the Her-2/neu gene.

[0122] PPARBP has been found in complex with the tumorsuppressor gene of the p53 family. Moreover, PPARBP also binds to PPAR-alpha (PPARA), RAR-alpha (RARA), RXR, THRA and TR-beta-1. Due to it's ability to bind to thyroid hormone receptors it has been named TRIP2 and TRAP220. In this complex PPARBP affects gene regulatory activities. Interestingly, PPARBP is located in genomic proximity to its interaction partners THRA and RARA. We have found PPARBP to be co-amplified with THRA and RARA in tumor tissue. THRA has been isolated from avian eryth-

robolastosis virus in conjunction with ERBB and therefore was named ERBA. ERBA potentiates ERBB by blocking differentiation of erythroblasts at an immature stage. ERBA has been shown to influence ERBB expression. In this setting deletions of C-terminal portions of the THRA gene product are of influence. Aberrant THRA expression has also been found in nonfunctioning pituitary tumors, which has been hypothesized to reflect mutations in the receptor coding and regulatory sequences. THRA function promotes tumor cell development by regulating gene expression of regulatory genes and by influencing metabolic activities (e.g. of key enzymes of alternative metabolic pathways in tumors such as malic enzyme and genes responsible for lipogenesis). The observed activities of nuclear receptors not only reflect their transactivating potential, but are also due to posttranscriptional activities in the absence or presence of ligands. Co-amplification of THRA/ERBA and ERBB has been shown, but its influence on tumor development has been doubted as no overexpression could be demonstrated in breast tumors [van de Vijver et al., 1987, (75)]. THRA and RARA are part of nuclear receptor family whose function can be mediated as monomers, homodimers or heterodimers. RARA regulates differentiation of a broad spectrum of cells. Interactions of hormones with ERBB expression has been investigated. Ligands of RARA can inhibit the expression of amplified ERBB genes in breast tumors [Offender et al., 1998, (76)]. As being part of this invention co-amplification and co-expression of THRA and RARA could be shown. It was also found that multiple genes, which are regulated by members of the thyroid hormone receptor - and retinoic acid receptor family, are differentially expressed in tumor samples, corresponding to their genomic alterations (amplification, mutation, deletion). These hormone receptor genes and respective target genes are useful to discriminate patient samples with respect to clinical features.

[0123] By expression analysis of multiple normal tissues, tumor samples and tumor cell lines and subsequent clustering of the 17q12 region, it was found that the expression profile of Her-2/neu positive tumor cells and tumor samples exhibits similarities with the expression pattern of tissue from the central nervous system (Fig. 2). This is in line with the observed malformations in the central nervous system of Her-2/neu and THRA knock-out mice. Moreover, it was found that NEUROD2, a nuclear factor involved specifically in neurogenesis, is commonly expressed in the respective samples. This led to the definition of the 17q12 Locus as being an "ARCHEON", whose primary function in normal organ development is defined to the central nervous system. Surprisingly, the expression of NEUROD2 was affected by therapeutic intervention. Strikingly, also ZNF144, TEM7, PIP5K and PPP1R1B are expressed in neuronal cells, where they display diverse tissue specific functions.

[0124] In addition Her-2/neu is often co-amplified with GRB7, a downstream member of the signaling cascade being involved in invasive properties of tumors. Surprisingly, we have found another member of the Her-2/neu signaling cascade being overexpressed in primary breast tumors TOB1 (= "Transducer of ERBB signaling"). Strong overexpression of TOB1 correlated with weaker overexpression of Her-2/neu, already indicating its involvement in oncogenic signaling activities. Amplification of Her-2/neu has been assigned to enhanced proliferative capacity, due to the identified downstream components of the signaling cascade (e.g. Ras-Raf-MAPK). In this respect it was surprising that some cdc genes, which are cell cycle dependent kinases, are part of the amplicons, which upon altered expression have great impact on cell cycle progression.

[0125] According to the observations described above the following examples of genes at 3q21-26 are offered by way of illustration, not by way of limitation.

→ WNT5A, CACNA1D, THRB, RARB, TOP2B, RAB5B, SMARCC1 (BAF155), RAF, WNT7A

[0126] The following examples of genes at 12q13 are offered by way of illustration, not by way of limitation.

→ CACNB3, Keratins, NR4A1, RAB5/13, RARgamma, STAT6, WNT10B, (GCN5), (SAS: Sarcoma Amplified Sequence), SMARCC2 (BAF170), SMARCD1 (BAF60A), (GAS41: Glioma Amplified Sequence), (CHOP), Her3, KRTHB, HOC X, IGFBP6, WNT5B

[0127] There is cross-talk between the amplified ARCHEONs described above and some other highly amplified genomic regions locate approximately at 1p13, 1q32, 2p16, 2q21, 3p12, 5p13, 6p12, 7p12, 7q21, 8q23, 11q13, 13q12, 19q13, 20q13 and 21q11. The above mentioned chromosomal regions are described by way of illustration not by way of limitation, as the amplified regions often span larger and/or overlapping positions at these chromosomal positions.

[0128] Additional alterations of non-transcribed genes, pseudogenes or intergenic regions of said chromosomal locations can be measured for prediction, diagnosis, prognosis, prevention and treatment of malignant neoplasia and breast cancer in particular. Some of the genes or genomic regions have no direct influence on the members of the ARCHEONs or the genes within distinct chromosomal regions but still retain marker gene function due to their chromosomal positioning in the neighborhood of functionally critical genes (e.g. Telethonin neighboring the Her-2/neu gene).

[0129] The invention further relates to the use of:

a) a polynucleotide comprising at least one of the sequences of SEQ ID NO: 1 to 26 or 53 to 75;

b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

c) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c)

e) an antisense molecule targeting specifically one of the polynucleotide sequences specified in (a) to (d);

f) a purified polypeptide encoded by a polynucleotide sequence specified in (a) to (d)

g) a purified polypeptide comprising at least one of the sequences of SEQ ID NO: 27 to 52 or 76 to 98;

h) an antibody capable of binding to one of the polynucleotide specified in (a) to (d) or a polypeptide specified in (f) and (g)

i) a reagent identified by any of the methods of claim 14 to 16 that modulates the amount or activity of a polynucleotide sequence specified in (a) to (d) or a polypeptide specified in (f) and (g)

in the preparation of a composition for the prevention, prediction, diagnosis, prognosis or a medicament for the treatment of malignant neoplasia and breast cancer in particular.

Polynucleotides

[0130] A "BREAST CANCER GENE" polynucleotide can be single- or double-stranded and comprises a coding sequence or the complement of a coding sequence for a "BREAST CANCER GENE" polypeptide. Degenerate nucleotide sequences encoding human "BREAST CANCER GENE" polypeptides, as well as homologous nucleotide sequences which are at least about 50, 55, 60, 65, 70, preferably about 75, 90, 96, or 98% identical to the nucleotide sequences of SEQ ID NO: 1 to 26 or 53 to 75 also are "BREAST CANCER GENE" polynucleotides. Percent sequence identity between the sequences of two polynucleotides is determined using computer programs such as ALIGN which employ the FASTA algorithm, using an affine gap search with a gap open penalty of -12 and a gap extension penalty of -2. Complementary DNA (cDNA) molecules, species homologues, and variants of "BREAST CANCER GENE" polynucleotides which encode biologically active "BREAST CANCER GENE" polypeptides also are "BREAST CANCER GENE" polynucleotides.

Preparation of Polynucleotides

[0131] A naturally occurring "BREAST CANCER GENE" polynucleotide can be isolated free of other cellular components such as membrane components, proteins, and lipids. Polynucleotides can be made by a cell and isolated using standard nucleic acid purification techniques, or synthesized using an amplification technique, such as the polymerase chain reaction (PCR), or by using an automatic synthesizer. Methods for isolating polynucleotides are routine and are known in the art. Any such technique for obtaining a polynucleotide can be used to obtain isolated "BREAST CANCER GENE" polynucleotides. For example, restriction enzymes and probes can be used to isolate polynucleotide fragments which comprises "BREAST CANCER GENE" nucleotide sequences. Isolated polynucleotides are in preparations which are free or at least 70, 80, or 90% free of other molecules.

[0132] "BREAST CANCER GENE" cDNA molecules can be made with standard molecular biology techniques, using "BREAST CANCER GENE" mRNA as a template. Any RNA isolation technique which does not select against the isolation of mRNA may be utilized for the purification of such RNA samples. See, for example, Sambrook et al., 1989, (77); and Ausubel, F. M. et al., 1989, (78), both of which are incorporated herein by reference in their entirety. Additionally, large numbers of tissue samples may readily be processed using techniques well known to those of skill in the art, such as, for example, the single-step RNA isolation process of Chomczynski, P. (1989, U.S. Pat. No. 4,843,155), which is incorporated herein by reference in its entirety.

[0133] "BREAST CANCER GENE" cDNA molecules can thereafter be replicated using molecular biology techniques known in the art and disclosed in manuals such as Sambrook et al., 1989, (77). An amplification technique, such as PCR, can be used to obtain additional copies of polynucleotides of the invention, using either human genomic DNA or

cDNA as a template.

[0134] Alternatively, synthetic chemistry techniques can be used to synthesize "BREAST CANCER GENE" polynucleotides. The degeneracy of the genetic code allows alternate nucleotide sequences to be synthesized which will encode a "BREAST CANCER GENE" polypeptide or a biologically active variant thereof.

Identification of differential expression

[0135] Transcripts within the collected RNA samples which represent RNA produced by differentially expressed genes may be identified by utilizing a variety of methods which are well known to those of skill in the art. For example, differential screening [Tedder, T. F. et al., 1988, (79)], subtractive hybridization [Hedrick, S. M. et al., 1984, (80); Lee, S. W. et al., 1984, (81)], and, preferably, differential display [Liang, P., and Pardee, A. B., 1993, U.S. Pat. No. 5,262,311, which is incorporated herein by reference in its entirety], may be utilized to identify polynucleotide sequences derived from genes that are differentially expressed.

[0136] Differential screening involves the duplicate screening of a cDNA library in which one copy of the library is screened with a total cell cDNA probe corresponding to the mRNA population of one cell type while a duplicate copy of the cDNA library is screened with a total cDNA probe corresponding to the mRNA population of a second cell type. For example, one cDNA probe may correspond to a total cell cDNA probe of a cell type derived from a control subject, while the second cDNA probe may correspond to a total cell cDNA probe of the same cell type derived from an experimental subject. Those clones which hybridize to one probe but not to the other potentially represent clones derived from genes differentially expressed in the cell type of interest in control versus experimental subjects.

[0137] Subtractive hybridization techniques generally involve the isolation of mRNA taken from two different sources, e.g., control and experimental tissue, the hybridization of the mRNA or single-stranded cDNA reverse-transcribed from the isolated mRNA, and the removal of all hybridized, and therefore double-stranded, sequences. The remaining non-hybridized, single-stranded cDNAs, potentially represent clones derived from genes that are differentially expressed in the two mRNA sources. Such single-stranded cDNAs are then used as the starting material for the construction of a library comprising clones derived from differentially expressed genes.

[0138] The differential display technique describes a procedure, utilizing the well known polymerase chain reaction (PCR), the experimental embodiment set forth in Mullis, K. B., 1987, U.S. Pat. No. 4,683,202 which allows for the identification of sequences derived from genes which are differentially expressed. First, isolated RNA is reverse-transcribed into single-stranded cDNA, utilizing standard techniques which are well known to those of skill in the art. Primers for the reverse transcriptase reaction may include, but are not limited to, oligo dT-containing primers, preferably of the reverse primer type of oligonucleotide described below. Next, this technique uses pairs of PCR primers, as described below, which allow for the amplification of clones representing a random subset of the RNA transcripts present within any given cell. Utilizing different pairs of primers allows each of the mRNA transcripts present in a cell to be amplified. Among such amplified transcripts may be identified those which have been produced from differentially expressed genes.

[0139] The reverse oligonucleotide primer of the primer pairs may contain an oligo dT stretch of nucleotides, preferably eleven nucleotides long, at its 5' end, which hybridizes to the poly(A) tail of mRNA or to the complement of a cDNA reverse transcribed from an mRNA poly(A) tail. Second, in order to increase the specificity of the reverse primer, the primer may contain one or more, preferably two, additional nucleotides at its 3' end. Because, statistically, only a subset of the mRNA derived sequences present in the sample of interest will hybridize to such primers, the additional nucleotides allow the primers to amplify only a subset of the mRNA derived sequences present in the sample of interest. This is preferred in that it allows more accurate and complete visualization and characterization of each of the bands representing amplified sequences.

[0140] The forward primer may contain a nucleotide sequence expected, statistically, to have the ability to hybridize to cDNA sequences derived from the tissues of interest. The nucleotide sequence may be an arbitrary one, and the length of the forward oligonucleotide primer may range from about 9 to about 13 nucleotides, with about 10 nucleotides being preferred. Arbitrary primer sequences cause the lengths of the amplified partial cDNAs produced to be variable, thus allowing different clones to be separated by using standard denaturing sequencing gel electrophoresis. PCR reaction conditions should be chosen which optimize amplified product yield and specificity, and, additionally, produce amplified products of lengths which may be resolved utilizing standard gel electrophoresis techniques. Such reaction conditions are well known to those of skill in the art, and important reaction parameters include, for example, length and nucleotide sequence of oligonucleotide primers as discussed above, and annealing and elongation step temperatures and reaction times. The pattern of clones resulting from the reverse transcription and amplification of the mRNA of two different cell types is displayed via sequencing gel electrophoresis and compared. Differences in the two banding patterns indicate potentially differentially expressed genes.

[0141] When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Randomly-primed libraries are preferable, in that they will contain more sequences which contain the

5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d (T) library does not yield a full-length cDNA. Genomic libraries can be useful for extension of sequence into 5' non-transcribed regulatory regions.

[0142] Commercially available capillary electrophoresis systems can be used to analyze the size or confirm the nucleotide sequence of PCR or sequencing products. For example, capillary sequencing can employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity can be converted to electrical signal using appropriate software (e.g. GENOTYPER and Sequence NAVIGATOR, Perkin Elmer, ABI), and the entire process from loading of samples to computer analysis and electronic data display can be computer controlled. Capillary electrophoresis is especially preferable for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

[0143] Once potentially differentially expressed gene sequences have been identified via bulk techniques such as, for example, those described above, the differential expression of such putatively differentially expressed genes should be corroborated. Corroboration may be accomplished via, for example, such well known techniques as Northern analysis and/or RT-PCR. Upon corroboration, the differentially expressed genes may be further characterized, and may be identified as target and/or marker genes, as discussed, below.

[0144] Also, amplified sequences of differentially expressed genes obtained through, for example, differential display may be used to isolate full length clones of the corresponding gene. The full length coding portion of the gene may readily be isolated, without undue experimentation, by molecular biological techniques well known in the art. For example, the isolated differentially expressed amplified fragment may be labeled and used to screen a cDNA library. Alternatively, the labeled fragment may be used to screen a genomic library.

[0145] An analysis of the tissue distribution of the mRNA produced by the identified genes may be conducted, utilizing standard techniques well known to those of skill in the art. Such techniques may include, for example, Northern analyses and RT-PCR. Such analyses provide information as to whether the identified genes are expressed in tissues expected to contribute to breast cancer. Such analyses may also provide quantitative information regarding steady state mRNA regulation, yielding data concerning which of the identified genes exhibits a high level of regulation in, preferably, tissues which may be expected to contribute to breast cancer.

[0146] Such analyses may also be performed on an isolated cell population of a particular cell type derived from a given tissue. Additionally, standard in situ hybridization techniques may be utilized to provide information regarding which cells within a given tissue express the identified gene. Such analyses may provide information regarding the biological function of an identified gene relative to breast cancer in instances wherein only a subset of the cells within the tissue is thought to be relevant to breast cancer.

Identification of co-amplified genes

[0147] Genes involved in genomic alterations (amplifications, insertions, translocations, deletions, etc.) are identified by PCR-based karyotyping in combination with database analysis. Of particular interest are gene amplifications, which account for gene copy numbers >2 per cell. Gene copy number and gene expression of the respective genes often correlates. Therefore clusters of genes being simultaneously overexpressed due to gene amplifications can be identified by expression analysis via DNA-chip technologies or quantitative RTPCR. For example, the altered expression of genes due to increased or decreased gene copy numbers can be determined by GeneArray™ technologies from Affymetrix or qRT-PCR with the TaqMan or iCycler Systems. Moreover combination of RNA with DNA analytic enables highly parallel and automated characterization of multiple genomic regions of variable length with high resolution in tissue or single cell samples. Furthermore these assays enable the correlation of gene transcription relative to gene copy number of target genes. As there is not necessarily a linear correlation of expression level and gene copy number and as there are synergistic or antagonistic effects in certain gene clusters, the identification on the RNA-level is easier and probably more relevant for the biological outcome of the alterations especially in tumor tissue.

Detection of co-amplified genes in malignant neoplasia

[0148] Chromosomal changes are commonly detected by FISH (=Fluorescence-In-Situ-Hybridization) and CGH (=Comparative Genomic Hybridization). For quantification of genomic regions genes or intergenic regions can be used. Such quantification measures the relative abundance of multiple genes with respect to each other (e.g. target gene vs. centromeric region or housekeeping genes). Changes in relative abundance can be detected in paraffin-embedded material even after extraction of RNA or genomic DNA. Measurement of genomic DNA has advantages compared to RNA-analysis due to the stability of DNA, which accounts for the possibility to perform also retrospective studies and offers multiple internal controls (genes not being altered, amplified or deleted) for standardization and exact calculations. Moreover, PCR-analysis of genomic DNA offers the advantage to investigate intergenic, highly variable regions

or combinations of SNPs (=Single Nucleotide Polymorphisms), RFLPs, VNTRs and STRs (in general polymorphic markers). Determination of SNPs or polymorphic markers within defined genomic regions (e.g. SNP analysis by "Pyrosequencing™") has impact on the phenotype of the genomic alterations. For example it is of advantage to determine combinations of polymorphisms or haplotypes in order to characterize the biological potential of genes being part of amplified alleles. Of particular interest are polymorphic markers in breakpoint regions, coding regions or regulatory regions of genes or intergenic regions. By determining predictive haplotypes with defined biological or clinical outcome it is possible to establish diagnostic and prognostic assays with non-tumor samples from patients. Depending on whether preferably one allele or both alleles to same extent are amplified (= linear or non-linear amplifications) haplotypes can be determined. Overrepresentation of specific polymorphic markers combinations in cells or tissues with gene amplifications facilitates haplotype determination, as e.g. combinations of heterozygous polymorphic markers in nucleic acids isolated from normal tissues, body fluids or biological samples of one patient become almost homozygous in neoplastic tissue of the very same patient. This "gain of homozygosity" corresponds to the measurement of altered genomic region due to amplification events and is suitable for identification of "gain of function"- alterations in tumors, which result in e.g. oncogenic or growth promoting activities. In contrast, the detection of "losses of heterozygosity" is used for identification of anti-oncogenes, gate keeper genes or checkpoint genes, that suppress oncogenic activities and negatively regulate cellular growth processes. This intrinsic difference clearly opposes the impact of the respective genomic regions for tumor development and emphasizes the significance of "gain of homozygosity" measurements disclosed in this invention. In addition to the analyses on SNPs, a comparative approach of blood leucocyte DNA and tumor DNA based on VNTR detection can reveal the existence of a formerly described ARCHEON. SNP and VNTR sequences and primer sets most suitable for detection of the ARCHEON at 17q11-21 are disclosed in Table 4 and Table 6. Detection, quantification and sizing of such polymorphic markers can be achieved by methods known to those with skill in the art. In one embodiment of this invention we disclose the comparative measurement of amount and size of any of the disclosed VNTRs (Table 6) by PCR amplification and capillary electrophoresis. PCR can be carried out by standard protocols favorably in a linear amplification range (low cycle number) and detection by CE should be carried out by suppliers protocols (e.g. Agilent). More favorably the detection of the VNTRs disclosed in Table 6 can be carried out in a multiplex fashion, utilizing a variety of labeled primers (e.g. fluorescent, radioactive, bioactive) and a suitable CE detection system (e.g. ABI 310). However the detection can also be performed on slab gels consisting of highly concentrated agarose or polyacrylamide with a monochromal DNA stain. Enhancement of resolution can be achieved by appropriate primer design and length variation to give best results in multiplex PCR.

[0149] It is also of interest to determine covalent modifications of DNA (e.g. methylation) or the associated chromatin (e.g. acetylation or methylation of associated proteins) within the altered genomic regions, that have impact on transcriptional activity of the genes. In general, by measuring multiple, short sequences (60-300 bp) these techniques enable high-resolution analysis of target regions, which cannot be obtained by conventional methods such as FISH analytic (2-100 kb). Moreover the PCR-based DNA analysis techniques offer advantages with regard to sensitivity, specificity, multiplexing, time consumption and low amount of patient material required. These techniques can be optimized by combination with microdissection or macrodissection to obtain purer starting material for analysis.

Extending Polynucleotides

[0150] In one embodiment of such a procedure for the identification and cloning of full length gene sequences, RNA may be isolated, following standard procedures, from an appropriate tissue or cellular source. A reverse transcription reaction may then be performed on the RNA using an oligonucleotide primer complementary to the mRNA that corresponds to the amplified fragment, for the priming of first strand synthesis. Because the primer is anti-parallel to the mRNA, extension will proceed toward the 5' end of the mRNA. The resulting RNA hybrid may then be "tailed" with guanines using a standard terminal transferase reaction, the hybrid may be digested with RNase H, and second strand synthesis may then be primed with a poly-C primer. Using the two primers, the 5' portion of the gene is amplified using PCR. Sequences obtained may then be isolated and recombined with previously isolated sequences to generate a full-length cDNA of the differentially expressed genes of the invention. For a review of cloning strategies and recombinant DNA techniques, see e.g., Sambrook et al., (77); and Ausubel et al., (78).

[0151] Various PCR-based methods can be used to extend the polynucleotide sequences disclosed herein to detect upstream sequences such as promoters and regulatory elements. For example, restriction site PCR uses universal primers to retrieve unknown sequence adjacent to a known locus [Sarkar, 1993, (82)]. Genomic DNA is first amplified in the presence of a primer to a linker sequence and a primer specific to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

[0152] Inverse PCR also can be used to amplify or extend sequences using divergent primers based on a known region [Triglia et al., 1988, (83)]. Primers can be designed using commercially available software, such as OLIGO 4.06

Primer Analysis software (National Biosciences Inc., Plymouth, Minn.), to be e.g. 2230 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68-72 °C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

[0153] Another method which can be used is capture PCR, which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA [Lagerstrom et al., 1991, (84)]. In this method, multiple restriction enzyme digestions and ligations also can be used to place an engineered double-stranded sequence into an unknown fragment of the DNA molecule before performing PCR.

[0154] Additionally, PCR, nested primers, and PROMOTERFINDER libraries (CLONTECH, Palo Alto, Calif.) can be used to walk genomic DNA (CLONTECH, Palo Alto, Calif.). This process avoids the need to screen libraries and is useful in finding intron/exon junctions.

[0155] The sequences of the identified genes may be used, utilizing standard techniques, to place the genes onto genetic maps, e.g., mouse [Copeland & Jenkins, 1991, (85)] and human genetic maps [Cohen, et al., 1993, (86)]. Such mapping information may yield information regarding the genes' importance to human disease by, for example, identifying genes which map near genetic regions to which known genetic breast cancer tendencies map.

Identification of polynucleotide variants and homologues or splice variants

[0156] Variants and homologues of the "BREAST CANCER GENE" polynucleotides described above also are "BREAST CANCER GENE" polynucleotides. Typically, homologous "BREAST CANCER GENE" polynucleotide sequences can be identified by hybridization of candidate polynucleotides to known "BREAST CANCER GENE" polynucleotides under stringent conditions, as is known in the art. For example, using the following wash conditions: 2 X SSC (0.3 M NaCl, 0.03 M sodium citrate, pH 7.0), 0.1% SDS, room temperature twice, 30 minutes each; then 2 X SSC, 0.1% SDS, 50 °C once, 30 minutes; then 2 X SSC, room temperature twice, 10 minutes each homologous sequences can be identified which contain at most about 25-30% basepair mismatches. More preferably, homologous polynucleotide strands contain 15-25% basepair mismatches, even more preferably 5-15% basepair mismatches.

[0157] Species homologues of the "BREAST CANCER GENE" polynucleotides disclosed herein also can be identified by making suitable probes or primers and screening cDNA expression libraries from other species, such as mice, monkeys, or yeast. Human variants of "BREAST CANCER GENE" polynucleotides can be identified, for example, by screening human cDNA expression libraries. It is well known that the T_m of a double-stranded DNA decreases by 1-1.5° C with every 1% decrease in homology [Bonner et al., 1973, (87)]. Variants of human "BREAST CANCER GENE" polynucleotides or "BREAST CANCER GENE" polynucleotides of other species can therefore be identified by hybridizing a putative homologous "BREAST CANCER GENE" polynucleotide with a polynucleotide having a nucleotide sequence of one of the sequences of the SEQ ID NO: 1 to 26 or 53 to 75 or the complement thereof to form a test hybrid. The melting temperature of the test hybrid is compared with the melting temperature of a hybrid comprising polynucleotides having perfectly complementary nucleotide sequences, and the number or percent of basepair mismatches within the test hybrid is calculated.

[0158] Nucleotide sequences which hybridize to "BREAST CANCER GENE" polynucleotides or their complements following stringent hybridization and/or wash conditions also are "BREAST CANCER GENE" polynucleotides. Stringent wash conditions are well known and understood in the art and are disclosed, for example, in Sambrook et al., (77). Typically, for stringent hybridization conditions a combination of temperature and salt concentration should be chosen that is approximately 12-20° C below the calculated T_m of the hybrid under study. The T_m of a hybrid between a "BREAST CANCER GENE" polynucleotide having a nucleotide sequence of one of the sequences of the SEQ ID NO: 1 to 26 or 53 to 75 or the complement thereof and a polynucleotide sequence which is at least about 50, preferably about 75, 90, 96, or 98% identical to one of those nucleotide sequences can be calculated, for example, using the equation below [Bolton and McCarthy, 1962, (88)]:

$$T_m = 81.5^\circ\text{C} - 16.6(\log_{10}[\text{Na}^+]) + 0.41(\%G + C) - 0.63(\%\text{formamide}) - 600/1,$$

where 1 = the length of the hybrid in basepairs.

[0159] Stringent wash conditions include, for example, 4 X SSC at 65° C, or 50% formamide, 4 X SSC at 28° C, or 0.5 X SSC, 0.1% SDS at 65° C. Highly stringent wash conditions include, for example, 0.2 X SSC at 65° C.

[0160] The biological function of the identified genes may be more directly assessed by utilizing relevant *in vivo* and *in vitro* systems. *In vivo* systems may include, but are not limited to, animal systems which naturally exhibit breast cancer predisposition, or ones which have been engineered to exhibit such symptoms, including but not limited to the apoE-deficient malignant neoplasia mouse model [Plump et al., 1992, (89)].

[0161] Splice variants derived from the same genomic region, encoded by the same pre mRNA can be identified by

hybridization conditions described above for homology search. The specific characteristics of variant proteins encoded by splice variants of the same pre transcript may differ and can also be assayed as disclosed. A "BREAST CANCER GENE" polynucleotide having a nucleotide sequence of one of the sequences of the SEQ ID NO: 1 to 26 or 53 to 75 or the complement thereof may therefor differ in parts of the entire sequence as presented for SEQ ID NO: 60 and the encoded splice variants SEQ ID NO: 61 to 66. These refer to individual proteins SEQ ID NO: 83 to 89. The prediction of splicing events and the identification of the utilized acceptor and donor sites within the pre mRNA can be computed (e.g. Software Package GRAIL or GenomeSCAN) and verified by PCR method by those with skill in the art.

Antisense oligonucleotides

[0162] Antisense oligonucleotides are nucleotide sequences which are complementary to a specific DNA or RNA sequence. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form complexes and block either transcription or translation. Preferably, an antisense oligonucleotide is at least 6 nucleotides in length, but can be at least 7, 8, 10, 12, 15, 20, 25, 30, 35, 40, 45, or 50 or more nucleotides long. Longer sequences also can be used. Antisense oligonucleotide molecules can be provided in a DNA construct and introduced into a cell as described above to decrease the level of "BREAST CANCER GENE" gene products in the cell.

[0163] Antisense oligonucleotides can be deoxyribonucleotides, ribonucleotides, peptide nucleic acids (PNAs; described in U.S. Pat. No. 5,714,331), locked nucleic acids (LNAs; described in WO 99/12826), or a combination of them. Oligonucleotides can be synthesized manually or by an automated synthesizer, by covalently linking the 5' end of one nucleotide with the 3' end of another nucleotide with non-phosphodiester internucleotide linkages such as alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, alkylphosphonates, phosphoramidates, phosphate esters, carbamates, acetamides, carboxymethyl esters, carbonates, and phosphate triesters [Brown, 1994, (126); Sonveaux, 1994, (127) and Uhlmann et al., 1990, (128)].

[0164] Modifications of "BREAST CANCER GENE" expression can be obtained by designing antisense oligonucleotides which will form duplexes to the control, 5', or regulatory regions of the "BREAST CANCER GENE". Oligonucleotides derived from the transcription initiation site, e.g., between positions 10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or chaperons. Therapeutic advances using triplex DNA have been described in the literature [Gee et al., 1994, (129)]. An antisense oligonucleotide also can be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

[0165] Precise complementarity is not required for successful complex formation between an antisense oligonucleotide and the complementary sequence of a "BREAST CANCER GENE" polynucleotide. Antisense oligonucleotides which comprise, for example, 2, 3, 4, or 5 or more stretches of contiguous nucleotides which are precisely complementary to a "BREAST CANCER GENE" polynucleotide, each separated by a stretch of contiguous nucleotides which are not complementary to adjacent "BREAST CANCER GENE" nucleotides, can provide sufficient targeting specificity for "BREAST CANCER GENE" mRNA. Preferably, each stretch of complementary contiguous nucleotides is at least 4, 5, 6, 7, or 8 or more nucleotides in length. Non-complementary intervening sequences are preferably 1, 2, 3, or 4 nucleotides in length. One skilled in the art can easily use the calculated melting point of an antisense-sense pair to determine the degree of mismatching which will be tolerated between a particular antisense oligonucleotide and a particular "BREAST CANCER GENE" polynucleotide sequence.

[0166] Antisense oligonucleotides can be modified without affecting their ability to hybridize to a "BREAST CANCER GENE" polynucleotide. These modifications can be internal or at one or both ends of the antisense molecule. For example, inter-nucleoside phosphate linkages can be modified by adding cholesteryl or diamine moieties with varying numbers of carbon residues between the amino groups and terminal ribose. Modified bases and/or sugars, such as arabinose instead of ribose, or a 3', 5' substituted oligonucleotide in which the 3' hydroxyl group or the 5' phosphate group are substituted, also can be employed in a modified antisense oligonucleotide. These modified oligonucleotides can be prepared by methods well known in the art [Agrawal et al., 1992, (130); Uhlmann et al., 1987, (131) and Uhlmann et al., (128)].

Ribozymes

[0167] Ribozymes are RNA molecules with catalytic activity [Cech, 1987, (132); Cech, 1990, (133) and Couture & Stinchcomb, 1996, (134)]. Ribozymes can be used to inhibit gene function by cleaving an RNA sequence, as is known in the art (e.g., Haseloff et al., U.S. Patent 5,641,673). The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Examples include engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of specific nucleotide sequences.

[0168] The transcribed sequence of a "BREAST CANCER GENE" can be used to generate ribozymes which will specifically bind to mRNA transcribed from a "BREAST CANCER GENE" genomic locus. Methods of designing and constructing ribozymes which can cleave other RNA molecules in trans in a highly sequence specific manner have been developed and described in the art [Haseloff et al., 1988, (135)]. For example, the cleavage activity of ribozymes can be targeted to specific RNAs by engineering a discrete "hybridization" region into the ribozyme. The hybridization region contains a sequence complementary to the target RNA and thus specifically hybridizes with the target [see, for example, Gerlach et al., EP 0 321201].

[0169] Specific ribozyme cleavage sites within a "BREAST CANCER GENE" RNA target can be identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target RNA containing the cleavage site can be evaluated for secondary structural features which may render the target inoperable. Suitability of candidate "BREAST CANCER GENE" RNA targets also can be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays. Longer complementary sequences can be used to increase the affinity of the hybridization sequence for the target. The hybridizing and cleavage regions of the ribozyme can be integrally related such that upon hybridizing to the target RNA through the complementary regions, the catalytic region of the ribozyme can cleave the target.

[0170] Ribozymes can be introduced into cells as part of a DNA construct. Mechanical methods, such as microinjection, liposome-mediated transfection, electroporation, or calcium phosphate precipitation, can be used to introduce a ribozyme-containing DNA construct into cells in which it is desired to decrease "BREAST CANCER GENE" expression. Alternatively, if it is desired that the cells stably retain the DNA construct, the construct can be supplied on a plasmid and maintained as a separate element or integrated into the genome of the cells, as is known in the art. A ribozyme-encoding DNA construct can include transcriptional regulatory elements, such as a promoter element, an enhancer or UAS element, and a transcriptional terminator signal, for controlling transcription of ribozymes in the cells.

[0171] As taught in Haseloff et al., U.S. Pat. No. 5,641,673, ribozymes can be engineered so that ribozyme expression will occur in response to factors which induce expression of a target gene. Ribozymes also can be engineered to provide an additional level of regulation, so that destruction of mRNA occurs only when both a ribozyme and a target gene are induced in the cells.

Polypeptides

[0172] "BREAST CANCER GENE" polypeptides according to the invention comprise a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by any of the polynucleotide sequences of the SEQ ID NO: 1 to 26 and 53 to 75 or derivatives, fragments, analogues and homologues thereof. A "BREAST CANCER GENE" polypeptide of the invention therefore can be a portion, a full-length, or a fusion protein comprising all or a portion of a "BREAST CANCER GENE" polypeptide.

Protein Purification

[0173] "BREAST CANCER GENE" polypeptides can be purified from any cell which expresses the enzyme, including host cells which have been transfected with "BREAST CANCER GENE" expression constructs. Breast tissue is an especially useful source of "BREAST CANCER GENE" polypeptides. A purified "BREAST CANCER GENE" polypeptide is separated from other compounds which normally associate with the "BREAST CANCER GENE" polypeptide in the cell, such as certain proteins, carbohydrates, or lipids, using methods well-known in the art. Such methods include, but are not limited to, size exclusion chromatography, ammonium sulfate fractionation, ion exchange chromatography, affinity chromatography, and preparative gel electrophoresis. A preparation of purified "BREAST CANCER GENE" polypeptides is at least 80% pure; preferably, the preparations are 90%, 95%, or 99% pure. Purity of the preparations can be assessed by any means known in the art, such as SDS-polyacrylamide gel electrophoresis.

Obtaining Polypeptides

[0174] "BREAST CANCER GENE" polypeptides can be obtained, for example, by purification from human cells, by expression of "BREAST CANCER GENE" polynucleotides, or by direct chemical synthesis.

Biologically Active Variants

[0175] "BREAST CANCER GENE" polypeptide variants which are biologically active, i.e., retain an "BREAST CANCER GENE" activity, also are "BREAST CANCER GENE" polypeptides. Preferably, naturally or non-naturally occurring "BREAST CANCER GENE" polypeptide variants have amino acid sequences which are at least about 60, 65, or 70,

preferably about 75, 80, 85, 90, 92, 94, 96, or 98% identical to the any of the amino acid sequences of the polypeptides of SEQ ID NO: 27 to 52 or 76 to 98 or the polypeptides encoded by any of the polynucleotides of SEQ ID NO: 1 to 26 or 53 to 75 or a fragment thereof. Percent identity between a putative "BREAST CANCER GENE" polypeptide variant and of the polypeptides of SEQ ID NO: 27 to 52 or 76 to 98 or the polypeptides encoded by any of the polynucleotides of SEQ ID NO: 1 to 26 or 53 to 75 or a fragment thereof is determined by conventional methods. [See, for example, Altschul *et al.*, 1986, (90 and Henikoff & Henikoff, 1992, (91)]. Briefly, two amino acid sequences are aligned to optimize the alignment scores using a gap opening penalty of 10, a gap extension penalty of 1, and the "BLOSUM62" scoring matrix of Henikoff & Henikoff, (91).

[0176] Those skilled in the art appreciate that there are many established algorithms available to align two amino acid sequences. The "FASTA" similarity search algorithm of Pearson & Lipman is a suitable protein alignment method for examining the level of identity shared by an amino acid sequence disclosed herein and the amino acid sequence of a putative variant [Pearson & Lipman, 1988, (92), and Pearson, 1990, (93)]. Briefly, FASTA first characterizes sequence similarity by identifying regions shared by the query sequence (e.g., SEQ ID NO: 1 to 26 or 53 to 75) and a test sequence that have either the highest density of identities (if the ktp variable is 1) or pairs of identities (if ktp=2), without considering conservative amino acid substitutions, insertions, or deletions. The ten regions with the highest density of identities are then rescored by comparing the similarity of all paired amino acids using an amino acid substitution matrix, and the ends of the regions are "trimmed" to include only those residues that contribute to the highest score. If there are several regions with scores greater than the "cutoff" value (calculated by a predetermined formula based upon the length of the sequence the ktp value), then the trimmed initial regions are examined to determine whether the regions can be joined to form an approximate alignment with gaps. Finally, the highest scoring regions of the two amino acid sequences are aligned using a modification of the Needleman-Wunsch-Sellers algorithm [Needleman & Wunsch, 1970, (94), and Sellers, 1974, (95)], which allows for amino acid insertions and deletions. Preferred parameters for FASTA analysis are: ktp=1, gap opening penalty=10, gap extension penalty=1, and substitution matrix=BLOSUM62. These parameters can be introduced into a FASTA program by modifying the scoring matrix file ("SMATRIX"), as explained in Appendix 2 of Pearson, (93).

[0177] FASTA can also be used to determine the sequence identity of nucleic acid molecules using a ratio as disclosed above. For nucleotide sequence comparisons, the ktp value can range between one to six, preferably from three to six, most preferably three, with other parameters set as default.

[0178] Variations in percent identity can be due, for example, to amino acid substitutions, insertions, or deletions. Amino acid substitutions are defined as one for one amino acid replacements. They are conservative in nature when the substituted amino acid has similar structural and/or chemical properties. Examples of conservative replacements are substitution of a leucine with an isoleucine or valine, an aspartate with a glutamate, or a threonine with a serine.

[0179] Amino acid insertions or deletions are changes to or within an amino acid sequence. They typically fall in the range of about 1 to 5 amino acids. Guidance in determining which amino acid residues can be substituted, inserted, or deleted without abolishing biological or immunological activity of a "BREAST CANCER GENE" polypeptide can be found using computer programs well known in the art, such as DNASTAR software. Whether an amino acid change results in a biologically active "BREAST CANCER GENE" polypeptide can readily be determined by assaying for "BREAST CANCER GENE" activity, as described for example, in the specific Examples, below. Larger insertions or deletions can also be caused by alternative splicing. Protein domains can be inserted or deleted without altering the main activity of the protein.

Fusion Proteins

[0180] Fusion proteins are useful for generating antibodies against "BREAST CANCER GENE" polypeptide amino acid sequences and for use in various assay systems. For example, fusion proteins can be used to identify proteins which interact with portions of a "BREAST CANCER GENE" polypeptide. Protein affinity chromatography or library-based assays for protein-protein interactions, such as the yeast two-hybrid or phage display systems, can be used for this purpose. Such methods are well known in the art and also can be used as drug screens.

[0181] A "BREAST CANCER GENE" polypeptide fusion protein comprises two polypeptide segments fused together by means of a peptide bond. The first polypeptide segment comprises at least 25, 50, 75, 100, 150, 200, 300, 400, 500, 600, 700 or 750 contiguous amino acids of an amino acid sequence encoded by any polynucleotide sequences of the SEQ ID NO: 1 to 26 or 53 to 75 or of a biologically active variant, such as those described above. The first polypeptide segment also can comprise full-length "BREAST CANCER GENE".

[0182] The second polypeptide segment can be a full-length protein or a protein fragment. Proteins commonly used in fusion protein construction include β -galactosidase, β -glucuronidase, green fluorescent protein (GFP), autofluorescent proteins, including blue fluorescent protein (BFP), glutathione-S-transferase (GST), luciferase, horseradish peroxidase (HRP), and chloramphenicol acetyltransferase (CAT). Additionally, epitope tags are used in fusion protein constructions, including histidine (His) tags, FLAG tags, influenza hemagglutinin (HA) tags, Myc tags, VSV-G tags,

and thioredoxin (Trx) tags. Other fusion constructions can include maltose binding protein (MBP), S-tag, Lex a DNA binding domain (DBD) fusions, GAL4 DNA binding domain fusions, and herpes simplex virus (HSV) BP16 protein fusions. A fusion protein also can be engineered to contain a cleavage site located between the "BREAST CANCER GENE" polypeptide-encoding sequence and the heterologous protein sequence, so that the "BREAST CANCER GENE" polypeptide can be cleaved and purified away from the heterologous moiety.

[0183] A fusion protein can be synthesized chemically, as is known in the art. Preferably, a fusion protein is produced by covalently linking two polypeptide segments or by standard procedures in the art of molecular biology. Recombinant DNA methods can be used to prepare fusion proteins, for example, by making a DNA construct which comprises coding sequences selected from any of the polynucleotide sequences of the SEQ ID NO: 1 to 26 and 53 to 75 in proper reading frame with nucleotides encoding the second polypeptide segment and expressing the DNA construct in a host cell, as is known in the art. Many kits for constructing fusion proteins are available from companies such as Promega Corporation (Madison, WI), Stratagene (La Jolla, CA), CLONTECH (Mountain View, CA), Santa Cruz Biotechnology (Santa Cruz, CA), MBL International Corporation (MIC; Watertown, MA), and Quantum Biotechnologies (Montreal, Canada; 1-888-DNA-KITS).

Identification of Species Homologues

[0184] Species homologues of human a "BREAST CANCER GENE" polypeptide can be obtained using "BREAST CANCER GENE" polypeptide polynucleotides (described below) to make suitable probes or primers for screening cDNA expression libraries from other species, such as mice, monkeys, or yeast, identifying cDNAs which encode homologues of a "BREAST CANCER GENE" polypeptide, and expressing the cDNAs as is known in the art.

Expression of Polynucleotides

[0185] To express a "BREAST CANCER GENE" polynucleotide, the polynucleotide can be inserted into an expression vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art can be used to construct expression vectors containing sequences encoding "BREAST CANCER GENE" polypeptides and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. Such techniques are described, for example, in Sambrook et al., (77) and in Ausubel et al., (78).

[0186] A variety of expression vector/host systems can be utilized to contain and express sequences encoding a "BREAST CANCER GENE" polypeptide. These include, but are not limited to, microorganisms, such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors, insect cell systems infected with virus expression vectors (e.g., baculovirus), plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids), or animal cell systems.

[0187] The control elements or regulatory sequences are those regions of the vector enhancers, promoters, 5' and 3' untranslated regions which interact with host cellular proteins to carry out transcription and translation. Such elements can vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, can be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the BLUESCRIPT phagemid (Stratagene, LaJolla, Calif.) or pSPORT1 plasmid (Life Technologies) and the like can be used. The baculovirus polyhedrin promoter can be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (e.g., heat shock, RUBISCO, and storage protein genes) or from plant viruses (e.g., viral promoters or leader sequences) can be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of a nucleotide sequence encoding a "BREAST CANCER GENE" polypeptide, vectors based on SV40 or EBV can be used with an appropriate selectable marker.

Bacterial and Yeast Expression Systems

[0188] In bacterial systems, a number of expression vectors can be selected depending upon the use intended for the "BREAST CANCER GENE" polypeptide. For example, when a large quantity of the "BREAST CANCER GENE" polypeptide is needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified can be used. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene). In a BLUESCRIPT vector, a sequence encoding the "BREAST CANCER GENE" polypeptide can be ligated into the vector in frame with sequences for the amino terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced. pIN vectors [Van Heeke & Schuster,

(17)] or pGEX vectors (Promega, Madison, Wis.) also can be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST).

[0189] In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems can be designed to include heparin, thrombin, or factor Xa protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

[0190] In the yeast *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH can be used. For reviews, see Ausubel et al., (4) and Grant et al., (18).

Plant and Insect Expression Systems

[0191] If plant expression vectors are used, the expression of sequences encoding "BREAST CANCER GENE" polypeptides can be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV can be used alone or in combination with the omega leader sequence from TMV [Takamatsu, 1987, (96)]. Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters can be used [Coruzzi et al., 1984, (97); Broglie et al., 1984, (98); Winter et al., 1991, (99)]. These constructs can be introduced into plant cells by direct DNA transformation or by pathogen-mediated transfection. Such techniques are described in a number of generally available reviews.

[0192] An insect system also can be used to express a "BREAST CANCER GENE" polypeptide. For example, in one such system *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. Sequences encoding "BREAST CANCER GENE" polypeptides can be cloned into a nonessential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of "BREAST CANCER GENE" polypeptides will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses can then be used to infect *S. frugiperda* cells or *Trichoplusia* larvae in which "BREAST CANCER GENE" polypeptides can be expressed [Engelhard et al., 1994, (100)].

Mammalian Expression Systems

[0193] A number of viral-based expression systems can be used to express "BREAST CANCER GENE" polypeptides in mammalian host cells. For example, if an adenovirus is used as an expression vector, sequences encoding "BREAST CANCER GENE" polypeptides can be ligated into an adenovirus transcription/translation complex comprising the late promoter and tripartite leader sequence. Insertion in a nonessential E1 or E3 region of the viral genome can be used to obtain a viable virus which is capable of expressing a "BREAST CANCER GENE" polypeptide in infected host cells [Logan & Shenk, 1984, (101)]. If desired, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, can be used to increase expression in mammalian host cells.

[0194] Human artificial chromosomes (HACs) also can be used to deliver larger fragments of DNA than can be contained and expressed in a plasmid. HACs of 6M to 10M are constructed and delivered to cells via conventional delivery methods (e.g., liposomes, polycationic amino polymers, or vesicles).

[0195] Specific initiation signals also can be used to achieve more efficient translation of sequences encoding "BREAST CANCER GENE" polypeptides. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding a "BREAST CANCER GENE" polypeptide, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals (including the ATG initiation codon) should be provided. The initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons can be of various origins, both natural and synthetic. The efficiency of expression can be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used [Scharf et al., 1994, (102)].

Host Cells

[0196] A host cell strain can be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed "BREAST CANCER GENE" polypeptide in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Posttranslational processing which cleaves a "prepro" form of the polypeptide also can be used to facilitate correct insertion, folding and/or function. Different host cells which have specific cellular machinery and characteristic mechanisms for Post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WJ38), are available from the American Type Culture Collection (ATCC; 10801 University Boulevard, Manassas, VA 20110-2209) and can be chosen

to ensure the correct modification and processing of the foreign protein.

[0197] Stable expression is preferred for long-term, high-yield production of recombinant proteins. For example, cell lines which stably express "BREAST CANCER GENE" polypeptides can be transformed using expression vectors which contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells can be allowed to grow for 12 days in an enriched medium before they are switched to a selective medium. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced "BREAST CANCER GENE" sequences. Resistant clones of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type [Freshney et al., 1986, (103)].

[0198] Any number of selection systems can be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase [Wigler et al., 1977, (104)] and adenine phosphoribosyltransferase [Lowy et al., 1980, (105)] genes which can be employed in tk- or apt- cells, respectively. Also, aminetalolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, dhfr confers resistance to methotrexate [Wigler et al., 1980, (106)], npt confers resistance to the aminoglycosides, neomycin and G418 [Colbere-Garapin et al., 1981, (107)], and als and pat confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. Additional selectable genes have been described. For example, trpB allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine [Hartman & Mulligan, 1988, (108)]. Visible markers such as anthocyanins, β -glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, can be used to identify transformants and to quantify the amount of transient or stable protein expression attributable to a specific vector system [Rhodes et al., 1995, (109)].

Detecting Expression and gene product

[0199] Although the presence of marker gene expression suggests that the "BREAST CANCER GENE" polynucleotide is also present, its presence and expression may need to be confirmed. For example, if a sequence encoding a "BREAST CANCER GENE" polypeptide is inserted within a marker gene sequence, transformed cells containing sequences which encode a "BREAST CANCER GENE" polypeptide can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding a "BREAST CANCER GENE" polypeptide under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the "BREAST CANCER GENE" polynucleotide.

[0200] Alternatively, host cells which contain a "BREAST CANCER GENE" polynucleotide and which express a "BREAST CANCER GENE" polypeptide can be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridization and protein bioassay or immunoassay techniques which include membrane, solution, or chip-based technologies for the detection and/or quantification of polynucleotide or protein. For example, the presence of a polynucleotide sequence encoding a "BREAST CANCER GENE" polypeptide can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes or fragments or fragments of polynucleotides encoding a "BREAST CANCER GENE" polypeptide. Nucleic acid amplification-based assays involve the use of oligonucleotides selected from sequences encoding a "BREAST CANCER GENE" polypeptide to detect transformants which contain a "BREAST CANCER GENE" polynucleotide.

[0201] A variety of protocols for detecting and measuring the expression of a "BREAST CANCER GENE" polypeptide, using either polyclonal or monoclonal antibodies specific for the polypeptide, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay using monoclonal antibodies reactive to two non-interfering epitopes on a "BREAST CANCER GENE" polypeptide can be used, or a competitive binding assay can be employed. These and other assays are described in Hampton et al., (110) and Maddox et al., (111).

[0202] A wide variety of labels and conjugation techniques are known by those skilled in the art and can be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding "BREAST CANCER GENE" polypeptides include oligo labeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, sequences encoding a "BREAST CANCER GENE" polypeptide can be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and can be used to synthesize RNA probes in vitro by addition of labeled nucleotides and an appropriate RNA polymerase such as T7, T3, or SP6. These procedures can be conducted using a variety of commercially available kits (Amersham Pharmacia Biotech, Promega, and US Biochemical). Suitable reporter molecules or labels which can be used for ease of detection include radionuclides, enzymes, and fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Expression and Purification of Polypeptides

[0203] Host cells transformed with nucleotide sequences encoding a "BREAST CANCER GENE" polypeptide can be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The polypeptide produced by a transformed cell can be secreted or stored intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode "BREAST CANCER GENE" polypeptides can be designed to contain signal sequences which direct secretion of soluble "BREAST CANCER GENE" polypeptides through a prokaryotic or eukaryotic cell membrane or which direct the membrane insertion of membrane-bound "BREAST CANCER GENE" polypeptide.

[0204] As discussed above, other constructions can be used to join a sequence encoding a "BREAST CANCER GENE" polypeptide to a nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). Inclusion of cleavable linker sequences such as those specific for Factor Xa or enterokinase (Invitrogen, San Diego, CA) between the purification domain and the "BREAST CANCER GENE" polypeptide also can be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing a "BREAST CANCER GENE" polypeptide and 6 histidine residues preceding a thiorodoxin or an enterokinase cleavage site. The histidine residues facilitate purification by IMAC (immobilized metal ion affinity chromatography [Porath et al., 1992, (112)], while the enterokinase cleavage site provides a means for purifying the "BREAST CANCER GENE" polypeptide from the fusion protein. Vectors which contain fusion proteins are disclosed in Kroll et al., (113).

Chemical Synthesis

[0205] Sequences encoding a "BREAST CANCER GENE" polypeptide can be synthesized, in whole or in part, using chemical methods well known in the art (see Caruthers et al., (114) and Horn et al., (115). Alternatively, a "BREAST CANCER GENE" polypeptide itself can be produced using chemical methods to synthesize its amino acid sequence, such as by direct peptide synthesis using solid-phase techniques [Merrifield, 1963, (116) and Roberge et al., 1995, (117)]. Protein synthesis can be performed using manual techniques or by automation. Automated synthesis can be achieved, for example, using Applied Biosystems 431 A Peptide Synthesizer (Perkin Elmer). Optionally, fragments of "BREAST CANCER GENE" polypeptides can be separately synthesized and combined using chemical methods to produce a full-length molecule.

[0206] The newly synthesized peptide can be substantially purified by preparative high performance liquid chromatography [Creighton, 1983, (118)]. The composition of a synthetic "BREAST CANCER GENE" polypeptide can be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure; see Creighton, (118). Additionally, any portion of the amino acid sequence of the "BREAST CANCER GENE" polypeptide can be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins to produce a variant polypeptide or a fusion protein.

Production of Altered Polypeptides

[0207] As will be understood by those of skill in the art, it may be advantageous to produce "BREAST CANCER GENE" polypeptide-encoding nucleotide sequences possessing non-natural occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce an RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

[0208] The nucleotide sequences disclosed herein can be engineered using methods generally known in the art to alter "BREAST CANCER GENE" polypeptide-encoding sequences for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the polypeptide or mRNA product. DNA shuffling by random fragmentation and PCR re-assembly of gene fragments and synthetic oligonucleotides can be used to engineer the nucleotide sequences. For example, site-directed mutagenesis can be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, introduce mutations, and so forth.

Predictive, Diagnostic and Prognostic Assays

[0209] The present invention provides method for determining whether a subject is at risk for developing malignant neoplasia and breast cancer in particular by detecting one of the disclosed polynucleotide markers comprising any of

the polynucleotides sequences of the SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19 or 21 to 26 or 53 to 75 and/or the polypeptide markers encoded thereby or polypeptide markers comprising any of the polypeptide sequences of the SEQ ID NO: 28 to 32, 34, 35, 37 to 42, 44, 45 or 47 to 52 or 76 to 98 or at least 2 of the disclosed polynucleotides selected from SEQ ID NO: 1 to 26 and 53 to 75 or the at least 2 of the disclosed polypeptides selected from SEQ ID NO: 28 to 32 and 76 to 98 for malignant neoplasia and breast cancer in particular.

[0210] In clinical applications, biological samples can be screened for the presence and/or absence of the biomarkers identified herein. Such samples are for example needle biopsy cores, surgical resection samples, or body fluids like serum, thin needle nipple aspirates and urine. For example, these methods include obtaining a biopsy, which is optionally fractionated by cryostat sectioning to enrich diseased cells to about 80% of the total cell population. In certain embodiments, polynucleotides extracted from these samples may be amplified using techniques well known in the art. The expression levels of selected markers detected would be compared with statistically valid groups of diseased and healthy samples.

[0211] In one embodiment the diagnostic method comprises determining whether a subject has an abnormal mRNA and/or protein level of the disclosed markers, such as by Northern blot analysis, reverse transcription-polymerase chain reaction (RT-PCR), in situ hybridization, immunoprecipitation, Western blot hybridization, or immunohistochemistry. According to the method, cells are obtained from a subject and the levels of the disclosed biomarkers, protein or mRNA level, is determined and compared to the level of these markers in a healthy subject. An abnormal level of the biomarker polypeptide or mRNA levels is likely to be indicative of malignant neoplasia such as breast cancer.

[0212] In another embodiment the diagnostic method comprises determining whether a subject has an abnormal DNA content of said genes or said genomic loci, such as by Southern blot analysis, dot blot analysis, fluorescence or colorimetric In Situ hybridization, comparative genomic hybridization, genotyping by VNTR, STS-PCR or quantitative PCR. In general these assays comprise the usage of probes from representative genomic regions. The probes contain at least parts of said genomic regions or sequences complementary or analogous to said regions. In particular intra- or intergenic regions of said genes or genomic regions. The probes can consist of nucleotide sequences or sequences of analogous functions (e.g. PNAs, Morpholino oligomers) being able to bind to target regions by hybridization. In general genomic regions being altered in said patient samples are compared with unaffected control samples (normal tissue from the same or different patients, surrounding unaffected tissue, peripheral blood) or with genomic regions of the same sample that don't have said alterations and can therefore serve as internal controls. In a preferred embodiment regions located on the same chromosome are used. Alternatively, gonosomal regions and/or regions with defined varying amount in the sample are used. In one favored embodiment the DNA content, structure, composition or modification is compared that lie within distinct genomic regions. Especially favored are methods that detect the DNA content of said samples, where the amount of target regions are altered by amplification and/or deletions. In another embodiment the target regions are analyzed for the presence of polymorphisms (e.g. Single Nucleotide Polymorphisms or mutations) that affect or predispose the cells in said samples with regard to clinical aspects, being of diagnostic, prognostic or therapeutic value. Preferably, the identification of sequence variations is used to define haplotypes that result in characteristic behavior of said samples with said clinical aspects.

[0213] The following examples of genes in 17q12-21.2 are offered by way of illustration, not by way of limitation.

[0214] One embodiment of the invention is a method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least 10, at least 5, or at least 4, or at least 3 and more preferably at least 2 markers whereby the markers are genes and fragments thereof and/or genomic nucleic acid sequences that are located on one chromosomal region which is altered in malignant neoplasia.

[0215] One further embodiment of the invention is method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least 10, at least 5, or at least 4, or at least 3 and more preferably at least 2 markers whereby the markers (a) are genes and fragments thereof and/or genomic nucleic acid sequences that are located on one or more chromosomal region(s) which is/are altered in malignant neoplasia and (b) functionally interact as (i) receptor and ligand or (ii) members of the same signal transduction pathway or (iii) members of synergistic signal transduction pathways or (iv) members of antagonistic signal transduction pathways or (v) transcription factor and transcription factor binding site.

[0216] In one embodiment, the method for the prediction, diagnosis or prognosis of malignant neoplasia and breast cancer in particular is done by the detection of:

(a) polynucleotide selected from the polynucleotides of the SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

(b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3;

(c) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the

generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3;

(d) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);

in a biological sample comprising the following steps: hybridizing any polynucleotide or analogous oligomer specified in (a) to (d) to a polynucleotide material of a biological sample, thereby forming a hybridization complex; and detecting said hybridization complex.

[0217] In another embodiment the method for the prediction, diagnosis or prognosis of malignant neoplasia is done as just described but, wherein before hybridization, the polynucleotide material of the biological sample is amplified.

[0218] In another embodiment the method for the diagnosis or prognosis of malignant neoplasia and breast cancer in particular is done by the detection of:

(a) a polynucleotide selected from the polynucleotides of the SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

(b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3;

(c) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3;

(d) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);

(e) a polypeptide encoded by a polynucleotide sequence specified in (a) to (d)

(f) a polypeptide comprising any polypeptide of SEQ ID NO: 28 to 32, 34, 35, 37 to 42, 44, 45, 47 to 52 or 76 to 98;

comprising the steps of contacting a biological sample with a reagent which specifically interacts with the polynucleotide specified in (a) to (d) or the polypeptide specified in (e).

DNA array technology

[0219] In one embodiment, the present invention also provides a method wherein polynucleotide probes are immobilized on a DNA chip in an organized array. Oligonucleotides can be bound to a solid support by a variety of processes, including lithography. For example a chip can hold up to 4100,00 oligonucleotides (GeneChip, Affymetrix). The present invention provides significant advantages over the available tests for malignant neoplasia, such as breast cancer, because it increases the reliability of the test by providing an array of polynucleotide markers on a single chip.

[0220] The method includes obtaining a biopsy of an affected person, which is optionally fractionated by cryostat sectioning to enrich diseased cells to about 80% of the total cell population and the use of body fluids such as serum or urine, serum or cell containing liquids (e.g. derived from fine needle aspirates). The DNA or RNA is then extracted, amplified, and analyzed with a DNA chip to determine the presence of absence of the marker polynucleotide sequences. In one embodiment, the polynucleotide probes are spotted onto a substrate in a two-dimensional matrix or array, samples of polynucleotides can be labeled and then hybridized to the probes. Double-stranded polynucleotides, comprising the labeled sample polynucleotides bound to probe polynucleotides, can be detected once the unbound portion of the sample is washed away.

[0221] The probe polynucleotides can be spotted on substrates including glass, nitrocellulose, etc. The probes can be bound to the substrate by either covalent bonds or by non-specific interactions, such as hydrophobic interactions. The sample polynucleotides can be labeled using radioactive labels, fluorophores, chromophores, etc. Techniques for constructing arrays and methods of using these arrays are described in EP 0 799 897; WO 97/29212; WO 97/27317; EP 0 785 280; WO 97/02357; U.S. Pat. No. 5,593,839; U.S. Pat. No. 5,578,832; EP 0 728 520; U.S. Pat. No. 5,599,695; EP 0 721 016; U.S. Pat. No. 5,556,752; WO 95/22058; and U.S. Pat. No. 5,631,734.

[0222] Further, arrays can be used to examine differential expression of genes and can be used to determine gene function. For example, arrays of the instant polynucleotide sequences can be used to determine if any of the polynucleotide sequences are differentially expressed between normal cells and diseased cells, for example. High expression

of a particular message in a diseased sample, which is not observed in a corresponding normal sample, can indicate a breast cancer specific protein.

[0223] Accordingly, in one aspect, the invention provides probes and primers that are specific to the unique polynucleotide markers disclosed herein.

[0224] In one embodiment, the method comprises using a polynucleotide probe to determine the presence of malignant or breast cancer cells in particular in a tissue from a patient. Specifically, the method comprises:

1) providing a polynucleotide probe comprising a nucleotide sequence at least 12 nucleotides in length, preferably at least 15 nucleotides, more preferably, 25 nucleotides, and most preferably at least 40 nucleotides, and up to all or nearly all of the coding sequence which is complementary to a portion of the coding sequence of a polynucleotide selected from the polynucleotides of SEQ ID NO: 1 to 26 and 53 to 75 or a sequence complementary thereto and is

2) differentially expressed in malignant neoplasia, such as breast cancer;

3) obtaining a tissue sample from a patient with malignant neoplasia;

4) providing a second tissue sample from a patient with no malignant neoplasia;

5) contacting the polynucleotide probe under stringent conditions with RNA of each of said first and second tissue samples (e.g., in a Northern blot or in situ hybridization assay); and

6) comparing (a) the amount of hybridization of the probe with RNA of the first tissue sample, with (b) the amount of hybridization of the probe with RNA of the second tissue sample;

wherein a statistically significant difference in the amount of hybridization with the RNA of the first tissue sample as compared to the amount of hybridization with the RNA of the second tissue sample is indicative of malignant neoplasia and breast cancer in particular in the first tissue sample.

Data analysis methods

[0225] Comparison of the expression levels of one or more "BREAST CANCER GENES" with reference expression levels, e.g., expression levels in diseased cells of breast cancer or in normal counterpart cells, is preferably conducted using computer systems. In one embodiment, expression levels are obtained in two cells and these two sets of expression levels are introduced into a computer system for comparison. In a preferred embodiment, one set of expression levels is entered into a computer system for comparison with values that are already present in the computer system, or in computer-readable form that is then entered into the computer system.

[0226] In one embodiment, the invention provides a computer readable form of the gene expression profile data of the invention, or of values corresponding to the level of expression of at least one "BREAST CANCER GENE" in a diseased cell. The values can be mRNA expression levels obtained from experiments, e.g., microarray analysis. The values can also be mRNA levels normalised relative to a reference gene whose expression is constant in numerous cells under numerous conditions, e.g., GAPDH. In other embodiments, the values in the computer are ratios of, or differences between, normalized or non-normalized mRNA levels in different samples.

[0227] The gene expression profile data can be in the form of a table, such as an Excel table. The data can be alone, or it can be part of a larger database, e.g., comprising other expression profiles. For example, the expression profile data of the invention can be part of a public database. The computer readable form can be in a computer. In another embodiment, the invention provides a computer displaying the gene expression profile data.

[0228] In one embodiment, the invention provides a method for determining the similarity between the level of expression of one or more "BREAST CANCER GENES" in a first cell, e.g., a cell of a subject, and that in a second cell, comprising obtaining the level of expression of one or more "BREAST CANCER GENES" in a first cell and entering these values into a computer comprising a database including records comprising values corresponding to levels of expression of one or more "BREAST CANCER GENES" in a second cell, and processor instructions, e.g., a user interface, capable of receiving a selection of one or more values for comparison purposes with data that is stored in the computer. The computer may further comprise a means for converting the comparison data into a diagram or chart or other type of output.

[0229] In another embodiment, values representing expression levels of "BREAST CANCER GENES" are entered into a computer system, comprising one or more databases with reference expression levels obtained from more than one cell. For example, the computer comprises expression data of diseased and normal cells. Instructions are provided to the computer, and the computer is capable of comparing the data entered with the data in the computer to determine

whether the data entered is more similar to that of a normal cell or of a diseased cell.

[0230] In another embodiment, the computer comprises values of expression levels in cells of subjects at different stages of breast cancer, and the computer is capable of comparing expression data entered into the computer with the data stored, and produce results indicating to which of the expression profiles in the computer, the one entered is most similar, such as to determine the stage of breast cancer in the subject.

[0231] In yet another embodiment, the reference expression profiles in the computer are expression profiles from cells of breast cancer of one or more subjects, which cells are treated *in vivo* or *in vitro* with a drug used for therapy of breast cancer. Upon entering of expression data of a cell of a subject treated *in vitro* or *in vivo* with the drug, the computer is instructed to compare the data entered to the data in the computer, and to provide results indicating whether the expression data input into the computer are more similar to those of a cell of a subject that is responsive to the drug or more similar to those of a cell of a subject that is not responsive to the drug. Thus, the results indicate whether the subject is likely to respond to the treatment with the drug or unlikely to respond to it.

[0232] In one embodiment, the invention provides a system that comprises a means for receiving gene expression data for one or a plurality of genes; a means for comparing the gene expression data from each of said one or plurality of genes to a common reference frame, and a means for presenting the results of the comparison. This system may further comprise a means for clustering the data.

[0233] In another embodiment, the invention provides a computer program for analyzing gene expression data comprising (i) a computer code that receives as input gene expression data for a plurality of genes and (ii) a computer code that compares said gene expression data from each of said plurality of genes to a common reference frame.

[0234] The invention also provides a machine-readable or computer-readable medium including program instructions for performing the following steps: (i) comparing a plurality of values corresponding to expression levels of one or more genes characteristic of breast cancer in a query cell with a database including records comprising reference expression or expression profile data of one or more reference cells and an annotation of the type of cell; and (ii) indicating to which cell the query cell is most similar based on similarities of expression profiles. The reference cells can be cells from subjects at different stages of breast cancer. The reference cells can also be cells from subjects responding or not responding to a particular drug treatment and optionally incubated *in vitro* or *in vivo* with the drug.

[0235] The reference cells may also be cells from subjects responding or not responding to several different treatments, and the computer system indicates a preferred treatment for the subject. Accordingly, the invention provides a method for selecting a therapy for a patient having breast cancer, the method comprising: (i) providing the level of expression of one or more genes characteristic of breast cancer in a diseased cell of the patient; (ii) providing a plurality of reference profiles, each associated with a therapy, wherein the subject expression profile and each reference profile has a plurality of values, each value representing the level of expression of a gene characteristic of breast cancer; and (iii) selecting the reference profile most similar to the subject expression profile, to thereby select a therapy for said patient. In a preferred embodiment step (iii) is performed by a computer. The most similar reference profile may be selected by weighing a comparison value of the plurality using a weight value associated with the corresponding expression data.

[0236] The relative abundance of an mRNA in two biological samples can be scored as a perturbation and its magnitude determined (i.e., the abundance is different in the two sources of mRNA tested), or as not perturbed (i.e., the relative abundance is the same). In various embodiments, a difference between the two sources of RNA of at least a factor of about 25% (RNA from one source is 25% more abundant in one source than the other source), more usually about 50%, even more often by a factor of about 2 (twice as abundant), 3 (three times as abundant) or 5 (five times as abundant) is scored as a perturbation. Perturbations can be used by a computer for calculating and expression comparisons.

[0237] Preferably, in addition to identifying a perturbation as positive or negative, it is advantageous to determine the magnitude of the perturbation. This can be carried out, as noted above, by calculating the ratio of the emission of the two fluorophores used for differential labeling, or by analogous methods that will be readily apparent to those of skill in the art.

[0238] The computer readable medium may further comprise a pointer to a descriptor of a stage of breast cancer or to a treatment for breast cancer.

[0239] In operation, the means for receiving gene expression data, the means for comparing the gene expression data, the means for presenting, the means for normalizing, and the means for clustering within the context of the systems of the present invention can involve a programmed computer with the respective functionalities described herein, implemented in hardware or hardware and software; a logic circuit or other component of a programmed computer that performs the operations specifically identified herein, dictated by a computer program; or a computer memory encoded with executable instructions representing a computer program that can cause a computer to function in the particular fashion described herein.

[0240] Those skilled in the art will understand that the systems and methods of the present invention may be applied to a variety of systems, including IBM-compatible personal computers running MS-DOS or Microsoft Windows.

[0241] The computer may have internal components linked to external components. The internal components may include a processor element interconnected with a main memory. The computer system can be an Intel Pentium®-based processor of 200 MHz or greater clock rate and with 32 MB or more of main memory. The external component may comprise a mass storage, which can be one or more hard disks (which are typically packaged together with the processor and memory). Such hard disks are typically of 1 GB or greater storage capacity. Other external components include a user interface device, which can be a monitor, together with an inputting device, which can be a "mouse", or other graphic input devices, and/or a keyboard. A printing device can also be attached to the computer.

[0242] Typically, the computer system is also linked to a network link, which can be part of an Ethernet link to other local computer systems, remote computer systems, or wide area communication networks, such as the Internet. This network link allows the computer system to share data and processing tasks with other computer systems.

[0243] Loaded into memory during operation of this system are several software components, which are both standard in the art and special to the instant invention. These software components collectively cause the computer system to function according to the methods of this invention. These software components are typically stored on a mass storage. A software component represents the operating system, which is responsible for managing the computer system and its network interconnections. This operating system can be, for example, of the Microsoft Windows® family, such as Windows 95, Windows 98, or Windows NT. A software component represents common languages and functions conveniently present on this system to assist programs implementing the methods specific to this invention. Many high or low level computer languages can be used to program the analytic methods of this invention. Instructions can be interpreted during run-time or compiled. Preferred languages include C/C++, and JAVA®. Most preferably, the methods of this invention are programmed in mathematical software packages which allow symbolic entry of equations and high-level specification of processing, including algorithms to be used, thereby freeing a user of the need to procedurally program individual equations or algorithms. Such packages include Matlab from Mathworks (Natick, Mass.), Mathematica from Wolfram Research (Champaign, Ill.), or S-Plus from Math Soft (Cambridge, Mass.). Accordingly, a software component represents the analytic methods of this invention as programmed in a procedural language or symbolic package. In a preferred embodiment, the computer system also contains a database comprising values representing levels of expression of one or more genes characteristic of breast cancer. The database may contain one or more expression profiles of genes characteristic of breast cancer in different cells.

[0244] In an exemplary implementation, to practice the methods of the present invention, a user first loads expression profile data into the computer system. These data can be directly entered by the user from a monitor and keyboard, or from other computer systems linked by a network connection, or on removable storage media such as a CD-ROM or floppy disk or through the network. Next the user causes execution of expression profile analysis software which performs the steps of comparing and, e.g., clustering co-varying genes into groups of genes.

[0245] In another exemplary implementation, expression profiles are compared using a method described in U.S. Patent No. 6,203,987. A user first loads expression profile data into the computer system. Geneset profile definitions are loaded into the memory from the storage media or from a remote computer, preferably from a dynamic geneset database system, through the network. Next the user causes execution of projection software which performs the steps of converting expression profile to projected expression profiles. The projected expression profiles are then displayed.

[0246] In yet another exemplary implementation, a user first leads a projected profile into the memory. The user then causes the loading of a reference profile into the memory. Next, the user causes the execution of comparison software which performs the steps of objectively comparing the profiles.

Detection of variant polynucleotide sequence

[0247] In yet another embodiment, the invention provides methods for determining whether a subject is at risk for developing a disease, such as a predisposition to develop malignant neoplasia, for example breast cancer, associated with an aberrant activity of any one of the polypeptides encoded by any of the polynucleotides of the SEQ ID NO: 1 to 26 or 53 to 75, wherein the aberrant activity of the polypeptide is characterized by detecting the presence or absence of a genetic lesion characterized by at least one of these:

- (i) an alteration affecting the integrity of a gene encoding a marker polypeptides, or
- (ii) the misexpression of the encoding polynucleotide.

[0248] To illustrate, such genetic lesions can be detected by ascertaining the existence of at least one of these:

- I. a deletion of one or more nucleotides from the polynucleotide sequence
- II. an addition of one or more nucleotides to the polynucleotide sequence

III. a substitution of one or more nucleotides of the polynucleotide sequence

IV. a gross chromosomal rearrangement of the polynucleotide sequence

V. a gross alteration in the level of a messenger RNA transcript of the polynucleotide sequence

VI. aberrant modification of the polynucleotide sequence, such as of the methylation pattern of the genomic DNA

VII. the presence of a non-wild type splicing pattern of a messenger RNA transcript of the gene

VIII. a non-wild type level of the marker polypeptide

IX. allelic loss of the gene

X. allelic gain of the gene

XI. inappropriate post-translational modification of the marker polypeptide

[0249] The present invention provides assay techniques for detecting mutations in the encoding polynucleotide sequence. These methods include, but are not limited to, methods involving sequence analysis, Southern blot hybridization, restriction enzyme site mapping, and methods involving detection of absence of nucleotide pairing between the polynucleotide to be analyzed and a probe.

[0250] Specific diseases or disorders, e.g., genetic diseases or disorders, are associated with specific allelic variants of polymorphic regions of certain genes, which do not necessarily encode a mutated protein. Thus, the presence of a specific allelic variant of a polymorphic region of a gene in a subject can render the subject susceptible to developing a specific disease or disorder. Polymorphic regions in genes, can be identified, by determining the nucleotide sequence of genes in populations of individuals. If a polymorphic region is identified, then the link with a specific disease can be determined by studying specific populations of individuals, e.g. individuals which developed a specific disease, such as breast cancer. A polymorphic region can be located in any region of a gene, e.g., exons, in coding or non coding regions of exons, introns, and promoter region.

[0251] In an exemplary embodiment, there is provided a polynucleotide composition comprising a polynucleotide probe including a region of nucleotide sequence which is capable of hybridizing to a sense or antisense sequence of a gene or naturally occurring mutants thereof, or 5' or 3' flanking sequences or intronic sequences naturally associated with the subject genes or naturally occurring mutants thereof. The polynucleotide of a cell is rendered accessible for hybridization, the probe is contacted with the polynucleotide of the sample, and the hybridization of the probe to the sample polynucleotide is detected. Such techniques can be used to detect lesions or allelic variants at either the genomic or mRNA level, including deletions, substitutions, etc., as well as to determine mRNA transcript levels.

[0252] A preferred detection method is allele specific hybridization using probes overlapping the mutation or polymorphic site and having about 5, 10, 20, 25, or 30 nucleotides around the mutation or polymorphic region. In a preferred embodiment of the invention, several probes capable of hybridizing specifically to allelic variants are attached to a solid phase support, e.g., a "chip". Mutation detection analysis using these chips comprising oligonucleotides, also termed "DNA probe arrays" is described e.g., in Cronin et al. (119). In one embodiment, a chip comprises all the allelic variants of at least one polymorphic region of a gene. The solid phase support is then contacted with a test polynucleotide and hybridization to the specific probes is detected. Accordingly, the identity of numerous allelic variants of one or more genes can be identified in a simple hybridization experiment.

[0253] In certain embodiments, detection of the lesion comprises utilizing the probe/primer in a polymerase chain reaction (PCR) (see, e.g. U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligase chain reaction (LCR) [Landegran et al., 1988, (120) and Nakazawa et al., 1994 (121)], the latter of which can be particularly useful for detecting point mutations in the gene; Abravaya et al., 1995, (122)]. In a merely illustrative embodiment, the method includes the steps of (i) collecting a sample of cells from a patient, (ii) isolating polynucleotide (e.g., genomic, mRNA or both) from the cells of the sample, (iii) contacting the polynucleotide sample with one or more primers which specifically hybridize to a polynucleotide sequence under conditions such that hybridization and amplification of the polynucleotide (if present) occurs, and (iv) detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. It is anticipated that PCR and/or LCR may be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described herein.

[0254] Alternative amplification methods include: self sustained sequence replication [Guatelli, J.C. et al., 1990, (123)], transcriptional amplification system [Kwoh, D.Y. et al., 1989, (124)], Q-Beta replicase [Lizardi, F.M. et al., 1988,

(125)], or any other polynucleotide amplification method, followed by the detection of the amplified molecules using techniques well known to those of skill in the art. These detection schemes are especially useful for the detection of polynucleotide molecules if such molecules are present in very low numbers.

[0255] In a preferred embodiment of the subject assay, mutations in, or allelic variants, of a gene from a sample cell are identified by alterations in restriction enzyme cleavage patterns. For example, sample and control DNA is isolated, amplified (optionally), digested with one or more restriction endonucleases, and fragment length sizes are determined by gel electrophoresis. Moreover, the use of sequence specific ribozymes (see, for example, U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

In situ hybridization

[0256] In one aspect, the method comprises *in situ* hybridization with a probe derived from a given marker polynucleotide, which sequence is selected from any of the polynucleotide sequences of the SEQ ID NO: 1 to 9, or 11 to 19 or 21 to 26 and 53 to 75 or a sequence complementary thereto. The method comprises contacting the labeled hybridization probe with a sample of a given type of tissue from a patient potentially having malignant neoplasia and breast cancer in particular as well as normal tissue from a person with no malignant neoplasia, and determining whether the probe labels tissue of the patient to a degree significantly different (e.g., by at least a factor of two, or at least a factor of five, or at least a factor of twenty, or at least a factor of fifty) than the degree to which normal tissue is labelled.

Polypeptide detection

[0257] The subject invention further provides a method of determining whether a cell sample obtained from a subject possesses an abnormal amount of marker polypeptide which comprises (a) obtaining a cell sample from the subject, (b) quantitatively determining the amount of the marker polypeptide in the sample so obtained, and (c) comparing the amount of the marker polypeptide so determined with a known standard, so as to thereby determine whether the cell sample obtained from the subject possesses an abnormal amount of the marker polypeptide. Such marker polypeptides may be detected by immunohistochemical assays, dot-blot assays, ELISA and the like.

Antibodies

[0258] Any type of antibody known in the art can be generated to bind specifically to an epitope of a "BREAST CANCER GENE" polypeptide. An antibody as used herein includes intact immunoglobulin molecules, as well as fragments thereof, such as Fab, F(ab)₂, and Fv, which are capable of binding an epitope of a "BREAST CANCER GENE" polypeptide. Typically, at least 6, 8, 10, or 12 contiguous amino acids are required to form an epitope. However, epitopes which involve non-contiguous amino acids may require more, e.g., at least 15, 25, or 50 amino acids.

[0259] An antibody which specifically binds to an epitope of a "BREAST CANCER GENE" polypeptide can be used therapeutically, as well as in immunochemical assays, such as Western blots, ELISAs, radioimmunoassays, immuno-histochemical assays, immunoprecipitations, or other immunochemical assays known in the art. Various immunoassays can be used to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays are well known in the art. Such immunoassays typically involve the measurement of complex formation between an immunogen and an antibody which specifically binds to the immunogen.

[0260] Typically, an antibody which specifically binds to a "BREAST CANCER GENE" polypeptide provides a detection signal at least 5-, 10-, or 20-fold higher than a detection signal provided with other proteins when used in an immunochemical assay. Preferably, antibodies which specifically bind to "BREAST CANCER GENE" polypeptides do not detect other proteins in immunochemical assays and can immunoprecipitate a "BREAST CANCER GENE" polypeptide from solution.

[0261] "BREAST CANCER GENE" polypeptides can be used to immunize a mammal, such as a mouse, rat, rabbit, guinea pig, monkey, or human, to produce polyclonal antibodies. If desired, a "BREAST CANCER GENE" polypeptide can be conjugated to a carrier protein, such as bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin. Depending on the host species, various adjuvants can be used to increase the immunological response. Such adjuvants include, but are not limited to, Freund's adjuvant, mineral gels (e.g., aluminum hydroxide), and surface active substances (e.g. lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinutrophenol). Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially useful.

[0262] Monoclonal antibodies which specifically bind to a "BREAST CANCER GENE" polypeptide can be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These techniques include, but are not limited to, the hybridoma technique, the human B cell hybridoma technique, and the EBV hybridoma technique [Kohler et al., 1985, (136); Kozbor et al., 1985, (137); Cote et al., 1983, (138) and Cole et

al., 1984, (139)].

[0263] In addition, techniques developed for the production of chimeric antibodies, the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used [Morrison et al., 1984, (140); Neuberger et al., 1984, (141); Takeda et al., 1985, (142)]. Monoclonal and other antibodies also can be humanized to prevent a patient from mounting an immune response against the antibody when it is used therapeutically. Such antibodies may be sufficiently similar in sequence to human antibodies to be used directly in therapy or may require alteration of a few key residues. Sequence differences between rodent antibodies and human sequences can be minimized by replacing residues which differ from those in the human sequences by site directed mutagenesis of individual residues or by grating of entire complementarity determining regions. Alternatively, humanized antibodies can be produced using recombinant methods, as described in GB218638B. Antibodies which specifically bind to a "BREAST CANCER GENE" polypeptide can contain antigen binding sites which are either partially or fully humanized, as disclosed in U.S. Patent 5,565,332.

[0264] Alternatively, techniques described for the production of single chain antibodies can be adapted using methods known in the art to produce single chain antibodies which specifically bind to "BREAST CANCER GENE" polypeptides. Antibodies with related specificity, but of distinct idiotype composition, can be generated by chain shuffling from random combinatorial immunoglobulin libraries [Burton, 1991, (143)].

[0265] Single-chain antibodies also can be constructed using a DNA amplification method, such as PCR, using hybridoma cDNA as a template [Thirion et al., 1996, (144)]. Single-chain antibodies can be mono- or bispecific, and can be bivalent or tetravalent. Construction of tetravalent, bispecific single-chain antibodies is taught, for example, in Coloma & Morrison, (145). Construction of bivalent, bispecific single-chain antibodies is taught in Mallender & Voss, (146).

[0266] A nucleotide sequence encoding a single-chain antibody can be constructed using manual or automated nucleotide synthesis, cloned into an expression construct using standard recombinant DNA methods, and introduced into a cell to express the coding sequence, as described below. Alternatively, single-chain antibodies can be produced directly using, for example, filamentous phage technology [Verhaar et al., 1995, (147); Nicholls et al., 1993, (148)].

[0267] Antibodies which specifically bind to "BREAST CANCER GENE" polypeptides also can be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature [Orlandi et al., 1989, (149) and Winter et al., 1991, (150)].

[0268] Other types of antibodies can be constructed and used therapeutically in methods of the invention. For example, chimeric antibodies can be constructed as disclosed in WO 93/03151. Binding proteins which are derived from immunoglobulins and which are multivalent and multispecific, such as the antibodies described in WO 94/13804, also can be prepared.

[0269] Antibodies according to the invention can be purified by methods well known in the art. For example, antibodies can be affinity purified by passage over a column to which a "BREAST CANCER GENE" polypeptide is bound. The bound antibodies can then be eluted from the column using a buffer with a high salt concentration.

[0270] Immunoassays are commonly used to quantify the levels of proteins in cell samples, and many other immunoassay techniques are known in the art. The invention is not limited to a particular assay procedure, and therefore is intended to include both homogeneous and heterogeneous procedures. Exemplary immunoassays which can be conducted according to the invention include fluorescence polarisation immunoassay (FPIA), fluorescence immunoassay (FIA), enzyme immunoassay (EIA), nephelometric inhibition immunoassay (NIA), enzyme linked immunosorbent assay (ELISA), and radioimmunoassay (RIA). An indicator moiety, or label group, can be attached to the subject antibodies and is selected so as to meet the needs of various uses of the method which are often dictated by the availability of assay equipment and compatible immunoassay procedures. General techniques to be used in performing the various immunoassays noted above are known to those of ordinary skill in the art.

[0271] In another embodiment, the level of at least one product encoded by any of the polynucleotide sequences of the SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19 or 21 to 26 or 53 to 75 or of at least 2 products encoded by a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or a sequence complementary thereto, in a biological fluid (e.g., blood or urine) of a patient may be determined as a way of monitoring the level of expression of the marker polynucleotide sequence in cells of that patient. Such a method would include the steps of obtaining a sample of a biological fluid from the patient, contacting the sample (or proteins from the sample) with an antibody specific for an encoded marker polypeptide, and determining the amount of immune complex formation by the antibody, with the amount of immune complex formation being indicative of the level of the marker encoded product in the sample. This determination is particularly instructive when compared to the amount of immune complex formation by the same antibody in a control sample taken from a normal individual or in one or more samples previously or subsequently obtained from the same person.

[0272] In another embodiment, the method can be used to determine the amount of marker polypeptide present in a cell, which in turn can be correlated with progression of the disorder, e.g., plaque formation. The level of the marker polypeptide can be used predictively to evaluate whether a sample of cells contains cells which are, or are predisposed

towards becoming, plaque associated cells. The observation of marker polypeptide level can be utilized in decisions regarding, e.g., the use of more stringent therapies.

[0273] As set out above, one aspect of the present invention relates to diagnostic assays for determining, in the context of cells isolated from a patient, if the level of a marker polypeptide is significantly reduced in the sample cells. The term "significantly reduced" refers to a cell phenotype wherein the cell possesses a reduced cellular amount of the marker polypeptide relative to a normal cell of similar tissue origin. For example, a cell may have less than about 50%, 25%, 10%, or 5% of the marker polypeptide that a normal control cell. In particular, the assay evaluates the level of marker polypeptide in the test cells, and, preferably, compares the measured level with marker polypeptide detected in at least one control cell, e.g., a normal cell and/or a transformed cell of known phenotype.

[0274] Of particular importance to the subject invention is the ability to quantify the level of marker polypeptide as determined by the number of cells associated with a normal or abnormal marker polypeptide level. The number of cells with a particular marker polypeptide phenotype may then be correlated with patient prognosis. In one embodiment of the invention, the marker polypeptide phenotype of the lesion is determined as a percentage of cells in a biopsy which are found to have abnormally high/low levels of the marker polypeptide. Such expression may be detected by immunohistochemical assays, dot-blot assays, ELISA and the like.

Immunohistochemistry

[0275] Where tissue samples are employed, immunohistochemical staining may be used to determine the number of cells having the marker polypeptide phenotype. For such staining, a multiblock of tissue is taken from the biopsy or other tissue sample and subjected to proteolytic hydrolysis, employing such agents as protease K or pepsin. In certain embodiments, it may be desirable to isolate a nuclear fraction from the sample cells and detect the level of the marker polypeptide in the nuclear fraction.

[0276] The tissues samples are fixed by treatment with a reagent such as formalin, glutaraldehyde, methanol, or the like. The samples are then incubated with an antibody, preferably a monoclonal antibody, with binding specificity for the marker polypeptides. This antibody may be conjugated to a Label for subsequent detection of binding. Samples are incubated for a time sufficient for formation of the immuno-complexes. Binding of the antibody is then detected by virtue of a Label conjugated to this antibody. Where the antibody is unlabelled, a second labeled antibody may be employed, e.g., which is specific for the isotype of the anti-marker polypeptide antibody. Examples of labels which may be employed include radionuclides, fluorescence, chemiluminescence, and enzymes.

[0277] Where enzymes are employed, the Substrate for the enzyme may be added to the samples to provide a colored or fluorescent product. Examples of suitable enzymes for use in conjugates include horseradish peroxidase, alkaline phosphatase, malate dehydrogenase and the like. Where not commercially available, such antibody-enzyme conjugates are readily produced by techniques known to those skilled in the art.

[0278] In one embodiment, the assay is performed as a dot blot assay. The dot blot assay finds particular application where tissue samples are employed as it allows determination of the average amount of the marker polypeptide associated with a Single cell by correlating the amount of marker polypeptide in a cell-free extract produced from a predetermined number of cells.

[0279] In yet another embodiment, the invention contemplates using one or more antibodies which are generated against one or more of the marker polypeptides of this invention, which polypeptides are encoded by any of the polynucleotide sequences of the SEQ ID NO: 1 to 26 or 53 to 75. Such a panel of antibodies may be used as a reliable diagnostic probe for breast cancer. The assay of the present invention comprises contacting a biopsy sample containing cells, e.g., macrophages, with a panel of antibodies to one or more of the encoded products to determine the presence or absence of the marker polypeptides.

[0280] The diagnostic methods of the subject invention may also be employed as follow-up to treatment, e.g., quantification of the level of marker polypeptides may be indicative of the effectiveness of current or previously employed therapies for malignant neoplasia and breast cancer in particular as well as the effect of these therapies upon patient prognosis.

[0281] The diagnostic assays described above can be adapted to be used as prognostic assays, as well. Such an application takes advantage of the sensitivity of the assays of the Invention to events which take place at characteristic stages in the progression of plaque generation in case of malignant neoplasia. For example, a given marker gene may be up- or down-regulated at a very early stage, perhaps before the cell is developing into a foam cell, while another marker gene may be characteristically up or down regulated only at a much later stage. Such a method could involve the steps of contacting the mRNA of a test cell with a polynucleotide probe derived from a given marker polynucleotide which is expressed at different characteristic levels in breast cancer tissue cells at different stages of malignant neoplasia progression, and determining the approximate amount of hybridization of the probe to the mRNA of the cell, such amount being an indication of the level of expression of the gene in the cell, and thus an indication of the stage of disease progression of the cell; alternatively, the assay can be carried out with an antibody specific for the gene

product of the given marker polynucleotide, contacted with the proteins of the test cell. A battery of such tests will disclose not only the existence of a certain arteriosclerotic plaque, but also will allow the clinician to select the mode of treatment most appropriate for the disease, and to predict the likelihood of success of that treatment.

[0282] The methods of the invention can also be used to follow the clinical course of a given breast cancer predisposition. For example, the assay of the invention can be applied to a blood sample from a patient; following treatment of the patient for BREAST CANCER, another blood sample is taken and the test repeated. Successful treatment will result in removal of demonstrate differential expression, characteristic of the breast cancer tissue cells, perhaps approaching or even surpassing normal levels.

10 Polypeptide activity

[0283] In one embodiment the present invention provides a method for screening potentially therapeutic agents which modulate the activity of one or more "BREAST CANCER GENE" polypeptides, such that if the activity of the polypeptide is increased as a result of the upregulation of the "BREAST CANCER GENE" in a subject having or at risk for malignant neoplasia and breast cancer in particular, the therapeutic substance will decrease the activity of the polypeptide relative to the activity of the same polypeptide in a subject not having or not at risk for malignant neoplasia or breast cancer in particular but not treated with the therapeutic agent. Likewise, if the activity of the polypeptide as a result of the downregulation of the "BREAST CANCER GENE" is decreased in a subject having or at risk for malignant neoplasia or breast cancer in particular, the therapeutic agent will increase the activity of the polypeptide relative to the activity of the same polypeptide in a subject not having or not at risk for malignant neoplasia or breast cancer in particular, but not treated with the therapeutic agent.

[0284] The activity of the "BREAST CANCER GENE" polypeptides indicated in Table 2 or 3 may be measured by any means known to those of skill in the art, and which are particular for the type of activity performed by the particular polypeptide. Examples of specific assays which may be used to measure the activity of particular polynucleotides are shown below.

a) G protein coupled receptors

[0285] In one embodiment, the "BREAST CANCER GENE" polynucleotide may encode a G protein coupled receptor. In one embodiment, the present invention provides a method of screening potential modulators (inhibitors or activators) of the G protein coupled receptor by measuring changes in the activity of the receptor in the presence of a candidate modulator.

35 1. G_i-coupled receptors

[0286] Cells (such as CHO cells or primary cells) are stably transfected with the relevant receptor and with an inducible CRE-luciferase construct. Cells are grown in 50% Dulbecco's modified Eagle medium / 50% F12 (DMEM/F12) supplemented with 10% FBS, at 37°C in a humidified atmosphere with 10% CO₂ and are routinely split at a ratio of 1:10 every 2 or 3 days. Test cultures are seeded into 384 - well plates at an appropriate density (e.g. 2000 cells / well in 35 µl cell culture medium) in DMEM/F12 with FBS, and are grown for 48 hours (range: ~ 24 - 60 hours, depending on cell line). Growth medium is then exchanged against serum free medium (SFM; e.g. Ultra-CHO), containing 0,1% BSA. Test compounds dissolved in DMSO are diluted in SFM and transferred to the test cultures (maximal final concentration 10 µmolar), followed by addition of forskolin (~ 1 µmolar, final conc.) in SFM + 0,1% BSA 10 minutes later. In case of antagonist screening both, an appropriate concentration of agonist, and forskolin are added. The plates are incubated at 37°C in 10% CO₂ for 3 hours. Then the supernatant is removed, cells are lysed with lysis reagent (25 mmolar phosphate-buffer, pH 7,8, containing 2 mmolar DDT, 10% glycerol and 3% Triton X100). The luciferase reaction is started by addition of substrate-buffer (e.g. luciferase assay reagent, Promega) and luminescence is immediately determined (e.g. Berthold luminometer or Hamamatsu camera system).

50 2. G_s-coupled receptors

[0287] Cells (such as CHO cells or primary cells) are stably transfected with the relevant receptor and with an inducible CRE-luciferase construct. Cells are grown in 50% Dulbecco's modified Eagle medium / 50% F12 (DMEM/F12) supplemented with 10% FBS, at 37°C in a humidified atmosphere with 10% CO₂ and are routinely split at a ratio of 1:10 every 2 or 3 days. Test cultures are seeded into 384 - well plates at an appropriate density (e.g. 1000 or 2000 cells / well in 35 µl cell culture medium) in DMEM/F12 with FBS, and are grown for 48 hours (range: ~ 24 - 60 hours, depending on cell line). The assay is started by addition of test-compounds in serum free medium (SFM; e.g. Ultra-CHO) containing 0,1% BSA: Test compounds are dissolved in DMSO, diluted in SFM and transferred to the test cultures (maximal final

concentration 10 μ molar, DMSO conc. < 0,6 %). In case of antagonist screening an appropriate concentration of agonist is added 5 - 10 minutes later. The plates are incubated at 37°C in 10% CO₂ for 3 hours. Then the cells are lysed with 10 μ l lysis reagent per well (25 mmolar phosphate-buffer, pH 7,8 , containing 2 mmolar DDT, 10% glycerol and 3% Triton X100) and the luciferase reaction is started by addition of 20 μ l substrate-buffer per well (e.g. luciferase assay reagent, Promega). Measurement of luminescence is started immediately (e.g. Berthold luminometer or Hamamatsu camera system).

3. G_i-coupled receptors

[0288] Cells (such as CHO cells or primary cells) are stably transfected with the relevant receptor. Cells expressing functional receptor protein are grown in 50% Dulbecco's modified Eagle medium / 50% F12 (DMEM/F12) supplemented with 10% FBS, at 37°C in a humidified atmosphere with 5% CO₂ and are routinely split at a cell line dependent ratio every 3 or 4 days. Test cultures are seeded into 384 - well plates at an appropriate density (e.g. 2000 cells / well in 35 μ l cell culture medium) in DMEM/F12 with FBS, and are grown for 48 hours (range: ~ 24 - 60 hours, depending on cell line). Growth medium is then exchanged against physiological salt solution (e.g. Tyrode solution). Test compounds dissolved in DMSO are diluted in Tyrode solution containing 0.1% BSA and transferred to the test cultures (maximal final concentration 10 molar). After addition of the receptor specific agonist the resulting Gq-mediated intracellular calcium increase is measured using appropriate read-out systems (e.g. calcium-sensitive dyes).

b) Ion channels

[0289] Ion channels are integral membrane proteins involved in electrical signaling, transmembrane signal transduction, and electrolyte and solute transport. By forming macromolecular pores through the membrane lipid bilayer, ion channels account for the flow of specific ion species driven by the electrochemical potential gradient for the permeating ion. At the single molecule level, individual channels undergo conformational transitions ("gating") between the 'open' (ion conducting) and 'closed' (non conducting) state. Typical single channel openings last for a few milliseconds and result in elementary transmembrane currents in the range of 10⁻⁹ - 10⁻¹² Ampere. Channel gating is controlled by various chemical and/or biophysical parameters, such as neurotransmitters and intracellular second messengers ('ligand-gated' channels) or membrane potential ('voltage-gated' channels). Ion channels are functionally characterized by their ion selectivity, gating properties, and regulation by hormones and pharmacological agents. Because of their central role in signaling and transport processes, ion channels present ideal targets for pharmacological therapeutics in various pathophysiological settings.

[0290] In one embodiment, the "BREAST CANCER GENE" may encode an ion channel. In one embodiment, the present invention provides a method of screening potential activators or inhibitors of channels activity of the "BREAST CANCER GENE" polypeptide. Screening for compounds interaction with ion channels to either inhibit or promote their activity can be based on (1.) binding and (2.) functional assays in living cells [Hille (183)].

1. For ligand-gated channels, e.g. ionotropic neurotransmitter/hormone receptors, assays can be designed detecting binding to the target by competition between the compound and a labeled ligand.

2. Ion channel function can be tested functionally in living cells. Target proteins are either expressed endogenously in appropriate reporter cells or are introduced recombinantly. Channel activity can be monitored by (2.1) concentration changes of the permeating ion (most prominently Ca²⁺ ions), (2.2) by changes in the transmembrane electrical potential gradient, and (2.3) by measuring a cellular response (e.g. expression of a reporter gene, secretion of a neurotransmitter) triggered or modulated by the target activity.

2.1 Channel activity results in transmembrane ion fluxes. Thus activation of ionic channels can be monitored by the resulting changes in intracellular ion concentrations using luminescent or fluorescent indicators. Because of its wide dynamic range and availability of suitable indicators this applies particularly to changes in intracellular Ca²⁺ ion concentration ([Ca²⁺]_i). [Ca²⁺]_i can be measured, for example, by aequorin luminescence or fluorescence dye technology (e.g. using Fluo-3, Indo-1, Fura-2). Cellular assays can be designed where either the Ca²⁺ flux through the target channel itself is measured directly or where modulation of the target channel affects membrane potential and thereby the activity of co-expressed voltage-gated Ca²⁺ channels.

2.2 Ion channel currents result in changes of electrical membrane potential (V_m) which can be monitored directly using potentiometric fluorescent probes. These electrically charged indicators (e.g. the anionic oxonol dye DiBAC₄(3)) redistribute between extra- and intracellular compartment in response to voltage changes. The equilibrium distribution is governed by the Nernst-equation. Thus changes in membrane potential results

in concomitant changes in cellular fluorescence. Again, changes in V_m might be caused directly by the activity of the target ion channel or through amplification and/or prolongation of the signal by channels co-expressed in the same cell.

2.3 Target channel activity can cause cellular Ca^{2+} entry either directly or through activation of additional Ca^{2+} channel (see 2.1). The resulting intracellular Ca^{2+} signals regulate a variety of cellular responses, e.g. secretion or gene transcription. Therefore modulation of the target channel can be detected by monitoring secretion of a known hormone/transmitter from the target-expressing cell or through expression of a reporter gene (e.g. luciferase) controlled by an Ca^{2+} -responsive promoter element (e.g. cyclic AMP/ Ca^{2+} -responsive elements; CRE).

c) DNA-binding proteins and transcription factors

[0291] In one embodiment, the "BREAST CANCER GENE" may encode a DNA-binding protein or a transcription factor. The activity of such a DNA-binding protein or a transcription factor may be measured, for example, by a promoter assay which measures the ability of the DNA-binding protein or the transcription factor to initiate transcription of a test sequence linked to a particular promoter. In one embodiment, the present invention provides a method of screening test compounds for its ability to modulate the activity of such a DNA-binding protein or a transcription factor by measuring the changes in the expression of a test gene which is regulated by a promoter which is responsive to the transcription factor.

d) Promotor assays

[0292] A promoter assay was set up with a human hepatocellular carcinoma cell HepG2 that was stably transfected with a luciferase gene under the control of a gene of interest (e.g. thyroid hormone) regulated promoter. The vector 2xIRoLuc, which was used for transfection, carries a thyroid hormone responsive element (TRE) of two 12 bp inverted palindromes separated by an 8 bp spacer in front of a tk minimal promoter and the luciferase gene. Test cultures were seeded in 96 well plates in serum - free Eagle's Minimal Essential Medium supplemented with glutamine, tricine, sodium pyruvate, non - essential amino acids, insulin, selen, transferrin, and were cultivated in a humidified atmosphere at 10 % CO_2 at 37°C. After 48 hours of incubation serial dilutions of test compounds or reference compounds (L-T3, L-T4 e.g.) and co-stimulator if appropriate (final concentration 1 nM) were added to the cell cultures and incubation was continued for the optimal time (e.g. another 4-72 hours). The cells were then lysed by addition of buffer containing Triton X100 and luciferin and the luminescence of luciferase induced by T3 or other compounds was measured in a luminometer. For each concentration of a test compound replicates of 4 were tested. EC_{50} values for each test compound were calculated by use of the Graph Pad Prism Scientific software.

Screening Methods

[0293] The invention provides assays for screening test compounds which bind to or modulate the activity of a "BREAST CANCER GENE" polypeptide or a "BREAST CANCER GENE" polynucleotide. A test compound preferably binds to a "BREAST CANCER GENE" polypeptide or polynucleotide. More preferably, a test compound decreases or increases "BREAST CANCER GENE" activity by at least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the test compound.

Test Compounds

[0294] Test compounds can be pharmacological agents already known in the art or can be compounds previously unknown to have any pharmacological activity. The compounds can be naturally occurring or designed in the laboratory. They can be isolated from microorganisms, animals, or plants, and can be produced recombinant, or synthesised by chemical methods known in the art. If desired, test compounds can be obtained using any of the numerous combinatorial library methods known in the art, including but not limited to, biological libraries, spatially addressable parallel solid phase or solution phase libraries, synthetic library methods requiring de-convolution, the one-bead one-compound library method, and synthetic library methods using affinity chromatography selection. The biological library approach is limited to polypeptide libraries, while the other four approaches are applicable to polypeptide, non-peptide oligomer, or small molecule libraries of compounds. [For review see Lam, 1997, (151)].

[0295] Methods for the synthesis of molecular libraries are well known in the art [see, for example, DeWitt et al., 1993, (152); Erb et al., 1994, (153); Zuckermann et al., 1994, (154); Cho et al., 1993, (155); Carell et al., 1994, (156) and Gallop et al., 1994, (157)]. Libraries of compounds can be presented in solution [see, e.g., Houghten, 1992, (158)],

or on beads [Lam, 1991, (159)], DNA-chips [Fodor, 1993, (160)], bacteria or spores [Ladner, U.S. Patent 5,223,409], plasmids [Cull et al., 1992, (161)], or phage [Scott & Smith, 1990, (162); Devlin, 1990, (163); Cwirla et al., 1990, (164); Felici, 1991, (165)].

High Throughput Screening

[0296] Test compounds can be screened for the ability to bind to "BREAST CANCER GENE" polypeptides or polynucleotides or to affect "BREAST CANCER GENE" activity or "BREAST CANCER GENE" expression using high throughput screening. Using high throughput screening, many discrete compounds can be tested in parallel so that large numbers of test compounds can be quickly screened. The most widely established techniques utilize 96-well, 384-well or 1536-well microtiter plates. The wells of the microtiter plates typically require assay volumes that range from 5 to 500 μ l. In addition to the plates, many instruments, materials, pipettors, robotics, plate washers, and plate readers are commercially available to fit the microwell formats.

[0297] Alternatively, free format assays, or assays that have no physical barrier between samples, can be used. For example, an assay using pigment cells (melanocytes) in a simple homogeneous assay for combinatorial peptide libraries is described by Jayawickreme et al., (166). The cells are placed under agarose in culture dishes, then beads that carry combinatorial compounds are placed on the surface of the agarose. The combinatorial compounds are partially released the compounds from the beads. Active compounds can be visualised as dark pigment areas because, as the compounds diffuse locally into the gel matrix, the active compounds cause the cells to change colors.

[0298] Another example of a free format assay is described by Chelsky, (167). Chelsky placed a simple homogenous enzyme assay for carbonic anhydrase inside an agarose gel such that the enzyme in the gel would cause a color change throughout the gel.

[0299] Thereafter, beads carrying combinatorial compounds via a photolinker were placed inside the gel and the compounds were partially released by UV light. Compounds that inhibited the enzyme were observed as local zones of inhibition having less color change.

[0300] In another example, combinatorial libraries were screened for compounds that had cytotoxic effects on cancer cells growing in agar [Salmon et al., 1996, (168)].

[0301] Another high throughput screening method is described in Beutel et al., U.S. Patent 5,976,813. In this method, test samples are placed in a porous matrix. One or more assay components are then placed within, on top of, or at the bottom of a matrix such as a gel, a plastic sheet, a filter, or other form of easily manipulated solid support. When samples are introduced to the porous matrix they diffuse sufficiently slowly, such that the assays can be performed without the test samples running together.

Binding Assays

[0302] For binding assays, the test compound is preferably a small molecule which binds to and occupies, for example, the ATP/GTP binding site of the enzyme or the active site of a "BREAST CANCER GENE" polypeptide, such that normal biological activity is prevented. Examples of such small molecules include, but are not limited to, small peptides or peptide-like molecules.

[0303] In binding assays, either the test compound or a "BREAST CANCER GENE" polypeptide can comprise a detectable label, such as a fluorescent, radioisotopic, chemiluminescent, or enzymatic label, such as horseradish peroxidase, alkaline phosphatase, or luciferase. Detection of a test compound which is bound to a "BREAST CANCER GENE" polypeptide can then be accomplished, for example, by direct counting of radioemission, by scintillation counting, or by determining conversion of an appropriate substrate to a detectable product.

[0304] Alternatively, binding of a test compound to a "BREAST CANCER GENE" polypeptide can be determined without labeling either of the interactants. For example, a microphysiometer can be used to detect binding of a test compound with a "BREAST CANCER GENE" polypeptide. A microphysiometer (e.g., Cytosensor[®]) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between a test compound and a "BREAST CANCER GENE" polypeptide [McConnell et al., 1992, (169)].

[0305] Determining the ability of a test compound to bind to a "BREAST CANCER GENE" polypeptide also can be accomplished using a technology such as real-time Bimolecular Interaction Analysis (BIA) [Sjölinder & Urbaniczky, 1991, (170), and Szabo et al., 1995, (171)]. BIA is a technology for studying biospecific interactions in real time, without labeling any of the interactants (e.g., BIAcore[™]). Changes in the optical phenomenon surface plasmon resonance (SPR) can be used as an indication of real-time reactions between biological molecules.

[0306] In yet another aspect of the invention, a "BREAST CANCER GENE" polypeptide can be used as a "bait protein" in a two-hybrid assay or three-hybrid assay [see, e.g., U.S. Patent 5,283,317; Zervos et al., 1993, (172); Madura et al., 1993, (173); Bartel et al., 1993, (174); Iwabuchi et al., 1993, (175) and Brent WO 94/10300], to identify

other proteins which bind to or interact with the "BREAST CANCER GENE" polypeptide and modulate its activity.

[0307] The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. For example, in one construct, polynucleotide encoding a "BREAST CANCER GENE" polypeptide can be fused to a polynucleotide encoding the DNA binding domain of a known transcription factor (e.g., GAL4). In the other construct a DNA sequence that encodes an unidentified protein ("prey" or "sample") can be fused to a polynucleotide that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact in vivo to form an protein-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ), which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected, and cell colonies containing the functional transcription factor can be isolated and used to obtain the DNA sequence encoding the protein which interacts with the "BREAST CANCER GENE" polypeptide.

[0308] It may be desirable to immobilize either a "BREAST CANCER GENE" polypeptide (or polynucleotide) or the test compound to facilitate separation of bound from unbound forms of one or both of the interactants, as well as to accommodate automation of the assay. Thus, either a "BREAST CANCER GENE" polypeptide (or polynucleotide) or the test compound can be bound to a solid support. Suitable solid supports include, but are not limited to, glass or plastic slides, tissue culture plates, microtiter wells, tubes, silicon chips, or particles such as beads (including, but not limited to, latex, polystyrene, or glass beads). Any method known in the art can be used to attach a "BREAST CANCER GENE" polypeptide (or polynucleotide) or test compound to a solid support, including use of covalent and non-covalent linkages, passive absorption, or pairs of binding moieties attached respectively to the polypeptide (or polynucleotide) or test compound and the solid support. Test compounds are preferably bound to the solid support in an array, so that the location of individual test compounds can be tracked. Binding of a test compound to a "BREAST CANCER GENE" polypeptide (or polynucleotide) can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and microcentrifuge tubes.

[0309] In one embodiment, a "BREAST CANCER GENE" polypeptide is a fusion protein comprising a domain that allows the "BREAST CANCER GENE" polypeptide to be bound to a solid support. For example, glutathione S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, Mo.) or glutathione derivatized microtiter plates, which are then combined with the test compound or the test compound and the nonadsorbed "BREAST CANCER GENE" polypeptide; the mixture is then incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components. Binding of the interactants can be determined either directly or indirectly, as described above. Alternatively, the complexes can be dissociated from the solid support before binding is determined.

[0310] Other techniques for immobilising proteins or polynucleotides on a solid support also can be used in the screening assays of the invention. For example, either a "BREAST CANCER GENE" polypeptide (or polynucleotide) or a test compound can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated "BREAST CANCER GENE" polypeptides (or polynucleotides) or test compounds can be prepared from biotin NHS (N-hydroxysuccinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals, Rockford, Ill.) and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies which specifically bind to a "BREAST CANCER GENE" polypeptide, polynucleotide, or a test compound, but which do not interfere with a desired binding site, such as the ATP/GTP binding site or the active site of the "BREAST CANCER GENE" polypeptide, can be derivatised to the wells of the plate. Unbound target or protein can be trapped in the wells by antibody conjugation.

[0311] Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies which specifically bind to a "BREAST CANCER GENE" polypeptide or test compound, enzyme-linked assays which rely on detecting an activity of a "BREAST CANCER GENE" polypeptide, and SDS gel electrophoresis under non-reducing conditions.

[0312] Screening for test compounds which bind to a "BREAST CANCER GENE" polypeptide or polynucleotide also can be carried out in an intact cell. Any cell which comprises a "BREAST CANCER GENE" polypeptide or polynucleotide can be used in a cell-based assay system. A "BREAST CANCER GENE" polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Binding of the test compound to a "BREAST CANCER GENE" polypeptide or polynucleotide is determined as described above.

Modulation of Gene Expression

[0313] In another embodiment, test compounds which increase or decrease "BREAST CANCER GENE" expression are identified. A "BREAST CANCER GENE" polynucleotide is contacted with a test compound, and the expression of an RNA or polypeptide product of the "BREAST CANCER GENE" polynucleotide is determined. The level of expression of appropriate mRNA or polypeptide in the presence of the test compound is compared to the level of expression of

mRNA or polypeptide in the absence of the test compound. The test compound can then be identified as a modulator of expression based on this comparison. For example, when expression of mRNA or polypeptide is greater in the presence of the test compound than in its absence, the test compound is identified as a stimulator or enhancer of the mRNA or polypeptide expression. Alternatively, when expression of the mRNA or polypeptide is less in the presence of the test compound than in its absence, the test compound is identified as an inhibitor of the mRNA or polypeptide expression.

[0314] The level of "BREAST CANCER GENE" mRNA or polypeptide expression in the cells can be determined by methods well known in the art for detecting mRNA or polypeptide. Either qualitative or quantitative methods can be used. The presence of polypeptide products of a "BREAST CANCER GENE" polynucleotide can be determined, for example, using a variety of techniques known in the art, including immunochemical methods such as radioimmunoassay, Western blotting, and immunohistochemistry. Alternatively, polypeptide synthesis can be determined in vivo, in a cell culture, or in an in vitro translation system by detecting incorporation of labeled amino acids into a "BREAST CANCER GENE" polypeptide.

[0315] Such screening can be carried out either in a cell-free assay system or in an intact cell. Any cell which expresses a "BREAST CANCER GENE" polynucleotide can be used in a cell-based assay system. A "BREAST CANCER GENE" polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Either a primary culture or an established cell line, such as CHO or human embryonic kidney 293 cells, can be used.

Therapeutic Indications and Methods

[0316] Therapies for treatment of breast cancer primarily relied upon effective chemotherapeutic drugs for intervention on the cell proliferation, cell growth or angiogenesis. The advent of genomics-driven molecular target identification has opened up the possibility of identifying new breast cancer-specific targets for therapeutic intervention that will provide safer, more effective treatments for malignant neoplasia patients and breast cancer patients in particular. Thus, newly discovered breast cancer-associated genes and their products can be used as tools to develop innovative therapies. The identification of the Her2/neu receptor kinase presents exciting new opportunities for treatment of a certain subset of tumor patients as described before. Genes playing important roles in any of the physiological processes outlined above can be characterized as breast cancer targets. Genes or gene fragments identified through genomics can readily be expressed in one or more heterologous expression systems to produce functional recombinant proteins. These proteins are characterized in vitro for their biochemical properties and then used as tools in high-throughput molecular screening programs to identify chemical modulators of their biochemical activities. Modulators of target gene expression or protein activity can be identified in this manner and subsequently tested in cellular and in vivo disease models for therapeutic activity. Optimization of lead compounds with iterative testing in biological models and detailed pharmacokinetic and toxicological analyses form the basis for drug development and subsequent testing in humans.

[0317] This invention further pertains to the use of novel agents identified by the screening assays described above. Accordingly, it is within the scope of this invention to use a test compound identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a modulating agent, an antisense polynucleotide molecule, a specific antibody, ribozyme, or a human "BREAST CANCER GENE" polypeptide binding molecule) can be used in an animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above described screening assays for treatments as described herein.

[0318] A reagent which affects human "BREAST CANCER GENE" activity can be administered to a human cell, either in vitro or in vivo, to reduce or increase human "BREAST CANCER GENE" activity. The reagent preferably binds to an expression product of a human "BREAST CANCER GENE". If the expression product is a protein, the reagent is preferably an antibody. For treatment of human cells ex vivo, an antibody can be added to a preparation of stem cells which have been removed from the body. The cells can then be replaced in the same or another human body, with or without clonal propagation, as is known in the art.

[0319] In one embodiment, the reagent is delivered using a liposome. Preferably, the liposome is stable in the animal into which it has been administered for at least about 30 minutes, more preferably for at least about 1 hour, and even more preferably for at least about 24 hours. A liposome comprises a lipid composition that is capable of targeting a reagent, particularly a polynucleotide, to a particular site in an animal, such as a human. Preferably, the lipid composition of the liposome is capable of targeting to a specific organ of an animal, such as the lung, liver, spleen, heart brain, lymph nodes, and skin.

[0320] A liposome useful in the present invention comprises a lipid composition that is capable of fusing with the plasma membrane of the targeted cell to deliver its contents to the cell. Preferably, the transfection efficiency of a liposome is about 0.5 µg of DNA per 16 nmol of liposome delivered to about 10⁶ cells, more preferably about 1.0 µg

of DNA per 16 nmol of liposome delivered to about 10^6 cells, and even more preferably about 2.0 μg of DNA per 16 nmol of liposome delivered to about 10^6 cells. Preferably, a liposome is between about 100 and 500 nm, more preferably between about 150 and 450 nm, and even more preferably between about 200 and 400 nm in diameter.

[0321] Suitable liposomes for use in the present invention include those liposomes usually used in, for example, gene delivery methods known to those of skill in the art. More preferred liposomes include liposomes having a polycationic lipid composition and/or liposomes having a cholesterol backbone conjugated to polyethylene glycol. Optionally, a liposome comprises a compound capable of targeting the liposome to a particular cell type, such as a cell-specific ligand exposed on the outer surface of the liposome.

[0322] Complexing a liposome with a reagent such as an antisense oligonucleotide or ribozyme can be achieved using methods which are standard in the art (see, for example, U.S. Patent 5,705,151). Preferably, from about 0.1 μg to about 10 μg of polynucleotide is combined with about 8 nmol of liposomes, more preferably from about 0.5 μg to about 5 μg of polynucleotides are combined with about 8 nmol liposomes, and even more preferably about 1.0 μg of polynucleotides is combined with about 8 nmol liposomes.

[0323] In another embodiment, antibodies can be delivered to specific tissues in vivo using receptor-mediated targeted delivery. Receptor-mediated DNA delivery techniques are taught in, for example, Findeis et al., 1993, (176); Chiou et al., 1994, (177); Wu & Wu, 1988, (178); Wu et al., 1994, (179); Zenke et al., 1990, (180); Wu et al., 1991, (181).

Determination of a Therapeutically Effective Dose

[0324] The determination of a therapeutically effective dose is well within the capability of those skilled in the art. A therapeutically effective dose refers to that amount of active ingredient which increases or decreases human "BREAST CANCER GENE" activity relative to the human "BREAST CANCER GENE" activity which occurs in the absence of the therapeutically effective dose.

[0325] For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays or in animal models, usually mice, rabbits, dogs, or pigs. The animal model also can be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

[0326] Therapeutic efficacy and toxicity, e.g., ED_{50} (the dose therapeutically effective in 50% of the population) and LD_{50} (the dose lethal to 50% of the population), can be determined by standard pharmaceutical procedures in cell cultures or experimental animals. The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the ratio, $\text{LD}_{50}/\text{ED}_{50}$.

[0327] Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that include the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

[0328] The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active ingredient or to maintain the desired effect. Factors which can be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions can be administered every 3 to 4 days, every week, or once every two weeks depending on the half-life and clearance rate of the particular formulation.

[0329] Normal dosage amounts can vary from 0.1 to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

[0330] If the reagent is a single-chain antibody, polynucleotides encoding the antibody can be constructed and introduced into a cell either ex vivo or in vivo using well-established techniques including, but not limited to, transfection-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated cellular fusion, intracellular transportation of DNA-coated latex beads, protoplast fusion, viral infection, electroporation, a gene gun, and DEAE- or calcium phosphate-mediated transfection.

[0331] Effective in vivo dosages of an antibody are in the range of about 5 μg to about 50 $\mu\text{g}/\text{kg}$, about 50 μg to about 5 mg/kg, about 100 μg to about 500 $\mu\text{g}/\text{kg}$ of patient body weight, and about 200 to about 250 $\mu\text{g}/\text{kg}$ of patient body weight. For administration of polynucleotides encoding single-chain antibodies, effective in vivo dosages are in the range of about 100 ng to about 200 ng, 500 ng to about 50 mg, about 1 μg to about 2 mg, about 5 μg to about 500 μg , and about 20 μg to about 100 μg of DNA.

[0332] If the expression product is mRNA, the reagent is preferably an antisense oligonucleotide or a ribozyme. Polynucleotides which express antisense oligonucleotides or ribozymes can be introduced into cells by a variety of methods, as described above.

[0333] Preferably, a reagent reduces expression of a "BREAST CANCER GENE" gene or the activity of a "BREAST CANCER GENE" polypeptide by at least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the reagent. The effectiveness of the mechanism chosen to decrease the level of expression of a "BREAST CANCER GENE" gene or the activity of a "BREAST CANCER GENE" polypeptide can be assessed using methods well known in the art, such as hybridization of nucleotide probes to "BREAST CANCER GENE"-specific mRNA, quantitative RT-PCR, immunologic detection of a "BREAST CANCER GENE" polypeptide, or measurement of "BREAST CANCER GENE" activity.

[0334] In any of the embodiments described above, any of the pharmaceutical compositions of the invention can be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy can be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents can act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

[0335] Any of the therapeutic methods described above can be applied to any subject in need of such therapy, including, for example, birds and mammals such as dogs, cats, cows, pigs, sheep, goats, horses, rabbits, monkeys, and most preferably, humans.

[0336] All patents and patent applications cited in this disclosure are expressly incorporated herein by reference. The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific examples which are provided for purposes of illustration only and are not intended to limit the scope of the invention.

Pharmaceutical Compositions

[0337] The invention also provides pharmaceutical compositions which can be administered to a patient to achieve a therapeutic effect. Pharmaceutical compositions of the invention can comprise, for example, a "BREAST CANCER GENE" polypeptide, "BREAST CANCER GENE" polynucleotide, ribozymes or antisense oligonucleotides, antibodies which specifically bind to a "BREAST CANCER GENE" polypeptide, or mimetics, agonists, antagonists, or inhibitors of a "BREAST CANCER GENE" polypeptide activity. The compositions can be administered alone or in combination with at least one other agent, such as stabilizing compound, which can be administered in any sterile, biocompatible pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water. The compositions can be administered to a patient alone, or in combination with other agents, drugs or hormones.

[0338] In addition to the active ingredients, these pharmaceutical compositions can contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Pharmaceutical compositions of the invention can be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intraarterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, parenteral, topical, sublingual, or rectal means. Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

[0339] Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethylcellulose, or sodium carboxymethylcellulose; gums including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents can be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

[0340] Dragee cores can be used in conjunction with suitable coatings, such as concentrated sugar solutions, which also can contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments can be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

[0341] Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds can be dissolved or suspended in suitable liquids,

such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

[0342] Pharmaceutical formulations suitable for parenteral administration can be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions can contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds can be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Non-lipid polycationic amino polymers also can be used for delivery. Optionally, the suspension also can contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions. For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

[0343] The pharmaceutical compositions of the present invention can be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes. The pharmaceutical composition can be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation can be a lyophilized powder which can contain any or all of the following: 150 mM histidine, 0.1 %2% sucrose, and 27% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

[0344] Further details on techniques for formulation and administration can be found in the latest edition of REMINGTON'S PHARMACEUTICAL SCIENCES (182). After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. Such labeling would include amount, frequency, and method of administration.

Material and Methods

[0345] One strategy for identifying genes that are involved in breast cancer is to detect genes that are expressed differentially under conditions associated with the disease versus non-disease conditions. The sub-sections below describe a number of experimental systems which may be used to detect such differentially expressed genes. In general, these experimental systems include at least one experimental condition in which subjects or samples are treated in a manner associated with breast cancer, in addition to at least one experimental control condition lacking such disease associated treatment. Differentially expressed genes are detected, as described below, by comparing the pattern of gene expression between the experimental and control conditions.

[0346] Once a particular gene has been identified through the use of one such experiment, its expression pattern may be further characterized by studying its expression in a different experiment and the findings may be validated by an independent technique. Such use of multiple experiments may be useful in distinguishing the roles and relative importance of particular genes in breast cancer. A combined approach, comparing gene expression pattern in cells derived from breast cancer patients to those of *in vitro* cell culture models can give substantial hints on the pathways involved in development and/or progression of breast cancer.

[0347] Among the experiments which may be utilized for the identification of differentially expressed genes involved in malignant neoplasia and breast cancer, for example, are experiments designed to analyze those genes which are involved in signal transduction. Such experiments may serve to identify genes involved in the proliferation of cells.

[0348] Below are methods described for the identification of genes which are involved in breast cancer. Such represent genes which are differentially expressed in breast cancer conditions relative to their expression in normal, or non-breast cancer conditions or upon experimental manipulation based on clinical observations. Such differentially expressed genes represent "target" and/or "marker" genes. Methods for the further characterization of such differentially expressed genes, and for their identification as target and/or marker genes, are presented below.

[0349] Alternatively, a differentially expressed gene may have its expression modulated, i.e., quantitatively increased or decreased, in normal versus breast cancer states, or under control versus experimental conditions. The degree to which expression differs in normal versus breast cancer or control versus experimental states need only be large enough to be visualized via standard characterization techniques, such as, for example, the differential display technique described below. Other such standard characterization techniques by which expression differences may be visualized include but are not limited to quantitative RT-PCR and Northern analyses, which are well known to those of skill in the art.

EXAMPLE 1*Expression profiling**a) Expression profiling utilizing quantitative RT-PCR*

[0350] For a detailed analysis of gene expression by quantitative PCR methods, one will utilize primers flanking the genomic region of interest and a fluorescent labeled probe hybridizing in-between. Using the PRISM 7700 Sequence Detection System of PE Applied Biosystems (Perkin Elmer, Foster City, CA, USA) with the technique of a fluorogenic probe, consisting of an oligonucleotide labeled with both a fluorescent reporter dye and a quencher dye, one can perform such a expression measurement. Amplification of the probe-specific product causes cleavage of the probe, generating an increase in reporter fluorescence. Primers and probes were selected using the Primer Express software and localized mostly in the 3' region of the coding sequence or in the 3' untranslated region (see Table 5 for primer- and probe- sequences) according to the relative positions of the probe sequence used for the construction of the Affymetrix HG_U95A-E or HG-U133A-B DNA-chips. All primer pairs were checked for specificity by conventional PCR reactions. To standardize the amount of sample RNA, GAPDH was selected as a reference, since it was not differentially regulated in the samples analyzed. TaqMan validation experiments were performed showing that the efficiencies of the target and the control amplifications are approximately equal which is a prerequisite for the relative quantification of gene expression by the comparative $\Delta\Delta C_T$ method, known to those with skills in the art.

[0351] As well as the technology provided by Perkin Elmer one may use other technique implementations like Lightcycler™ from Roche Inc. or iCycler from Stratagene Inc..

b) Expression profiling utilizing DNA microarrays

[0352] Expression profiling can be carried out using the Affymetrix Array Technology. By hybridization of mRNA to such a DNA-array or DNA-Chip, it is possible to identify the expression value of each transcripts due to signal intensity at certain position of the array. Usually these DNA-arrays are produced by spotting of cDNA, oligonucleotides or sub-cloned DNA fragments. In case of Affymetrix technology app. 400.000 individual oligonucleotide sequences were synthesized on the surface of a silicon wafer at distinct positions. The minimal length of oligomers is 12 nucleotides, preferable 25 nucleotides or full length of the questioned transcript. Expression profiling may also be carried out by hybridization to nylon or nitrocellulose membrane bound DNA or oligonucleotides. Detection of signals derived from hybridization may be obtained by either colorimetric, fluorescent, electrochemical, electronic, optic or by radioactive readout. Detailed description of array construction have been mentioned above and in other patents cited. To determine the quantitative and qualitative changes in the chromosomal region to analyze, RNA from tumor tissue which is suspected to contain such genomic alterations has to be compared to RNA extracted from benign tissue (e.g. epithelial breast tissue, or micro dissected ductal tissue) on the basis of expression profiles for the whole transcriptome. With minor modifications, the sample preparation protocol followed the Affymetrix GeneChip Expression Analysis Manual (Santa Clara, CA). Total RNA extraction and isolation from tumor or benign tissues, biopsies, cell isolates or cell containing body fluids can be performed by using TRIzol (Life Technologies, Rockville, MD) and Oligotex mRNA Midi kit (Qiagen, Hilden, Germany), and an ethanol precipitation step should be carried out to bring the concentration to 1 mg/ml. Using 5-10 mg of mRNA to create double stranded cDNA by the Superscript system (Life Technologies). First strand cDNA synthesis was primed with a T7-(dT24) oligonucleotide. The cDNA can be extracted with phenol/chloroform and precipitated with ethanol to a final concentration of 1 mg /ml. From the generated cDNA, cRNA can be synthesized using Enzo's (Enzo Diagnostics Inc., Farmingdale, NY) *in vitro* Transcription Kit. Within the same step the cRNA can be labeled with biotin nucleotides Bio-11-CTP and Bio-16-UTP (Enzo Diagnostics Inc., Farmingdale, NY). After labeling and cleanup (Qiagen, Hilden (Germany) the cRNA then should be fragmented in an appropriated fragmentation buffer (e.g., 40 mM Tris-Acetate, pH 8.1, 100 mM KOAc, 30 mM MgOAc, for 35 minutes at 94 °C). As per the Affymetrix protocol, fragmented cRNA should be hybridized on the HG_U133 arrays A and B, comprising app. 40.000 probed transcripts each, for 24 hours at 60 rpm in a 45 °C hybridization oven. After Hybridization step the chip surfaces have to be washed and stained with streptavidin phycoerythrin (SAPE; Molecular Probes, Eugene, OR) in Affymetrix fluids stations. To amplify staining, a second labeling step can be introduced, which is recommended but not compulsive. Here one should add SAPE solution twice with an antistreptavidin biotinylated antibody. Hybridization to the probe arrays may be detected by fluorometric scanning (Hewlett Packard Gene Array Scanner; Hewlett Packard Corporation, Palo Alto, CA).

[0353] After hybridization and scanning, the microarray images can be analyzed for quality control, looking for major chip defects or abnormalities in hybridization signal. Therefore either Affymetrix GeneChip MAS 5.0 Software or other microarray image analysis software can be utilized. Primary data analysis should be carried out by software provided by the manufacturer..

[0354] In case of the genes analyses in one embodiment of this invention the primary data have been analyzed by further bioinformatic tools and additional filter criteria. The bioinformatic analysis is described in detail below.

c) Data analysis

[0355] According to Affymetrix measurement technique (Affymetrix GeneChip Expression Analysis Manual, Santa Clara, CA) a single gene expression measurement on one chip yields the average difference value and the absolute call. Each chip contains 16-20 oligonucleotide probe pairs per gene or cDNA clone. These probe pairs include perfectly matched sets and mismatched sets, both of which are necessary for the calculation of the average difference, or expression value, a measure of the intensity difference for each probe pair, calculated by subtracting the intensity of the mismatch from the intensity of the perfect match. This takes into consideration variability in hybridization among probe pairs and other hybridization artifacts that could affect the fluorescence intensities. The average difference is a numeric value supposed to represent the expression value of that gene. The absolute call can take the values 'A' (absent), 'M' (marginal), or 'P' (present) and denotes the quality of a single hybridization. We used both the quantitative information given by the average difference and the qualitative information given by the absolute call to identify the genes which are differentially expressed in biological samples from individuals with breast cancer versus biological samples from the normal population. With other algorithms than the Affymetrix one we have obtained different numerical values representing the same expression values and expression differences upon comparison.

[0356] The differential expression E in one of the breast cancer groups compared to the normal population is calculated as follows. Given n average difference values d_1, d_2, \dots, d_n in the breast cancer population and m average difference values c_1, c_2, \dots, c_m in the population of normal individuals, it is computed by the equation:

$$E = \exp\left(\frac{1}{m} \sum_{i=1}^m \ln(c_i) - \frac{1}{n} \sum_{i=1}^n \ln(d_i)\right)$$

If $d_i < 50$ or $c_i < 50$ for one or more values of i and j, these particular values c_i and/or d_j are set to an "artificial" expression value of 50. These particular computation of E allows for a correct comparison to TaqMan results.

[0357] A gene is called up-regulated in breast cancer versus normal if $E \geq 1.5$ and if the number of absolute calls equal to 'P' in the breast cancer population is greater than $n/2$.

[0358] A gene is called down-regulated in breast cancer versus normal if $E \leq 1.5$ and if the number of absolute calls equal to 'P' in the normal population is greater than $m/2$.

[0359] The final list of differentially regulated genes consists of all up-regulated and all down-regulated genes in biological samples from individuals with breast cancer versus biological samples from the normal population. Those genes on this list which are interesting for a pharmaceutical application were finally validated by TaqMan. If a good correlation between the expression values/behavior of a transcript could be observed with both techniques, such a gene is listed in Tables 1 to 3.

[0360] Since not only the information on differential expression of a single gene within an identified ARCHEON, but also the information on the co-regulation of several members is important for predictive, diagnostic, preventive and therapeutic purposes we have combined expression data with information on the chromosomal position (e.g. golden path) taken from public available databases to develop a picture of the overall transcriptome of a given tumor sample. By this technique not only known or suspected regions of genomes can be inspected but even more valuable, new regions of dysregulation with chromosomal linkage can be identified. This is of value in other types of neoplasia or viral integration and chromosomal rearrangements. By SQL based database searches one can retrieve information on expression, qualitative value of a measurement (denoted by Affymetrix MAS 5.0 Software), expression values derived from other techniques than DNA-chip hybridization and chromosomal linkage.

EXAMPLE 2

Identification of the ARCHEON

a) Identification and localization of genes or gene probes (represented by the so called probe sets on Affymetrix arrays HG-U95A-E or HG-U133A-B) in their chromosomal context and order on the human genome.

[0361] For identification of larger chromosomal changes or aberrations, as they have been described in detail above,

a sufficient number of genes, transcripts or DNA-fragments is needed. The density of probes covering a chromosomal region is not necessarily limited to the transcribed genes, in case of the use of array based CGH but by utilizing RNA as probe material the density is given by the distance of genes on a chromosome. The DNA-microarrays provided by Affymetrix Inc. do contain hitherto all transcripts from the known humane genome, which are represented by 40.000
 5 -60.000 probe sets. By BLAST mapping and sorting the sequences of these short DNA-oligomers to the public available sequence of the human genome represented by the so called "golden path", available at the university of California in Santa Cruz or from the NCBI, a chromosomal display of the whole Transcriptome of a tissue specimen evolves. By graphical display of the individual chromosomal regions and color coding of over or under represented transcripts, compared to a reference transcriptome regions with DNA gains and losses can be identified.

10 b) Quantification of gene copy numbers by combined IHC and quantitative PCR (PCR karyotyping) or directly by quantitative PCR

15 [0362] Usually one to three paraffin-embedded tissue sections that are 5 µm thick are used to obtain genomic DNA from the samples. Tissue section are stained by colorimetric IHC after deparaffinization to identify regions containing disease associated cells. Stained regions are macrodissected with a scalpel and transferred into a microcentrifuge tube. The genomic DNA of these isolated tissue sections is extracted using appropriate buffers. The isolated DNA is then used for quantitative PCR with appropriate primers and probes. Optionally the IHC staining can be omitted and the genomic DNA can be directly isolated with or without prior deparaffinization with appropriate buffers. Those who
 20 are skilled in the art may vary the conditions and buffers described below to obtain equivalent results.

[0363] Reagents from DAKO (HerceptTest Code No. K 5204) and TaKaRa were used (Biomedicals Cat.: 9091) according to the manufactures protocol.

[0364] It is convenient to prepare the following reagents prior to staining:

25 **Solution No. 7**

[0365] Epitope Retrieval Solution (Citrate buffer + antimicrobial agent) (10xconc.) 20 ml ad 200 ml aqua dest. (stable for 1month at 2-8°C)

30 **Solution No. 8**

[0366] Washing-buffer (Tris-HCl + antimicrobial agent) (10 x conc.) 30 ml ad 300 ml destilled water (stable for 1month at 2-8°C)

35 **Staining solution: DAB**

[0367] 1 ml solution is sufficient for 10 slides. The solution were prepared immediately before usage.:

40 [0368] 1 ml DAB buffer (Substrate Buffer solution, pH 7.5, containing H₂O₂, stabilizer, enhancers and an antimicrobial agent) + 1 drop (25-3 µl) DAB-Chromogen (3,3'-diaminobenzidine chromogen solution). This solution is stable for up to 5 days at 2-8°C. Precipitated substances do not influence the staining result. Additionally required are: 2x approx. 100 ml Xyol, 2x approx. 100 ml Ethanol 100%, 2x Ethanol 95%, aqua dest. These solution can be used for up to 40 stainings. A water bath is required for the epitope retrieval step.

45 Staining procedure:

[0369] All reagents are pre-warmed to room temperature (20-25°C) prior to immuno-staining. Likewise all incubations were performed at room temperature. Except the epitope retrieval which is performed in at 95°C water bath. Between the steps excess of liquid is tapped off from the slides with lintless tissue (Kim Wipe).

50 Deparaffinization

55 [0370] Slides are placed in a xylene bath and incubated for 5 minutes. The bath is changed and the step repeated once. Excess of liquid is tapped off and the slides are placed in absolute ethanol for 3 minutes. The bath is changed and the step repeated once. Excess of liquid is tapped off and the slides are placed in 95% ethanol for 3 minutes. The bath is changed and the step repeated once. Excess of liquid is tapped off and the slides are placed in distilled water for a minimum of 30 seconds.

Epitope Retrieval

[0371] Staining jars are filled with diluted epitope retrieval solution and preheated in a water bath at 95° C. The deparaffinized sections are immersed into the preheated solution in the staining jars and incubated for 40 minutes at 95° C. The entire jar is removed from the water bath and allowed to cool down at room temperature for 20 minutes. The epitope retrieval solution is decanted, the sections are rinsed in distilled water and finally soaked in wash buffer for 5 minutes.

Peroxidase Blocking:

[0372] Excess of buffer is tapped off and the tissue section encircled with a DAKO pen. The specimen is covered with 3 drops (100 µl) Peroxidase-Blocking solution and incubated for 5 minutes. The slides are rinsed in distilled water and placed into a fresh washing buffer bath.

Antibody Incubation

[0373] Excess of liquid is tapped off and the specimen are covered with 3 drops (100 µl) of Anti-Her-2/neu reagent (Rabbit Anti-Human Her2 Protein in 0.05 mol/L Tris/HCl, 0.1 mol/L NaCl, 15 mmol/L pH7.2 Na₂S₂O₃ containing stabilizing protein) or negative control reagent (= IGG fraction of normal rabbit serum at an equivalent protein concentration as the Her2 Ab). After 30 minutes of incubation the slide is rinsed in water and placed into a fresh water bath.

Visualization

[0374] Excess of liquid is tapped off and the specimen are covered with 3 drops (100 µl) of visualization reagent. After 30 minutes of incubation the slide is rinsed in water and placed into a fresh water bath. Excess of liquid is tapped off and the specimen are covered with 3 drops (100 µl) of Substrate-Chromogen solution (DAB) for 10 minutes. After rinsing the specimen with distilled water, photographs are taken with a conventional Olympus microscope to document the staining intensity and tumor regions within the specimen. Optionally a counterstain with hematoxylin was performed.

DNA extraction

[0375] The whole specimens or dissected subregions are transferred into a microcentrifuge tubes. Optionally a small amount (10µl) of preheated TaKaRa solution (DEXPAT™) is preheated and placed onto the specimen to facilitate sample transfer with a scalpel. 50 to 150 µl of TaKaRa solution were added to the samples depending on the size of the tissue sample selected. The sample are incubated at 100° C for 10 minutes in a block heater, followed by centrifugation at 12.000 rpm in a microcentrifuge. The supernatant is collected using a micropipette and placed in a separate microcentrifuge tube. If no deparaffinization step has been undertaken one has to be sure not to withdraw tissue debris and resin. Genomic DNA left in the pellet can be collected by adding resin-free TaKaRa buffer and an additional heating and centrifugation step. Samples are stored at -20° C.

[0376] Genomic DNA from different tumor cell lines (MCF-7, BT-20, BT-474, SKBR-3, AU-565, UACC-812, UACC-893, HCC-1008, HCC-2157, HCC-1954, HCC-2218, HCC-1937, HCC1599, SW480), or from lymphocytes is prepared with the QIAamp® DNA Mini Kits or the QIAamp® DNA Blood Mini Kits according to the manufacturers protocol. Usually between 1ng up to 1µg DNA is used per reaction.

Quantitative PCR

[0377] To measure the gene copy number of the genes within the patient samples the respective primer/probes (see table below) are prepared by mixing 25 µl of the 100 µM stock solution "Upper Primer", 25 µl of the 100 µM stock solution "Lower Primer" with 12,5 µl of the 100 µM stock solution Taq Man Probe (Quencher Tamra) and adjusted to 500 µl with aqua dest. For each reaction 1,25 µl DNA-Extract of the patient samples or 1,25 µl DNA from the cell lines were mixed with 8,75 µl nuclease-free water and added to one well of a 96 Well-Optical Reaction Plate (Applied Biosystems Part No. 4306737). 1,5 µl Primer/Probe mix, 12, µl Taq Man Universal-PCR Mix (2x) (Applied Biosystems Part No. 4318157) and 1 µl Water are then added. The 96 well plates are closed with 8 Caps/Strips (Applied Biosystems Part Number 4323032) and centrifuged for 3 minutes. Measurements of the PCR reaction are done according to the instructions of the manufacturer with a TaqMan 7900 HT from Applied Biosystems (No. 20114) under appropriate conditions (2 min. 50° C, 10 min. 95° C, 0.15min. 95° C, 1 min. 60° C; 40 cycles). Software SDS 2.0 from Applied Biosystems is used according to the respective instructions. CT-values are then further analyzed with appropriate software (Microsoft Excel™).

REFERENCES

[0378]

5	<i>Patents cited</i>	
	U.S. Pat. No. 4,843,155	Chomczynski, P.
	U.S. Pat. No. 5,262,31	Liang, P., and Pardee, A. B., 1993
	U.S. Pat. No. 4,683,202	Mullis, K. B., 1987
10	U.S. Pat. No. 5,593,839	
	U.S. Pat. No. 5,578,832	
	U.S. Pat. No. 5,556,752	
	U.S. Pat. No. 5,631,734	
15	U.S. Pat. No. 5,599,695	
	U.S. Pat. No. 4,683,195	
	U.S. Pat. No. 5,498,531	
	U.S. Pat. No. 5,714,331	
	U.S. Pat. No. 5,641,673	Haseloff et al.,
20	U.S. Pat. No. 5,223,409	Lander, E.,
	U.S. Pat. No. 5,976,813	Beutel et al.
	U.S. Pat. No. 5,283,317	
	U.S. Pat. No. 6,203,987	
25	U.S. Pat. No. 6,379,895	
	WO 97/29212	
	WO 97/27317	
	WO 95/22058	
	WO 99/12826	
30	WO 97/02357	
	WO 94/13804	
	WO 94/10300	
	WO 97/14028	
	WO 99/52708	
35	EP 0 785 280	
	EP 0 799 897	
	EP 0 728 520	
	EP 0 721 016	
40	EP 0 321 201	

GB2188638B

Publications cited

- 45
- [0379]
- (1) Gusterson et al., Journal of Clinical Oncology 10, 1049-1056, 1992
- (2) Achuthan et al., Cancer Genet Cytogenet. 130:166-72, 2001
- 50 (3) Tomasetto et al., FEBS Lett. 373: 245-249, 1995
- (4) Pragnelli et al., FEBS Lett. 291: 253-258, 1991
- (5) Nakamichi et al. 1986
- (6) Feo et al., Proc. Nat. Acad. Sci. 86: 6691-6695, 1989
- (7) Davies et al., Proc. Nat. Acad. Sci. 86: 6691-6695, 1989
- 55 (8) Lee et al., Molec. Endocr. 9: 243-254, 1995
- (9) Drane et al., Oncogene 15: 3013-3024, 1997
- (10) Zhu et al., J. Biol. Chem. 272: 25500-25506, 1997
- (11) Yuan et al., Proc. Nat. Acad. Sci. 95: 7939-7944, 1998

- (12) Zhu et al., Proc. Nat. Acad. Sci. 96: 10848-10853, 1999
- (13) Lee et al., Science 268: 836-844, 1995
- (14) McCormick et al., Molec. Cell. Biol. 16: 5792-5800, 1996
- (15) Tamimi et al., Genomics 40: 355-357, 1997
- 5 (16) Valle et al., *FEBS Lett.* 415: 163-168, 1997
- (17) Kaneda et al., J. Biol. Chem. 263: 7672-7677, 1988
- (18) Hoehe et al., Hum. Molec. Genet. 1:175-178, 1992
- (19) Yang-Feng et al., Abstract Cytogenet. Cell Genet. 40: 784, 1985
- (20) Coussens et al., Science 230: 1132-1139, 1985
- 10 (21) van de Vijver et al., New Eng. J. Med. 319: 1239-1245, 1988
- (22) Slamon et al., Science 244: 707-712, 1989
- (23) Fukushima et al., Res. Commun. 134: 477-483, 1986
- (24) Kaneko et al., Jpn. J. Cancer Res. 78: 16-19, 1987
- (25) Di Fiore et al., Science 237: 178-182, 1987
- 15 (26) Popescu et al., Genomics 4: 362-366, 1989
- (27) Qiu et al., Nature 393: 83-85, 1998
- (28) Yu et al., Molec. Cell 2: 581-591, 1998
- (29) Doherty et al., Proc. Nat. Acad. Sci. 96: 10869-10874, 1999
- (30) Slamon et al., New Eng. J. Med. 344: 783-792, 2001
- 20 (31) Margolis et al., J. Clin. Invest. 102: 821-827, 1998
- (32) Tanaka et al., J. Clin. Invest. 102: 821-827, 1998
- (33) Dong et al., J. Biol. Chem. 272: 29104-29112, 1997
- (34) Stein et al., EMBO J.13:1331-40, 1994
- (35) Nagata et al., Nature 319: 415-418, 1986
- 25 (36) Le Beau et al., Leukemia 1: 795-799, 1987,
- (37) Jansson et al., EMBO J. 2: 561-565, 1983
- (38) Thompson et al., Science 237:1610-1614, 1987
- (39) Nakai et al., Proc. Nat. Acad. Sci. 85: 2781-2785, 1988
- (40) Miyajima et al., Cell 57: 31-39, 1989
- 30 (41) Debuire et al., Science 224: 1456-1459, 1984
- (42) Petkovich et al., Nature 330: 444-450, 1987
- (43) Mattei et al., Hum. Genet. 80: 186-188, 1988,
- (44) Williams et al., Molec. Cell. Biol. 18: 2758-2767, 1998
- (45) Saha et al., Molec. Cell. Biol. 18: 2758-2767, 1998
- 35 (46) Yan et al., Proc. Nat. Acad. Sci. 95: 3603-3608, 1998
- (47) Singh et al., Nucleic Acids Res. 16: 3919-3929, 1988
- (48) Tsai-Pflugfelder et al., Proc. Nat. Acad. Sci. 85: 7177-7181, 1988
- (49) Chung et al., Proc. Nat. Acad. Sci. 86: 9431-9435, 1989
- (50) Lang et al., Gene 221: 255-266, 1998
- 40 (51) Keith et al., Genes Chromosomes Cancer 4: 169-175, 1992
- (52) Kingsmore et al., Mammalian Genome 4: 288-289, 1993
- (53) Watt et al., Biochem. J. 303: 681-695, 1994
- (54) Kiefer et al., J. Biol. Chem. 267: 12692-12699, 1992
- (55) Shimasaki et al., Molec. Endocr. 4: 1451-1458, 1990
- 45 (56) Zazzi et al., Genomics 49: 401-410, 1998
- (57) Bajalica et al., Hum. Genet. 89: 234-236, 1992
- (58) Tonin et al., Genomics 18: 414-417, 1993
- (59) Birkenbach et al., J. Virol. 67: 2209-2220, 1993
- (60) Wang et al., Proc. Nat. Acad. Sci. 95: 492-498, 1998
- 50 (61) Klochendler-Yeivin et al., Curr Opin Genet Dev 12:73-9,2002
- (62) Ring et al., Genomics 51:140-3, 1998
- (63) Darmon et al., Molec. Biol. Rep. 12: 277-283, 1987
- (64) Zhou et al., J. Biol. Chem. 263: 15584-15589, 1988
- (65) Korge et al., Proc. Nat. Acad. Sci. 89: 910-914, 1992
- 55 (66) Lessin et al., J. Invest. Derm. 91: 572-578, 1988
- (67) Romano et al., Cytogenet. Cell Genet. 58: 2009-2010, 1991.
- (68) Fuchs et al., Proc. Nat. Acad. Sci. 89: 6906-6910, 1992
- (69) Rogaev et al., Nature Genet. 5: 158-162, 1993

- (70) Liu et al., Curr. Eye Res. 12: 963-974, 1993
- (71) Nishida et al., Invest. Ophthalm. Vis. Sci. 37: 1800-1809, 1996
- (72) Nishida et al., Am. J. Hum. Genet. 61: 1268-1275, 1997
- (73) Meesmann and Wilke, 1939
- 5 (74) Corden et al., Exp. Eye Res. 70: 41-49, 2000
- (75) van de Vijver et al., Mol Cell Biol 7, 2091-23, 1987
- (76) Offlerding et al., Biochem Biophys Res Comm 251, 907-13, 1988
- (77) Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2d ed., 1989
- (78) Ausubel et al., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, N.Y., 1989.
- 10 (79) Tedder, T. F. et al., Proc. Natl. Acad. Sci. U.S.A. 85:208-212, 1988
- (80) Hedrick, S. M. et al., Nature 308:149-153, 1984
- (81) Lee, S. W. et al., Proc. Natl. Acad. Sci. U.S.A. 88:4225, 1984
- (82) Sarkar, PCR Methods Applic. 2, 318-322, 1993
- 15 (83) Triglia et al., Nucleic Acids Res. 16, 81-86, 1988
- (84) Lagerstrom et al., PCR Methods Applic. 1, 111-119, 1991
- (85) Copeland & Jenkins, Trends in Genetics 7: 113-118, 1991
- (86) Cohen, et al., Nature 366: 698-701, 1993
- (87) Bonner et al., J. Mol. Biol. 81, 123 1973
- 20 (88) Bolton and McCarthy, Proc. Natl. Acad. Sci. U.S.A. 48, 1390 1962
- (89) Plump et al., Cell 71: 343-353, 1992
- (90) Altschul et al., Bull. Math. Bio. 48:603, 1986,
- (91) Henikoff & Henikoff, Proc. Natl. Acad. Sci. USA 89:10915, 1992
- (92) Pearson & Lipman, Proc. Nat'l Acad. Sci. USA 85:2444, 1988
- 25 (93) Pearson et al., Meth. Enzymol. 183:63, 1990
- (94) Needleman & Wunsch, J. Mol. Biol.48:444, 1970
- (95) Sellers, SIAM J. Appl. Math.26:787, 1974
- (96) Takamatsu, EMBO J. 6, 307-311, 1987
- (97) Coruzzi et al., EMBO J. 3, 1671-1680, 1984
- 30 (98) Broglie et al., Science 224, 838-843, 1984
- (99) Winter et al., Results Probl. Cell Differ. 17, 85-105, 1991
- (100) Engelhard et al., Proc. Nat. Acad. Sci. 91, 3224-3227, 1994
- (101) Logan & Shenk, Proc. Natl. Acad. Sci. 81, 3655-3659, 1984
- (102) Scharf et al., Results Probl. Cell Differ. 20, 125-162, 1994
- 35 (103) Freshney R.L., ed., ANIMAL CELL CULTURE , 1986
- (104) Wigler et al., Cell 11, 223-232, 1977
- (105) Lowy et al., Cell 22, 817-823, 1980
- (106) Wigler et al., Proc. Natl. Acad. Sci. 77, 3567-3570, 1980
- (107) Colbere-Garapin et al., J. Mol. Biol. 150, 114, 1981
- 40 (108) Hartman & Mulligan, Proc. Natl. Acad. Sci. 85, 8047-8051, 1988
- (109) Rhodes et al., Methods Mol. Biol. 55, 121-131, 1995
- (110) Hampton et al., SEROLOGICAL METHODS: A LABORATORY MANUAL, APS Press, St. Paul, Minn., 1990
- (111) Maddox et al., J. Exp. Med. 158, 1211-1216, 1983
- (112) Porath et al., Prot. Exp. Purif. 3, 263-281, 1992
- 45 (113) Kroll et al., DNA Cell Biol. 12, 441-453, 1993
- (114) Caruthers et al., Nud. Acids Res. Symp. Ser. 215-223, 1980
- (115) Horn et al. Nud. Acids Res. Symp. Ser. 225-232, 1980
- (116) Merrifield, J. Am. Chem. Soc. 85, 2149-2154, 1963
- (117) Roberge et al., Science 269, 202-204, 1995
- 50 (118) Creighton, PROTEINS: STRUCTURES AND MOLECULAR PRINCIPLES, WH and Co., New York, N.Y., 1983
- (119) Cronin et al., Human Mutation 7:244, 1996
- (120) Landegran et al., Science 241:1077-1080, 1988
- (121) Nakazawa et al., PNAS 91:360-364, 1994
- (122) Abravaya et al., Nuc Acid Res 23:675-682, 1995
- 55 (123) Guatelli, J.C. et al., Proc. Natl. Acad. Sci. USA 87:1874-1878, 1990
- (124) Kwoh, D.Y. et al., Proc. Natl. Acad. Sci. USA 86:1173-1177, 1989
- (125) Lizardi, P.M. et al., Bio/Technology 6:1197, 1988
- (126) Brown, Meth. Mol. Biol. 20, 18, 1994

- (127) Sonveaux, Meth. Mol. Biol. 26, 1-72, 1994
 (128) Uhlmann et al., Chem. Rev. 90, 543-583, 1990
 (129) Gee et al., in Huber & Carr, MOLECULAR AND IMMUNOLOGIC APPROACHES, Publishing Co., Mt. Kisco, N.Y., 1994
- 5 (130) Agrawal et al., Trends Biotechnol. 10, 152-158, 1992
 (131) Uhlmann et al., Tetrahedron. Lett. 215, 3539-3542, 1987
 (132) Cech, Science 236, 1532-1539, 1987
 (133) Cech, Ann. Rev. Biochem. 59, 543-568, 1990
 (134) Couture & Stinchcomb, Trends Genet. 12, 510-515, 1996
- 10 (135) Haseloff et al. Nature 334, 585-591, 1988
 (136) Kohler et al., Nature 256,495-497,1985
 (137) Kozbor et al., J. Immunol. Methods 81, 3142, 1985
 (138) Cole et al., Proc. Natl. Acad. Sci. 80, 2026-2030, 1983
 (139) Cole et al., Mol. Cell Biol. 62, 109-120, 1984
- 15 (140) Morrison et al., Proc. Natl. Acad. Sci. 81, 6851-6855, 1984
 (141) Neuberger et al., Nature 312, 604-608, 1984
 (142) Takeda et al., Nature 314, 452-454, 1985
 (143) Burton, Proc. Natl. Acad. Sci. 88, 11120-11123, 1991
 (144) Thirion et al., Eur. J. Cancer Prev. 5, 507-11, 1996
- 20 (145) Coloma & Morrison, Nat. Biotechnol. 15, 159-63, 1997
 (146) Mallender & Voss, J. Biol. Chem. 269, 199-206, 1994
 (147) Verhaar et al., Int. J. Cancer 61, 497-501, 1995
 (148) Nicholls et al., J. Immunol. Meth. 165, 81-91, 1993
 (149) Orlandi et al., Proc. Natl. Acad. Sci. 86, 3833-3837, 1989
- 25 (150) Winter et al., Nature 349, 293-299, 1991
 (151) Lam, Anticancer Drug Des. 12, 145, 1997
 (152) DeWitt et al., Proc. Natl. Acad. Sci. U.S.A. 90, 6909, 1993
 (153) Erb et al. Proc. Natl. Acad. Sci. U.S.A. 91, 11422, 1994
 (154) Zuckermann et al., J. Med. Chem. 37, 2678, 1994
- 30 (155) Cho et al., Science 261, 1303, 1993
 (156) Carell et al., Angew. Chem. Int. Ed. Engl. 33, 2059 & 2061, 1994
 (157) Gallop et al., J. Med. Chem. 37, 1233, 1994
 (158) Houghten, BioTechniques 13, 412-421, 1992
 (159) Lam, Nature 354, 8284, 1991
- 35 (160) Fodor, Nature 364, 555-556, 1993
 (161) Cull et al., Proc. Natl. Acad. Sci. U.S.A. 89, 1865-1869, 1992
 (162) Scott & Smith, Science 249, 386-390, 1990
 (163) Devlin, Science 249, 404-406, 1990
 (164) Cwiria et al., Proc. Natl. Acad. Sci. 97, 6378-6382, 1990
- 40 (165) Felici, J. Mol. Biol. 222, 301-310, 1991
 (166) Jayawickreme et al., Proc. Natl. Acad. Sci. U.S.A. 19, 1614-1618, 1994
 (167) Chelsky, Strategies for Screening Combinatorial Libraries 1995
 (168) Salmon et al., Molecular Diversity 2, 57-63, 1996
 (169) McConnell et al., Science 257, 1906-1912, 1992
- 45 (170) Sjolander & Urbaniczky, Anal. Chem. 63, 2338-2345, 1991
 (171) Szabo et al., Curr. Opin. Struct. Biol. 5, 699-705, 1995
 (172) Zervos et al., Cell 72, 223-232, 1993
 (173) Madura et al., J. Biol. Chem. 268, 12046-12054, 1993
 (174) Bartel et al., BioTechniques 14, 920-924, 1993
- 50 (175) Iwabuchi et al., Oncogene 8, 1693-1696, 1993
 (176) Findeis et al. Trends in Biotechnol. 11, 202-205, 1993
 (177) Chiou et al., GENE THERAPEUTICS: METHODS AND APPLICATIONS OF DIRECT GENE TRANSFER J. A. Wolf ed., 1994
- (178) Wu & Wu, J. Biol. Chem. 263, 621-24, 1988
 (179) Wu et al., J. Biol. Chem. 269, 542-46, 1994
 (180) Zenke et al., Proc. Natl. Acad. Sci. U.S.A. 87, 3655-59, 1990
 (181) Wu et al., J. Biol. Chem. 266, 338-42, 1991
 (182) REMINGTON'S PHARMACEUTICAL SCIENCES Maack Publishing Co., Easton, Pa.

EP 1 365 034 A2

(183) Hille, Excitable Membranes, Sunderland, MA, Sinauer Associates, Inc.

Table 1

DNA SEQ ID NO:	Protein SEQ ID NO:	Genbank ID	Unigene_v133_ID	Locus Link ID	Gene Name
1	27	NM_006148.1	75080	3927	LASP1
2	28	NM_000723.1	635	782	CACNB1
3	29	NM_000981.1	252723	6143	RPL19RPL19
4	30	Y13467	15589	5469	PPARGBP
5	31	NM_016507.1	123073		CrkRS
6	32	AB021742.1	322431	4761	NEUROD2
7	33	NM_006804.1	77628	10948	MLN64
8	34	NM_003673.1	111110	8557	TELETHONIN
9	35	NM_002686.1	1892	5409	PNMT
10	36	X03363.1	323910	2064	ERBB2
11	37	AB008790.1	86859	2886	GRB7
12	38	NM_002809.1	9736	5709	PSMD3
13	39	NM_000759.1	2233	1440	GCSFG
14	40	A1023317	23106	9862	KIAA0130
15	41	X55005		7067	c-erbA-1
16	42	X72631	211606	9572	NR1D1
17	43	NM_007359.1	83422	22794	MLN51
18	44	U77949.1	69563	990	CDC6
19	45	U41742.1		5914	RARA
20	46	NM_001067.1	156346	7153	TOP2A
21	47	NM_001552.1	1516		IGFBP4
22	48	NM_001838.1	1652		CCR7 EBI1
23	49	NM_003079.1	332848	6605	SMARCE1 BAF57
24	50	X14487	99936	3858	KRT10
25	51	NM_000223.1	66739		KRT12
26	52	NM_002279.2	32950	3884	hHka3-II
53	76	NM_005937	349196	4302	MLLT6
54	77	XM_008147	184669	7703	ZNF144
55	78	NM_138687	432736	8396	PIP5K2B
56	79	NM_020405	125036	57125	TEM7
57	80	XM_012694	258579	22806	ZNFN1A3
58	81	XM_085731	13996	147179	WIRE
59	82	NM_002795	82793	5691	PSMB3
60	83	NM_033419	91668	93210	MGC9753 Variant a

Table 1 (continued)

DNA SEQ ID NO:	Protein SEQ ID NO:	Genbank ID	Unigene_v133_ID	Locus Link ID	Gene Name
61	84				MGC9753 Variant c
62	85				MGC9753 Variant d
63	86				MGC9753 Variant e
64	87				MGC9753 Variant g
65	88				MGC9753 Variant h
66	89				MGC9753 Variant i
67	90	AF395708	374824	94103	ORMDL3
68	91	NM_032875	194498	84961	MGC15482
69	92	NM_032192	286192	84152	PPP1R1B
70	93	NM_032339	333526	84299	MGC14832
71	94	NM_057555	12101	51242	LOC51242
72	95	NM_017748	8928	54883	FLJ20291
73	96	NM_018530	19054	55876	Pro2521
74	97	NM_016339	118562	51195	Link-GEFII
75	98	NM_032865	294022	84951	CTEN

Table 2

DNA SEQ ID NO:	Gene description
1	Member of a subfamily of LIM proteins that contains a LIM domain and an SH3 (Src homology region 3) domain
2	Beta 1 subunit of a voltage-dependent calcium channel (dihydropyridine receptor), involved in coupling of excitation and contraction in muscle, also acts as a calcium channel in various other tissues
3	Ribosomal protein L19, component of the large 60S ribosomal subunit
4	Protein with similarity to nuclear receptor-interacting proteins; binds and co-activates the nuclear receptors PPARalpha (PPARA), RARalpha (RARA), RXR, TRbeta1, and VDR
5	we26e0 CDC2-related protein kinase 7
6	Neurogenic differentiation, a basic-helix-loop-helix transcription factor that mediates neuronal differentiation
7	Protein that is overexpressed in malignant tissues, contains a putative trans-membrane region and a SIAR Homology Domain (SHD), may function in steroidogenesis and contribute to tumor progression
8	Telethonin, a sarcomeric protein specifically expressed in skeletal and heart muscle, caps titin (TTN) and is important for structural integrity of the sarcomere

Table 2 (continued)

DNA SEQ ID NO:	Gene description
5 9	Phenylethanolamine N-methyltransferase, acts in catecholamine biosynthesis to convert norepinephrine to epinephrine
10 10	Tyrosine kinase receptor that has similarity to the EGF receptor, a critical component of IL-6 signaling through the MAP kinase pathway, overexpression associated with prostate, ovary and breast cancer
10 11	Growth factor receptor-bound protein, an SH2 domain-containing protein that has isoforms which may have a role in cell invasion and metastatic progression of esophageal carcinomas
12	Non-ATPase subunit of the 26S proteasome (prosome, macropain)
15 13	Granulocyte colony stimulating factor, a glycoprotein that regulates growth, differentiation, and survival of neutrophilic granulocytes
14	Member of the Vitamin D Receptor Interacting Protein co-activator complex, has strong similarity to thyroid hormone receptor-associated protein (murine Trap 100) which function as a transcriptional coregulator
20 15	Thyroid hormone receptor alpha, a high affinity receptor for thyroid hormone that activates transcription; homologous to avian erythroblastic leukemia virus oncogene
16	encoding Rev-ErbAa1p nuclear receptor subfamily 1, group D, member 1
17	Protein that is overexpressed in breast carcinomas
25 18	Protein which interacts with the DNA replication proteins PCNA and Orc1, translocates from the nucleus following onset of S phase; S. cerevisiae homolog Cdc6p is required for initiation of S phase
19	Retinoic acid receptor alpha, binds retinoic acid and stimulates transcription in a ligand-dependent manner
30 20	DNA topoisomerase II alpha, member of a family of proteins that relieves torsional stress created by DNA replication, transcription, and cell division;
35 21	Insulin-like growth factor binding protein, the major IGFBP of osteoblast-like cells, binds IGF1 and IGF2 and inhibits their effects on promoting DNA and glycogen synthesis in osteoblastic cells
22	HUMEB103 G protein-coupled receptor (EBI 1) gene exon 3 chemokine (C-C motif) receptor 7 G protein-coupled receptor
40 23	Protein with an HMG 1/2 DNA-binding domain that is subunit of the SNF/SWI complex associated with the nuclear matrix and implicated in regulation of transcription by affecting chromatin structure
45 24	Keratin 10, a type I keratin that is a component of intermediate filaments and is expressed in terminally differentiated epidermal cells; mutation of the corresponding gene causes epidermolytic hyperkeratosis
25	Keratin 12, a component of intermediate filaments in corneal epithelial cells; mutation of the corresponding gene causes Meesmann corneal dystrophy
50 26	Hair keratin 3B, a type I keratin that is a member of a family of structural proteins that form intermediate filaments
53	MLLT6 Myeloid/lymphoid or mixed-lineage leukemia (trithorax homolog, Drosophila); translocated to, 6
54	zinc finger protein 144 (Mel-18)
55 55	phosphatidylinositol-4-phosphate 5-kinase type II beta isoform a
56	tumor endothelial marker 7 precursor

EP 1 365 034 A2

Table 2 (continued)

DNA SEQ ID NO:	Gene description
57	zinc finger protein, subfamily 1A, 3
58	WASP-binding protein putative cr16 and wip like protein similar to Wiskott-Aldrich syndrome protein
59	proteasome (prosome, macropain) subunit, beta type, 3
60	Predicted
67	ORM1-like 3 (<i>S. cerevisiae</i>)
68	F-box domain A Receptor for Ubiquitination Targets
69	protein phosphatase 1, regulatory (inhibitor) subunit 1B (dopamine and cAMP regulated phosphoprotein, DARPP-32)
70	Predicted Protein
71	Predicted Protein
72	Predicted Protein
73	Predicted Protein
74	Link-GEFII: Link guanine nucleotide exchange factor II
75	C-terminal tensin-like

Table 3

DNA SEQ ID NO:	Gene function	Subcellular localization
1	SH3/SH2 adapter protein	-
2	voltage-gated calcium channel membrane fraction Channel [passive transporter]	Plasma membrane
3	RNA binding structural protein of ribosome protein biosynthesis	Cytoplasm
4	transcription co-activator nucleus Pol II transcription	Nucleus
5	-	-
6	transcription factor transcription regulation from Pol II promoter neurogenesis	-
7	mitochondrial transport steroid and lipid metabolism	Cytoplasm
8	structural protein of muscle sarcomere alignment	Cytoplasm
9	phenylethanolamine N-methyltransferase Transferase	-
10	Neu/ErbB-2 receptor receptor signaling protein tyrosine kinase	Plasma membrane
11	SH3/SH2 adapter protein EGF receptor signaling pathway	Cytoplasm
12	26S proteasome Protein degradation Proteasome subunit	Cytoplasm
13	developmental processes positive control of cell proliferation	Extracellular space
14	fatty acid omega-hydroxylase fatty acid omega-hydroxylase	-
15	DNA-binding protein Transcription factor	Nucleus
16	steroid hormone receptor transcription co-repressor	Nucleus
17	-	-
18	nucleotide binding cell cycle regulator DNA replication checkpoint regulation of CDK activity	nucleus
19	retinoic acid receptor transcription co-activator transcription factor	nucleus
20	DNA binding DNA topoisomerase (ATP-hydrolyzing)	nucleus
21	skeletal development DNA metabolism signal transduction cell proliferation	-
22	-	-
23	chromatin binding transcription co-activator nucleosome disassembly transcription	plasma membrane nucleus nuclear chromosome
24	Cell structure Cytoskeletal Epidermal Development and Maintenance	cytoplasm
25	structural protein vision cell shape and cell size control intermediate filament	cytoplasm
26	cell shape and cell size control Cell structure	cytoplasm
53	leucine-zipper containing fusion	-

Table 3 (continued)

DNA SEQ ID NO:	Gene function	Subcellular localization
54		
55	Tumor endothelial marker 7 precursor, may be involved in angiogenesis	-
56	Aiolos; DNA binding protein that may be a transcription factor; has strong similarity to murine Znf133, contains zinc finger domain	-
57	The WASP-binding protein WIRE has a role in the regulation of the actin filament system downstream of the platelet-derived growth factor receptor	-
58		-
59		-
60		-
67		-
68		-
69	Midbrain dopaminergic neurons play a critical role in multiple brain functions, and abnormal signaling through dopaminergic pathways has been implicated in several major neurologic and psychiatric disorders. One well-studied target for the actions of dopamine is DARPP32.	-
70		-
71		-
72		-
73		-
74	Brain-specific guanine nucleotide exchange factor; activates the ERK/MAP kinase cascade plus R-Ras and H-ras; activates targets through a Ca2+- and diacylglycerol-sensitive mechanism; active protein associates with membranes	-
75	C-terminal tensin-like Phosphotyrosine-binding domain, phosphotyrosine-interaction (PTI) domain	-

Table 4

DNA SEQ ID NO:	Protein SEQ ID NO:	Gene Name	DBSN ^{PI} D	Type	Codon	AA-Seq
9	34	ERBB2	rs2230698	coding-synon	TCA TCG	S S
9	34	ERBB2	rs2230700	noncoding		
9	34	ERBB2	rs1058808	coding-nonsynon	CCC GCC	P A
9	34	ERBB2	rs1801200	noncoding		
9	34	ERBB2	rs903506	noncoding		
9	34	ERBB2	rs2313170	noncoding		
9	34	ERBB2	rs1136201	coding-nonsynon	ATC GTC	I V
9	34	ERBB2	rs2934968	noncoding		
9	34	ERBB2	rs2172826	noncoding		
9	34	ERBB2	rs1810132	coding-nonsynon	ATC GTC	I V
9	34	ERBB2	rs1801201	noncoding		
14	39	c-erbA-1	rs2230702	coding-synon	TCC TCT	S S
14	39	c-erbA-1	rs2230701	coding-synon	GCC GCT	A A
14	39	c-erbA-1	rs1126503	coding-nonsynon	ACC AGC	T S
14	39	c-erbA-1	rs3471	noncoding		
19	44	TOP2A	rs13695	noncoding		
19	44	TOP2A	rs471692	noncoding		
19	44	TOP2A	rs558068	noncoding		
19	44	TOP2A	rs1064288	noncoding		
19	44	TOP2A	rs1061692	coding-synon	GGA GGG	G G
19	44	TOP2A	rs520630	noncoding		
19	44	TOP2A	rs782774	coding-nonsynon	AAT ATT ATT TTT	N I I F
19	44	TOP2A	rs565121	noncoding		
19	44	TOP2A	rs2586112	noncoding		
19	44	TOP2A	rs532299	coding-nonsynon	TTT GTT	F V
19	44	TOP2A	rs2732786	noncoding		
19	44	TOP2A	rs1804539	noncoding		
19	44	TOP2A	rs1804538	noncoding		
19	44	TOP2A	rs1804537	noncoding		
19	44	TOP2A	rs1141364	coding-synon	AAA AAG	K K
23	48	KRT10	rs12231	noncoding		
23	48	KRT10	rs1132259	coding-nonsynon	CAT CGT	H R
23	48	KRT10	rs1132257	coding-synon	CTG TTG	L L
23	48	KRT10	rs1132256	coding-synon	GCC GCT	A A
23	48	KRT10	rs1132255	coding-synon	CTG TTG	L L
23	48	KRT10	rs1132254	coding-synon	GGC GGT	G G
23	48	KRT10	rs1132252	coding-synon	TTC TTT	F F

Table 4 (continued)

DNA SEQ ID NO:	Protein SEQ ID NO:	Gene Name	dbSNP ID	Type	Codon	AA-Seq
23	48	KRT10	rs1132268	coding-nonsynon	CAG GAG	Q E
23	48	KRT10	rs1132258	coding-nonsynon	CGG TGG	R W

Table 5

PRIMER	SEQUENCE
CACNB1	FAM 5' CCATATATAAACCCTCTCTGCTCTTTGGCT 3'TAMRA
CACNB1 FOR	5' CCCCACATCTGCTGCTCTATATTGTC 3'
CACNB1 REV	5' TGCCTACGCTGACGACTATGTG 3'
CDC6	FAM 5' TTTGGTTTCTACAACCTGTGCTAT 3'TAMRA
CDC6 FOR	5' GGGCTCCACACACCAAGT 3'
CDC6 REV	5' AGCTCTGAGCACCTCTACA 3'
EBI1-1	FAM 5' TGTCACAGGACTGAACCTCTCTCATGT 3'TAMRA
EBI1-1 FOR	5' CCCAAGGCCACGAGCTT 3'
EBI1-1 REV	5' TGTTCCTCTTAACGAATCGAAA 3'
EBI1-2	FAM 5' CTGGTCAACAACTCTGTGACCCCTCC 3'TAMRA
EBI1-2 FOR	5' TGGTGAGGAAAGCGGACAT 3'
EBI1-2 REV	5' CTGGCTTGAGGACAGTGAAG 3'
GCSF	FAM 5' CCAAGCCCTCCCATCCCATGTAT 3'TAMRA
GCSF FOR	5' GAGGTGTCGTACCGGTTCTA 3'
GCSF REV	5' CGGTTCTGCTCTTCCTGTCT 3'
GRB7	FAM 5' CCAGACCCGCTTCACTGACCTGC 3'TAMRA
GRB7 FOR	5' CGCCTGTACTCAGCATGGA 3'
GRB7 REV	5' GCGGTTCACTGTTGGA 3'
HKA3	FAM 5' ACCCGAGGATCAACCAATCAT 3'TAMRA
HKA3 FOR	5' AGTTCTGCTCTGTGACACCAT 3'
HKA3 REV	5' TAGGCTCAGTCAAGCCCAAC 3'
MLN50	FAM 5' CCTCTGGGCTTGTCTCGG 3'TAMRA
MLN50 FOR	5' AAGCCCGAGTTCACTCTTTT 3'
MLN50 REV	5' CTGTGGTTCAAGTCAATGTTCA 3'
MLN64-1	FAM 5' TCTGCTGGCTCTCTCGGT 3'TAMRA
MLN64-1 FOR	5' GGGCTGGCACCTGACTT 3'

Table 5 (continued)

PRIMER	SEQUENCE	
MLN64-1REV	5' CCCAACAGGGTCCAGACT	3'
MLN64-2	FAM 5' CGGCGCATTTAGCGGCG	3'TAMRA
MLN64-2 FOR	5' CCCAAGGACTTCGTGAATG	3'
MLN64-2REV	5' GCGCATCCTGATGACAAGTA	3'
PPARBP	FAM 5' AGCACCAACTGTGAACCGGTACAATGGC	3'TAMRA
PPARBP FOR	5' GAGGAGGCTCTGCTTTGG	3'
PPARBP REV	5' TCACAACTAGCGGGTGAGGAG	3'
PSMD3	FAM 5' TGCAGAGGAACGGCGTGAGCG	3'TAMRA
PSMD3 FOR	5' TGAGGTTTCTCCCAATCGTA	3'
PSMD3 REV	5' CAGCTCAGGGAAGCTGCATC	3'
RAR	FAM 5' CCCCCACATGTTCCCAAGATGCT	3'TAMRA
RAR FOR	5' GGAGGCGTAAAGGTTACGT	3'
RAR REV	5' TGATGTTGCGAGGTCAGTAA	3'
RPL23A	FAM 5' CTCCTGCGCCTCCTAAAGCTGAAGCC	3'TAMRA
RPL23A FOR	5' GGACGCGTGGCTTTTC	3'
RPL23A REV	5' TGTGCTGTGGACACCTTTC	3'
RPL19	FAM 5' CCACAGGCTGAAGGCAGACAAGGCC	3'TAMRA
RPL19 FOR	5' GCGGATCTCATGGACACA	3'
RPL19 REV	5' GGTCCAGCAGGAGCTTTTG	3'
NEUROD2	FAM 5' ACCACCTTGGCAGGTTGTCCAG	3'TAMRA
NEUROD2 FOR	5' CGCATGCACGACTGAAC	3'
NEUROD2 REV	5' GTCTCGATCTTGACAGCTTCTG	3'
TELE TELETHONIN	FAM 5' ACACCTGCCACGCGCCGAGG	3'TAMRA
TELE TELETHONIN FOR	5' CTGGGCAGAAATGAAGGATCT	3'
TELE TELETHONIN REV	5' GGGACTCTAGCAGACCCACACT	3'
PENT PNMT	FAM 5' CACCCACCTGGATTCCTGTTTC	3'TAMRA
PENT PNMT FOR	5' CCTTCAGACAGGCGTAGATGATG	3'
PENT PNMT REV	5' GGGTATTATTCTTTTATTAGTGCCACTT	3'

Table 5 (continued)

PRIMER	SEQUENCE	
HER2/NEU;ERBB2	FAM 5' TCCCTAAGGCTTTTCAGTACCAGGATCTG	3'TAMRA
HER2/NEU;ERBB FOR	5' CGAGCTTGGCCCTTTCCT	3'
HER2/NEU;ERBB REV	5' GAATGGGTGCGTTTGTCTTAG	3'
KIA0130	FAM 5' TCACGACCTCAGCTGCCCT	3'TAMRA
KIA0130 FOR	5' TGGTGAAGGTTCAGCCATGT	3'
KIA0130 REV	5' TCAGAGTCGCCAATGGCTTT	3'
THRA	FAM 5' ACCTCTGCCAGCTCCC	3'TAMRA
THRA FOR	5' GGCAATCTTACTTGTCTTTGA	3'
THRA REV	5' CCAGGAAGCACAGCAACTATTC	3'
MLN51	FAM 5' TCCTCCCTATCCATGGCCTAACCACTTC	3'TAMRA
MLN51 FOR	5' TGGGCAAGGGCTCCTATCT	3'
MLN51 REV	5' GTTACCCCTGGCAGACGTATG	3'
TOP2A	FAM 5' TGCTCTGATCTGAATCTCCCAAGAGAGA	3'TAMRA
TOP2A FOR	5' GAGTAGTTATGTGATTATTCAGCTCTTGAC	3'
TOP2A REV	5' TCAATGTTGTCGCCGAGTCT	3'
KRT10	FAM 5' CAGAAATCGGAAGACAGAACTATTGTCATGCCCT	3'TAMRA
KRT10 FOR	5' GATTAGTAAACCATAGCAGTTGAAGGT	3'
KRT10 REV	5' ATTACTGACGGTGGCTCGAACATAC	3'
K12 KRT12	FAM 5' TGACAGATCCAAATCAAGCAAGCAGTCAAC	3'TAMRA
K12 KRT12 FOR	5' TGATGTTTGGAGGAAGTTTATT	3'
K12 KRT12 REV	5' TTTGGTGGGTCTTTAGAGGAATC	3'
NR1D1	FAM 5' TGCCAAACCATGATCAGGTAGCCC	3'TAMRA
NR1D1 FOR	5' CAGCTCACCTGGCACTCA	3'
NR1D1 REV	5' CCTGATTTCCCGCAGTGT	3'
HSERBT1	FAM 5' CGCGGCTCCCGTTCTGCT	3'TAMRA
HSERBT FOR	5' TGGCCAAGCGTAAGCTGATT	3'
HSERBT REV	5' GCTGCGATGATCGGATCATCT	3'
MLL16	FAM 5' CACCATGGAGCCCATCGTGC	3'TAMRA
MLL16 FOR	5' ATCCCGAGGTGCAATTTG	3'

Table 5 (continued)

PRIMER	SEQUENCE	
MLL176 REV	5' AGCGATCATGAGCGACGTACT	3'
ZNF144	FAM 5' CTTGCAGAGATAGGAGCCGACAGCT	3' TAMRA
ZNF144 FOR	5' ATCCCTCGAGCCTTTTCA	3'
ZNF144 REV	5' CAGCCTCTGGTCCCACT	3'
PIP5K2B	FAM 5' TGATCATCAATCCAACTCTCCGGA	3' TAMRA
PIP5K2B FOR	5' CCCATGGTGTCCGAAC	3'
PIP5K2B REV	5' TGCAGGAGCCTCCATACC	3'
TEM7	FAM 5' CAGCCTTCTAAACACAATGATTCATGT	3' TAMRA
TEM7 FOR	5' CCTGAACCTTAATGGTAGAATTCAGATC	3'
TEM7 REV	5' TATTAACACTGAGAATCCATCAGAGA	3'
ZNFN1A3	FAM 5' TATCTGTCTCAGGATGCTCTATGATTCAGC	3' TAMRA
ZNFN1A3 FOR	5' CACAGAGCCTCTGAAATG	3'
ZNFN1A3 REV	5' GCGAGGTCATGCTTTTAGAA	3'
WIRE	FAM 5' CTGTGATCCGAATGGTGCAG	3' TAMRA
WIRE FOR	5' CCGTCTCCATCCAACT	3'
WIRE REV	5' ACCCATGCATTGGTATGGT	3'
PSMB3	FAM 5' AGTGGCCCTGGCCGAACAA	3' TAMRA
PSMB3 FOR	5' CCCATGGTGAATGATGACTT	3'
PSMB3 REV	5' CCAGAGGACTCACATTC	3'
MGC9753	FAM 5' CCAGAACTTTCATCCCAAGGAGTCT	3' TAMRA
MGC9753 FOR	5' CTGCCCCACAGGATAGATG	3'
MGC9753 REV	5' AAAAATCCAGTCTGTTCAACA	3'
ORMDL3	FAM 5' AGTGCACAGCTCCACGGA	3' TAMRA
ORMDL3 FOR	5' TCCCTGATGAGCGTGTATC	3'
ORMDL3 REV	5' TCTCAGTACTTATTGATCCAAAATCC	3'
MGC15482	FAM 5' TCCAGTGAAGCAACCCAGTGTTC	3' TAMRA
MGC15482 FOR	5' CACTTCTAGAGTACCGTGGATCT	3'
MGC15482 REV	5' CCCTCACTTTGTAACCCCTTGT	3'
PPP1R1B	FAM 5' CAGCGTGGCCCAACCAACA	3' TAMRA

Table 5 (continued)

PRIMER	SEQUENCE	
PP1R1B FOR	5' GGGATTGTTTTCGCCACACATA	3'
PP1R1B REV	5' CCGATGTTAAGGCCATAGC	3'
MGC14832	FAM 5' TAAATGTCCGGCCACATGAGTTCCC	3'TAMRA
MGC14832 FOR	5' CGCAGTCCTGGCACAT	3'
MGC14832 REV	5' GACACCCCTGACCTATGGA	3'
LOC51242	FAM 5' CAGTGACCTCTCCGTTCCCTTGGA	3'TAMRA
LOC51242 FOR	5' TGGTCCCTGTGCTCTTC	3'
LOC51242 REV	5' AGGTCAGGAGGAGAAAAAC	3'
FLJ20291	FAM 5' CCAGTCCACCCGTTAAGAGATCAA	3'TAMRA
FLJ20291 FOR	5' TTGTGGGACACTCAGTAACTTTGG	3'
FLJ20291 REV	5' ACAAGCACTCCACCCGAGAT	3'
PRO3521	FAM 5' AGTCTGTCTCACTGCCATCGCCA	3'TAMRA
PRO3521 FOR	5' AAGCCTCTGGGTTTTCCTTT	3'
PRO3521 REV	5' CCCACTGGTGACAGATGGT	3'
Link-GEFII	FAM 5' CATCTGACATCTTCCCGTGGAG	3'TAMRA
Link-GEFII FOR	5' CTTTGCACGATGTCACCA	3'
Link-GEFII REV	5' TTTCCCGTGGAGCAGGAA	3'
CTEN	FAM 5' CCGCGGCTAATATGCAACATTAGG	3'TAMRA
CTEN FOR	5' CGAGTATCCAAAGTGGTATCG	3'
CTEN REV	5' ATCACAGAGATGGCCCTTATCT	3'

Table 6

No.	ID	forward	reverse	PCR size (bp)	GB ID
1	D17S946	ACAGTCTATCAAGCGAGAAATCCT	TGCGTGCACAGAGA	128-142	Z24029
2	D17S1181	GACAAACAGACGAGACTCC	GCCAGCCTGTGCATTATTC	122	-
3	D17S2026	TGCTCATTTGACACAGAA	CAGCATGTGATGCAATCC	171-318	G05498 X53777
4	D17S838	CTCCAGATCCAGACATGA	AGCAGATCTGAGCCCTTC	71-103	Z51080
5	D17S230	GGAGAAATCAATAGACAAAT	GCTGCCATATATATATTTAAACC	151	-
6	D17S1818	CATAGGTATGTTCCAGAAATGTGA	TGCTACTCGAAACACAGA	119-151	Z52895
7	D17S614	AAAGGGAAGGGCTTCTCAAGCT	NGGAGTTTCAGCTGAGCAAGAT	136	L29873
8	D17S2019	CAAAAGCTTATGATGCTCAACC	TTGTTTCCCTTTGACTTTCTGTA	151-152	G07286 Z39013
9	D17S608	TAGGTTCACTCTCATTTTCTTCAG	GTCTGGTCTTTTATGNGCTTGTG	136	L29870
10	D17S1655	CGGACGAGAGTGTTCATGG	GCATACAGACCCCTCTACCT	240	-
11	D17S2147	AGGGGAGATAAATAAATCTGTGG	CAGGATGAGACACTCTCCATG	138	G15195
12	D17S754	TGGATTCACTGACTAGCCTGC	GCCTGTCTCTCCATGTGTGC	145	-
13	D17S1814	TCCCATGACGGTGTG	CTGGAGTTGGCTTTGTGGAT	150-166	Z52854
14	D17S2007	GGTCCACGAAATTTGCTG	CCACCCAGAAAACACAGAGA	102-103	G07073 X03438
15	D17S1246	TCGATCTCTGACCTTGTGA	TTGTCAACCCATTGCTCTTC	115	-
16	D17S1979	CCTTGTGATAGATTCAGTCCC	CTTGTCCCTTCTCAATCCCTCC	199	G11172 X55068
17	D17S1984	TTAAGCAAGGTTTAAATTAAGTGC	GATTACAGTCTCCCTCTCCC	134	G14779 T50487
18	D17S1984	GGTTTAAATTAAGCTCGATGGC	GATTACAGTCTCCCTCTCCC	126	G11580 T50487
19	D17S1867	AGTTTGACCTGAGGGCTTTG	TTTAGACTTGGTAACTGGCG	94	Z51301
20	D17S1788	TGCAGATGCCCTAAGAACTTTTCAG	GCCATGATCTCCCAAGCG	156-168	Z52160
21	D17S1836	TCGAGGTATGTTGAGCC	AAACTGTGTGTCTCAAGGATACT	167-173	Z53182
22	D17S1787	GCTGATCTGAAGCCATGA	TACATGAAGCCATGGTCTG	239-251	Z52130
23	D17S1660	CTAATATAATCTCGGCACATGG	GCTCGGACCAACACAGAT	201	G06069
24	D17S2154	GATAAAAACAGACCTGGCTCC	CCACCGCTTTTCTGATCTA	137	G15440
25	D17S1953	TGTAAATGAAGCCCATGAGG	CACCTCAACTCAACAGCTCAAGGTG	180	G11900
26	D17S2098	GTGATTCAGCATATGTAATATCC	ATTTCAGCTCAGTTCACCTGCTTC	181	G13994
27	D17S518	GATCCGTGGAGACTCAGAG	TAGTCTCTGGACACCCAGA	88 - 100	X60690
28	D17S1851	ATTCTGAGTGTCTACCTGTTGAG	ACTGACTGGCCACTGC	237 - 253	Z53675
29	D17S4358	TCGAGAGGACAAAATCAACC	GAAACGGTTAGTCCATTGG	58	-

Table 6 (continued)

No.	ID	forward	reverse	PCR size (bp)	GB ID
30	D17S964	GTTCTTTCCCTTTGGGG	AGTCAGCTGAGATTGTGCC	224	L36695
31	D19S1091	CAAGCCAAAGACATCCAGTT	CCCCACACAGCTCATATG	238	G14589
32	D17S1179	TTTCTCTCATCTCCATTGGG	GCAACAGGGAGACTCCAA	113 - 125	-
33	D10S2160	TCCATCCCGTAGACCTC	TATGGAGTACCTACTCTATGCCAGG	349	G06592
34	D17S1230	ATTCAAAGCTGGATCCCTTT	AGCTGTGACAAATGCCTGTA	108	L32949
35	D17S1338	TCACCTGAGATTGGGAGACC	AAGATGGGGCAGGAATGG	178 - 200	-
36	D17S2011	TCACCTGCTCCCAAGCCAG	AAACACCACTCTCCCTCTG	115	G07143
37	D17S1237	TTCTTGGGCTTCCCGTAGCC	GGGGCAGACGACTTCTCCTT	186	L32947
38	D17S2038	GGGATACACCTTTAAAGTTCC	ATTACCTTAATGAGGATTTCTTTT	228	G6219
39	D17S2091	GCTGAATAGCCATCTTGAGCTAC	TCCGCATCTCTTTTAAAGGGCAC	157	G13941
40	D17S649	CTTTTCACTCTTTTCAGCTGAAGAGG	TGACGTGCTATTTCCTGTTTGTCT	146	L36685
41	D17S1190	GTTTGTGTGTATGCGCTGC	CAACACACTACCCCAAGGA	122	L18197
42	M87506	ACTCCTCATCTGTAGGGTCT	GAGTCGGCTACCTGAGTGCT	102 - 120	m87506

SEQUENCE LISTING

<110> Bayer AG

<120> METHODS AND COMPOSITIONS FOR THE PREDICTION, DIAGNOSIS, PROGNOSIS, PREVENTION AND TREATMENT OF MALIGNANT NEOPLASIA

<130> LeA 36108.1 EP

<150> EP02010291.9

<151> 2002-05-21

<160> 314

<170> PatentIn version 3.1

<210> 1

<211> 3846

<212> DNA

<213> Homo sapiens

<400> 1
gctcccgccg agctogccct ggggaacagg aogcgctga gctcaggcgt ccccgcccca 60
gcttttctcg gaacctgaa ccccaactgc gccgggtgog gcaagatogt gtatcccaag 120
gagagggtag actgtctgga taagtctctg cataaagcat gcttccattg cgagacctgc 180
aagatgacac tgaacatgaa gaactacaag ggctacgaga agaagcccta ctgcaacgca 240
cactacccca agcagctcct caocatgggt goggacaacc oggaaacctc tcgcctcaag 300
cagacagtg agctccagag tcagggtgogc tacaaggagg agtttgagaa gaacaagggc 360
aaaggtttca gogtagtgcc agacaagccc gagctocaga gaatcaagaa gaacagggac 420
cagatocagta atataaata ccattgaggag ttgagaaga googcatggg cctagcggg 480
ggcgagggca tggagccaga gogtogggat tcaaggagc gcagcagcta ccggcgccgc 540
ctggagcagc agcagcccca ccacatcccg aocagtgccc oggtttaoca gacgcccag 600
cagcagccgg tggccagcto ctatggtggc tacaaggagc ctgcagcccc agtctccata 660
cagcagccgg ccccaagggt tggggggaag oggtaccggc oggtgtatga ctacagccc 720
gocgacgagg acgaggtctc ctccaggac ggggacacca togtcaacgt gcacagatc 780
gaogacggct ggaatgtagc gacggtggag ogcaccggcg acaoggggat gctgcgggc 840
aaotacgtgg aggcacatc aaccgggag gcccccatct gtcttcagca cttccacgg 900
catgcatacc gtctgtggcg tgagccgtcc attcttcaagt gtctctgttt tttaaaaact 960

	gogacagact	gtgattctcta	ccocctcttcc	agcttctttt	gccaactgaa	goccttcttct	1020
	gocactctctg	ogggctccct	ccctctggcag	gcttcccccg	tgatogactt	ctctggtttct	1080
	ctctctggatg	gaacgggtat	gggocctctct	gggggaggca	gggctggaa	gggagacctg	1140
5	ttggctctgtg	ggcctcaacct	gocctctctgt	ctctccccc	caatccctcc	tgccagctc	1200
	ctccacatacc	caacacattcc	agggctctggg	tgagcctgac	tgccaggagcc	ccaggtccagg	1260
	ggctccctac	attccccaga	gtgggatacca	cttcttggtt	ctctgggatgg	cgatgggggct	1320
	ctctgcccgtg	gttgaggagac	agtgggatggt	gctctacctc	ctcttctccaa	agaggggggct	1380
	ctgcccacact	ggggctctctc	ctccctacctc	ccctctcagg	ggcaacacaca	ggagaattggg	1440
10	gttctctgctg	tggggcgaat	tcactccctc	cccgoggtt	ctcttcgaca	ctgtgatttt	1500
	gocctctctgc	ccacgcagac	ctgcagcggg	caaagagctc	ccaggagagc	acagcttggg	1560
	ctagggttctt	goccttctcta	atcttgaggga	cagctacccg	aaggaggagga	acagggaatt	1620
	ctctctccgca	gcccccttcc	ccacgcacac	ccccagctct	caggggaccc	tgccctgctc	1680
	ctaggctcga	agccatggctc	ccgaagtgtta	gggcaagggt	gocctcaggac	cttttggtct	1740
15	tcagcctccc	tcagccccca	ggatctgggt	taggtggcgg	ctcctccctg	ctcctcatgg	1800
	agctgcacaa	agtcagcagg	catgacctcc	ccctccacgc	ccaatgocaa	gtggactgg	1860
	catgtatgtc	caggagccttc	gacacactaaa	ctctccaaag	ctgtgtgtcg	gaggcaggag	1920
	gggggttgtgt	ctgtagggtgt	tgacacgcga	gtgtgttgagt	gtgtgttgagt	gaggttgggtg	1980
	agagctgtct	tcocctcgaga	ctctgggocct	gtgtgttggt	ggggttatgt	gaggttatga	2040
20	gcagagaagt	ctcttaggtgt	gtttctccag	aaocccacgt	gagggggag	ggctcctggg	2100
	ggaggaagcc	caaccaatctg	ttctttggaa	tgaattctct	ccctccccc	atctctaggt	2160
	tggaagtggg	ggctctgcagc	ccctttgga	tggtctcagg	tggaaggttaa	gaggttgggtg	2220
	tgctacactca	ggcatatgtt	tcoccatctc	tgctctgggg	agacccctcc	aggaaggtcc	2280
	gtgtccacct	gtgggtgtct	ccctcgggg	ctctccccc	ccactctaca	ccactctaca	2340
25	taagctccct	ttgaagcaag	aaaggggttc	ccagggtctc	aaaactggaa	gcacagctcc	2400
	gggagtgggg	agggaaagac	gggtctatat	ccagttccct	ctctctgtct	atgggttggtg	2460
	gtgcaaacct	tgccctcact	tgattcatct	ctggtttct	tgccacccct	tgggagtccc	2520
	catcccaatt	tcactcctgag	cccaaacagg	ccctgcact	ggcctcttgt	cccttgagac	2580
	acottgtacc	acaggttgagg	ggcaggacct	gaaggtattg	gocctgttca	caatcagtc	2640
30	tcattgggtgt	ttttgtccaa	tgcttgttaa	ttgatttggg	gatgtttgoc	ccgaattgaga	2700
	gggtgaggaa	aagactgtgg	gtggggaggc	ccctgctgac	ccatcccttt	tcctttctgg	2760
	ccccagccca	gggtggaggca	agtggaatat	cttatattgg	ccatcccttt	tcctttctgg	2820
	ggcagagaa	ctcttgggg	cttgggttgg	ccctgctgac	ccatcccttt	tcctttctgg	2880
	aaagoccaat	gtggatttgg	ggacccaaag	ccctgctgac	ccatcccttt	tcctttctgg	2940
35	tttgggagag	tggtgagtg	aagggaggag	ccagggacac	atccccctag	aggaactgag	3000
	taattctctc	tcocagataa	ggcagccag	ccagggccgt	cagcctgttt	tcactcagct	3060
	actcttctct	tcctaaaaat	acggggggct	ccaggaagac	tgctgtcagg	ccctccctct	3120
	gtgcacactg	agagcacaga	gocctgagg	ccctgctgac	cccccaagaga	gagggaggac	3180
	catccctaga	ttctccacag	ctcttccag	ccctgctgac	tgcttcttcc	ccaaaacatc	3240
40	ttctctacag	ttccacagag	ctcttccag	ccctgctgac	tgcttcttcc	ccaaaacatc	3300
	tagaatactg	ataatacttt	tgctggccaa	taactagaat	cagggaagag	gcataaagc	3360
	ggaaagtggg	ttctccacag	actcctcctc	gttttctccc	cccaagagag	tgctggttga	3420
	aatgactctc	ttgcttaatt	ctctgcttcc	ccctccacac	ctccagctgac	ctccagggaa	3480
	ttttatagctg	gaaaaaacaa	aatccacccc	tacaaaactc	gtttcccttt	tgccgctttt	3540
	tgacacagtg	aaatttgctc	ctgtccaaac	attctcatcg	tatttaaaaa	gaaaacagaa	3600
45	agtggttgaa	gcggccagct	catcttttta	tattctatcg	tggtgtatgt	tatgtgtgtg	3660
	ttggtccccc	ccctctgggg	ctgtgtctct	tattctatcg	ttgtctcact	ttggtctgtt	3720
	tgacacattg	acttgaaacca	caagtgaaac	ggatccaaac	ttctctcctg	gttatgattc	3780
	ttttta				tgactcaaat	aaaagtataa	3840

<210> 2

<211> 1711

<212> DNA

<213> Homo sapiens

5	<400> 2						
	gagggaaggg	aggaagagg	cagccgaagg	cogagctggg	tggtctggacc	gggtgctggc	60
	tgccgcggcgt	gctttcggcgt	cccacggcct	ctccatcgcg	ctgaggggagc	cggcgtcgcg	120
	gcccgggggg	ggagggggagg	ctcctctcca	tggtccagaa	gaccagcatg	tcggcggggcc	180
10	cttaaccacc	ctccaggagg	atccccatgg	aggtctttoga	ccccagcccg	cagggcaaat	240
	acagcaagag	gaaaggggga	ttcaaacggt	caga tggggag	caogtctocg	gataccacat	300
	ccaacagcctt	tgtoocgcaag	ggctcagcggt	agtcctacac	cagccgtcca	tcagactctg	360
	atgtatctct	ggagagggac	cgggaagcct	taaggagga	agcagagcg	cagggcattag	420
	cgcagctcga	gaaggccaa	acaaagccag	tgccatttgc	tgctgggaca	aatgttggct	480
	acaatccgtc	tcacggggat	gaggtgcctg	tgccagggagt	ggccatccac	ttcgagccca	540
	aagacttct	gcacatcaag	gagaaataca	ataatgactg	gtggatcggg	cggtctggta	600
15	aggagggct	tgaggttggc	ttcatccca	gcccctcaa	actggacagc	cttcgctcgc	660
	tgccaggaca	gaagctgcgc	cagaaacggc	tcggctccag	caaatcaggc	gataactcca	720
	gttccagctc	gggagatgtg	gtgactggca	cccgcgcgc	cacacccct	gccagtcca	780
	aacagagca	gaagtgcaca	gagcatgtgc	cccctatga	cgtggtgcct	tcacatgagg	840
	ccatctact	ggtgggacgg	tcgctcaagg	gctacagagt	tacagacatg	atgcagaaag	900
20	ctttatttga	cttccttgaag	catcggtttg	atggcaggat	ctccatccat	cgtgtgacgg	960
	cagatatctc	ccctggctaa	cgctcagttc	tcacacaccc	cagcaaacac	atccatcttg	1020
	agcgtccaa	gaacccctcc	agcctggctg	aggtgcagag	tgaaatcgag	cgaactcttc	1080
	agctggccgc	gaacccctcc	tggtgcctc	tgga tgcgt	ccacatccat	cacccagccc	1140
	agctgtccaa	gaacccctcc	tggtgcctc	tgga tgcgt	ccacatccat	cacccagccc	1200
25	tacttcaagg	gctccatcaag	tcocgagga	accttcagtc	caaacacccc	aatgtcccaa	1260
	tagcgccctg	ggaaagctg	gcacagtgcc	cccttgaaat	gtttgacatc	atctggagtg	1320
	agaaacattt	ggagga tgc	tgccagcatc	tgccggagta	cttggaagcc	tattggaggg	1380
	ccacacaccc	ccacatcgca	acgcacacca	atccgctgct	gaacccgccc	gatgtatccg	1440
	cagccctgag	cctgagccct	gcccctgtct	ccaaactcca	ggtacaggtg	ctacactcgc	1500
	tcaggagaaa	ctcggccttc	tggggctggc	tgga tgcct	acagcggggc	agtggtgtgc	1560
30	ccacaggaca	ggaacatgcc	atgtagtggg	cgccctgcgc	gtcttccctc	ctgctctggg	1620
	gtcgggaactg	gagtcgagg	aacatggagg	aggaaggga	gagctttatt	ttgtaaaaa	1680
	ataagatgag	cggcaaaaa	aaaaaaaaa	a			1711
<210> 3							
35	<211> 698						
	<212> DNA						
40	<213> Homo sapiens						
45	<400> 3						
	ttttctcttc	gctgctggcg	cgcagccat	gagtatgctc	aggcttcaga	agaggtctgc	60
	ctctagtgtc	ctccgctgtg	gcaagaagaa	ggtctggtta	gaccccaatg	agaccaatga	120
	aatgcacat	gccaactccc	gtccagagat	cogggaagctc	atccaaagatg	ggctgataat	180
	cgcgaagcct	gtgacggtcc	attcccgggc	togattgcgg	aaaaacacct	tgccocggcg	240
	gaagggcagg	ccacatggca	taggttaagc	gaaggttaca	gccaatgccc	gattgccaga	300
	gaaggtcaca	tggtatgagg	gaatgaggat	tttgccggcg	ctgctcagaa	gataccttga	360
50	atctaagaa	atcgatgcgc	acatgtatca	cagcctgtac	ctgaaggtga	aggggaatgt	420
	gttcaaaaac	aagcggatct	tcattggaac	catccacaag	ctgaagggag	acaagggccc	480
	caagaagctc	ctggctgacc	agcctgaggc	cgcgaaggtct	aagacacaga	aagcagcaga	540
	cgcgcgtgaa	gagcgcctcc	agcccaagaa	ggaggaagat	atccagactt	tatccagaga	600
	ggaagagacc	aagaaataaa	acctccacct	ttgtctgtac	atactggcct	ctgtgattac	660
55	atagatcagc	cattaaaaa	aaacaagcct	taattctgc			698

<210> 4

<211> 5810

<212> DNA

<213> Homo sapiens

<400> 4

gggaagatgg	cgccgggctc	gagcacccct	ctctctcttc	cgccggggac	ttoagattga	60
tocttccggg	gaagagtagg	gaactgtggg	gcootgcgtc	cgggagatcc	gagccaaact	120
gttttccggg	ttagtggtgg	ggaagggtct	atccttttgt	ggcggatcta	gcttctcttc	180
gcoctcagga	tgaaagctca	gggggggaaa	cgaggagctc	gaaaagctga	gtaagatgag	240
ttctctctcg	gaaoggtctc	atgcacaaat	taacacaaat	agacccctga	gtgaaaacct	300
taagcttggt	cgctcaagtc	tgagagaagag	ggttggtgat	agttctggag	ggcatcaaca	360
tttggctcgg	tgcttggaga	caatgcagaa	ggctctcaaa	gtaacatctt	taccagcaat	420
gaactgtcgt	ttggagtcca	tagcaggaca	gaatggactg	ggctctcttc	tcagtgcacg	480
ttggacatgaa	tgctacatca	cgctcagatc	gttctatgtg	gaagtgcagt	tagatctctg	540
aggacagcttt	ttgtgatgtaa	aagtggctca	ccatggggag	aatcctgtga	gctgtccagg	600
gcttgctacag	cagctaaagg	aaaaaaatct	tgaatgaatt	tcataagcaac	ttaaaggcct	660
tgcttaacttc	tataaccttc	caggggacaa	caaaactgaag	actaaaatgt	acttggctct	720
ccaatctctta	gaacaagatc	ttctctaaat	ggcaattatg	tactggaaag	caactaatgc	780
tggtccctctg	gataagatct	ttcatgggaag	tgctggctat	ctcacaccaa	ggagtggggg	840
ctatttaactg	aaactgaagt	actatgtctc	tcctctctgc	ctactggatg	acaagactgc	900
atctccatct	atcttgcagt	agaataatgt	ttctcgtatc	ttgggcatga	atgcactcgt	960
gacaaactgaa	ggaacatact	ctgtgtacaa	actcccaatt	gcaacataaa	ttatgggggtc	1020
acatccatgtg	gacaaataat	ggaccccttc	ctctctctca	atcaacagtg	ccaacagctg	1080
tgatctctctc	gctgtgtctc	ctctgaaatt	tcctcagcca	atcaacagta	ctagagcaatt	1140
tgcttoagaaa	ctcaggaact	gcacaggaat	tcacttgctt	gcaactcaac	caacttatgc	1200
acccctgtat	gaactgtaca	ctcagtttga	gctatcaaa	gacccatgac	ccaatccctt	1260
gaatcaccaac	atgagatctt	atgctgctct	tcctggctga	cagccactgc	atttctctca	1320
caaggatgctc	ctctctccag	attggcgaag	tcctcagggg	accctgttta	gcaaaaatca	1380
ctttcagcaac	attggaaagt	ttctctctat	ccctaaatct	aaagaagatt	ctcctggggt	1440
taacacccctc	attggaaagt	gtgtccaaag	aactattctg	agcgtatctt	ttcagcacc	1500
ctctccatctt	gaagtgtgct	ctctctcaga	gtctcgtttc	agcgtatctt	ttcagcacc	1560
tgctgaatgac	tcctctgtgt	gtgtggtaat	ggatgtgaag	ggcttaacac	atgtgagctg	1620
taaaactctac	aaaggggtgt	cggtatgcact	gatctgcaca	gatgacttca	ttgcacaagt	1680
tgctcaaga	tgctatgtca	tcctctgtac	gatgagggct	attcgggaga	aagctgaaac	1740
oattcaagcc	gacacccagc	caactgtccct	catgtgcagag	acaattgaag	acatgggtga	1800
aaagaagcc	cccccggcta	gcagcccgag	gtatggcatg	acaaacaggc	acaaacccat	1860
gaagtgttaac	actacataaa	ccaaacacct	tcgggggggt	ccatgtgcac	ctctgtttaa	1920
tatgacacatg	agcaataaag	atcggcatag	gtcgtgtggc	catggggagg	acttcagcaa	1980
gggtctctcag	aaacccatct	ttcaacagtt	gttgcaaatc	acaggggaag	gggggtctac	2040
caatgtctctg	agctcagacc	ctctctctca	ccctctctct	ccatgtctct	ccatgtctct	2100
caacacccag	aaaccccgga	tgctcatgaa	ccctctctca	gataatctct	ccacggattt	2160
ctcaacccctt	tatggaagca	gcoctttaga	aaaggcagaac	tcctctctcc	gctcaccccg	2220
ctcgggaata	tgctcgggga	gcaacaagac	caagaaaag	aagctatcaa	gattacccac	2280
tgagaaaccca	aagccacaga	ctcagagtg	ctctcagagg	gagctattct	caatggatgt	2340
tgactcacaag	aacctatctc	ttgatgtcaa	ctcagacgtg	gacacgctgg	atacggcaac	2400
catcactcaca	gctccaaagc	agtgtagcac	tcacccacaa	acttacccac	aaacagtaac	2460
tcaccccaaa	ccaggtatgt	caagactatc	agaggaagct	aggtcagaca	catctggccc	2520
agatgttaact	gacatctctt	cagacatctg	tcagagatct	tcataaactc	ccatcacatg	2580
tgatgatttgc	caagcaattg	gcaacccctc	aactaaccaat	aatgaaaaat	ctacacatga	2640
gagtaacccct	tttgactctg	atgtctttca	aagccacagc	agtgactctc	ctacacatga	2700
tcacgctgagt	cttatgtcag	atgtcgtctg	tgatttatgt	aacagccaga	gocaaagtgg	2760
tttttttctat	gatggagtag	atttcaatcc	ccaaagtggg	gataatgatg	atttcaaaag	2820
ttttggagaaa	gaatattttg	atgaaagcag				2880

	atttgcatct	caggcactaa	atactttggg	ggtgccaatg	ottggagggtg	ataatggggg	2940
	gaccaagttt	aagggcgaata	accaagccga	cacagctgat	ttcagtat	ttccagtagc	3000
	cggaagaagt	ttagctctgt	cagatcttat	ggagcatcac	atgtgttagtc	agggctcttt	3060
5	actgacacat	ggggacttag	ggaaagaaaa	gactcaaaag	agggtaaaag	aaggcgaatg	3120
	caccagtaat	agtactctct	cggggcccgg	attagacagc	aaaccaggga	agggcagctg	3180
	gaccctctgt	aatgatggga	aaagcaaaaga	taagcctoca	aagcggaaag	gtgttgcac	3240
	tgaggggaag	ttctcatctc	atagttcttc	taacagacat	tttaccocac	ctaccagtac	3300
	aggttgatct	aaatggccga	gaagtgcagg	aaagtctcag	actccccacg	gtgttgcac	3360
	accacccatt	cccaaaatca	ctattccgat	ttctaaaggga	acagtgtatg	tggggcaagc	3420
10	ttctctctac	gttcagctata	ccagcagtg	ttctgtgtct	ttctcaggga	gcaaaagcca	3480
	ccatagccat	ttctctctct	cttctctctc	tgcttccacc	tcagggaaga	tgaaaggaag	3540
	taaatcgaat	ggttcaatca	gttcaagttt	aagttagcgt	atgtattcta	gccaaggtac	3600
	ttctggatct	agccagctca	aaaattctac	ccagtctggg	gggaagccag	gtctctctcc	3660
	cataaaccag	catggactga	gaagtggctc	tagcagcacc	aagatgaaac	ctcctctcag	3720
15	gccatcatca	cttatgaaat	cttttttaag	taaaccaaac	atatccocct	ctcatccaag	3780
	gccacctgga	ggctctgaca	agcttgcttc	ttcaatgaag	ctgtttctct	tgcttggaa	3840
	atctcttaaa	gccaaagtcc	ctatcagttc	aggttctgtg	gggttctcta	ttgttggaa	3900
	tagttcaagc	cttggaatga	agttctcttc	aggtttagga	ttctcaggtc	gtgttctcag	3960
	gaaaaactccc	ccatcatctc	atctctgtac	ggcattcttc	ttctctcttt	ctccaaagtg	4020
20	ctctctccatg	ttatctctct	agaaaccagc	tgaggatct	aagcatgggg	ttgtccaccg	4080
	aaacaagaag	ccgtctctga	cagctgtcat	ccagatgggg	gtgagccaa	attctctcag	4140
	tgccctctgg	ggtgaagacc	cactggcagg	agggagagag	tttcagggca	aggttgagaa	4200
	ccatctctatg	ttctccaaac	ataacatgtc	ctccgggagt	tcagttgatt	ctcttaagaa	4260
	aagtgtataa	gacaaatcaa	aggtttccac	aggtgtggca	aaaaattatc	tcagtaagca	4320
	gacctcagag	tcaaaaaatg	tggggagcac	agtgactttg	cagaaacctg	gggaagtag	4380
25	tgatggaggc	ttccctagca	ttaaagccaa	ttctaaaaac	tatggctctc	cactcatcag	4440
	tggaagaagg	cttagggctc	aaatggcttc	ttccagccat	agtaagtcc	cagcatatac	4500
	tggtttccat	ccaagcagtg	agcgtggctc	aggtctctcc	atagcagaga	catcttatac	4560
	ccccccagat	ctggcagctg	aaagttagtc	accacttcca	gaatacagca	cagagaacca	4620
	gaatagctcc	agctcagagc	atggatctcg	aaaagacaaa	gatagggaac	gagaccggga	4680
30	taagaagaac	aaaaagaaaa	agaagaaagt	catcaagcca	gagagtgtgt	ccaaatccac	4740
	caaaagccga	gaccagaaaa	aatctcatag	aagtaaacaca	atcttatctg	cagacagacc	4800
	catctctctc	gaccagctct	tgtctatgac	ggagggaagat	gatgatctta	tggaatgtgc	4860
	ctcaaggctc	agccagagct	ttatgatctg	aaaagaaaca	gggccagagg	aaaaaaaact	4920
	ctctgatggg	aatttagaac	cttattctct	ggggtagtgc	agacaggtct	gatttgggtta	4980
35	attgataagt	ttataggcaa	accaaccata	agctgggtga	attagaaagg	catctccaga	5040
	agaatccata	atggcatggc	tttgacatca	agctgggtga	gaagtctctc	ttgtctctgt	5100
	ccctattata	gaaaccacag	gggtttgatc	tggttaccag	gaagtctctc	ttgtctctgt	5160
	gcagagaaga	aagttaaaat	acttgcttaa	gaaagggagg	ggggtggagg	gggtgttagg	5220
	agaggggaag	gaggggaaca	gttttggggg	aaattatctc	atatatttct	ttctctctct	5280
	ttccattttt	agggcatggt	ttaaactcat	tttagtgcat	gtatatgaag	gggtggggag	5340
40	aaaaagaaa	agccatacat	ttcttgatgc	atttgataga	aggtgtgtca	acttgttttg	5400
	aggtagtgtg	ccgtttgagt	catggggcaa	tgaaggactt	tggtcaattt	gggcaactaa	5460
	tgatgtgtgt	gtgtgtgtgt	cttaggagtg	actggggagg	gggaagattt	tttagctatt	5520
	tatttgtaat	attttaaccc	tttatctgtt	tggttttata	cagtgttctg	ttctaaatct	5580
	atgaggttta	gggttcaaaa	tgatgggaag	cogaagacga	aggtttatat	gggtgttagg	5640
	agctttatagc	ttgtgctaat	actgtagcat	caagcccaag	caaattagtc	agagcccgcc	5700
45	tttagagtta	aatataatga	aaaaaccaaa	atgatatatt	tatttttaga	gggttttaaa	5760
	aggggttoaga	gatcatagga	atattaggag	ttactctctc	gtggagggtat		5810

<210> 5

<211> 5515

<212> DNA

<213> Homo sapiens

[illegible]

	cagctgaatc	aaagtgaatt	ggcagtgtaa	ttaaacctgc	tgacagagca	aaccgacctg	3420
	agcatccctc	aaatggcaca	gctgcttaac	atccactcca	accacagagt	gcagcagcag	3480
	ctggaaagcc	tgaaccaato	catcagtgcc	ctgaaggaa	ctactcccca	gcagcagcag	3540
5	ctgagagcca	tggccccaga	ggagttottt	aggaagcaca	ccctctgccc	gcagcagcag	3600
	ccctcagcag	aacagatgac	ccttgaagct	tcagagcaca	cagctgcacat	gcagcagcag	3660
	ttggcagttc	tcttgagtca	gctgatgaaa	acccaagagc	cagcagggag	tctggaggaa	3720
	aacaacagtg	acaagaacag	tgggccacag	ggggcccgaa	gaactccccc	aatgcacacg	3780
	gaggaggcag	cagcatgtcc	tcctcacatt	cttcccccag	agaaggagcc	ccctgagccc	3840
	cccgagacct	caccccgccc	acctccaccc	ccctctggtg	aaggcgatct	ctccagcgcc	3900
10	cccccagggt	tgaacccagc	cgtagacagc	gccttgctgc	aaccttttct	ccagcctgaa	3960
	gcagagcctc	ctggccacct	gccacatgag	caccagcctc	tgagaccaat	ggagtactcc	4020
	accgcccccc	gtccaaacag	gaacttatga	aacactgatg	ggcctgaaac	aggggttcagt	4080
	gcatttgaca	ctgatgaacg	aaactctggt	ccagccttga	cagaatcctt	ggctccagac	4140
	ctgggtgaaga	acaggacctt	ctcaggctct	ctgagccccc	ttggggagtc	cagcagttac	4200
15	cagggcacag	ggctcagtgca	gtttccaggg	gaccaggacc	tcogttttgc	cagggtccccc	4260
	ttagcgttac	acccggtggg	ogggccaccca	ttcctgaagg	ctgagggaag	cagcaattct	4320
	gtggatccat	cagagaccaa	attgcaaaac	tatggggagc	tggggcccag	aaccactggg	4380
	gcagcagagt	caggagcagc	ccctcactgg	ggggccccc	ctcagctctc	tgcttatgga	4440
	aaactctatc	gggggctcac	aagagtcoca	ccagaggggg	gaagggggag	agggagtctc	4500
20	tactcaaccca	gagactctcag	tgctcctgaa	gattccttcc	ctatccatcc	ttccatccag	4560
	ttctctgaa	ctttaatgaa	atcatttgcc	agagcggagt	aatcatctgc	atttgctcac	4620
	tgcagaagctg	tcctgtgtat	tcctgtgtca	cttgctctac	gcaggcgagc	gttggaataa	4680
	tgatgtttgc	acaggttccc	ccctggatgg	ctatagccag	aacatttact	taaacctcac	4740
	cttagtagat	acaagttagag	aatatggaga	ggatcatcac	attgaaaaag	aaatgtttta	4800
	ttagttctat	gcctgcactt	actggtcgga	agagagaaag	aacagtttca	gtatctgagc	4860
25	ggctcaggag	aggtctcttg	atttttaaa	ttttgggttg	gggggtgtgt	tggtgtttca	4920
	ttcttttgaa	ttttaattta	gggtgttttg	gtttttttcc	tttaagaga	atagttttca	4980
	caaaaattga	gtgtgtcttt	ggcttttgtc	ataagggaag	cagagtggcc	tggtctgatt	5040
	gataaaatgt	tttctctctc	tcacacacac	cacattttgc	ttttaagtga	cttacttttc	5100
	ccacttgagc	atcttgaaac	tacttttttt	ccaaataaat	tactcatcct	taaaagttaa	5160
30	tcacatttga	caaaagatac	gcctctctcc	ctgcaacata	agcaggttgt	agaacgtggc	5220
	attctttggc	aagttagtag	actttaaccc	gtctctttcc	ttttttgtgt	atgtgtgtcc	5280
	ttctctctct	ttctctctct	ttctctctct	ttctctctct	ttctctctct	ttctctctct	5340
	gtctctctct	ttctctctct	tttgagcact	ttgttttgaa	aaaatogttg	agatgcocaa	5400
	gaaccttgag	taattcttta	cttttttttg	aataaaggaa	aggaatttca	aaaaaataaa	5460
	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaa	5515
35	<210> 6						
	<211> 6131						
40	<212> DNA						
	<213> Homo sapiens						
45	<400> 6						
	gaattctatgg	ccaggtttct	tgtttccccc	gtgtgttctc	aggcagggtc	gtttccctcc	60
	atgggctctc	gtaagatgag	gagttggaga	ggtaacttct	caggctactc	tcacatccca	120
	gccaaagtac	tcagagtgcc	caggccagac	cagcacccct	atctccaaag	ccctctgagt	180
	tggtctctct	tttagaacct	aatccacact	acccaacac	agtcagtgct	cttaaccagc	240
50	caaggtccct	ggggagccct	ctagagggag	agagccctgc	ccaccagatg	tgagggttaa	300
	ggcctccccc	tgctcatttt	tgtaaccaca	agctgcttgc	cacttggtag	acatcaaaaa	360
	gtgtgtgtgt	aagaataaat	tgaagtgtgt	tatatatgtc	agctagagtg	tctggagggg	420
	cagaaatgtg	ggtctcaaac	atacaaatgc	tcacaaatgg	gtgtgggcaa	gggtctgtct	480
	acacccagct	gtgattcact	gtccacatgc	atgtgtctat	ctgagttagg	gtatgttctc	540
55	tattttttta	ccaccagggg	tgaggaaacg	gtatgtgtgt	gcattgtgat	cactccogtg	600
	gtgtgtgtgt	gtgtgtgtgt	atgagtgtgt	gtgtgtgtgt	ccaaaggccc	ctcttccaac	660

	tgtgccattt	gta totgtgt	ctggcccaat	gagagtgttg	aaaggtgago	cacaagataa	720
	aacagcaact	tctactctcc	cttatcaaga	cagctgtctg	acctaactcc	ccttgcccaac	780
	tcttggaatt	actgggggtg	gcttcagtat	tttcagattt	ttcagaaggg	gaggagaaatg	840
5	ctcaggtctc	atccaggaaac	ttaggcagtt	ctcagcaactg	ctcgtctctc	ctccctcaaa	900
	taaacaagtc	tgaagacacg	gagagaaagc	cgctggttga	ctggtcaact	gtctggcagtg	960
	gagtgaggga	gagtgtgagc	ttttatagta	ggaaatccaga	cttagacact	ccctcccaac	1020
	cccgatctgg	tggtgcacag	gctcatctcg	cgcccccctc	ccactccacc	ctaacatctga	1080
	tacgccccca	acaaaccaag	aaagatctcc	catcgctga	ctccacagat	caacacatgt	1140
10	cccccaagac	aaacacacgc	ccatgcacag	gcacagacat	ccaggacacat	ctttcccttt	1200
	ctctgtcttt	cccttggttt	gaatttctgt	tagccacata	tgttgttgtg	gcgtggagggt	1260
	gggtggggga	ggggccagaca	gggagtgggg	atggcatgtg	gccaacatct	acctatctggg	1320
	ctcggggcag	gtagcgccct	tacagccatc	ctggggaggg	gtctcagctg	tccttttgtg	1380
	gccaagggag	ccctctctgg	gagtgggggc	aagcaacagag	gtctttctc	cccaacccgg	1440
	ggctctggcc	ctgacccacg	ttgggggccc	gcaggggagg	aaatggacag	agcggggaccc	1500
15	tgaggggaga	tagaattggc	caacacagag	ccccagtgct	cagccttgcc	acccatttgt	1560
	tcocgtgagg	gggtctctat	atcacagggg	caactccctc	caacttctc	tcacttctc	1620
	cttccctgc	gttggggcgg	gagggggagg	cggcagaaat	atattattat	ttcctttatt	1680
	tatttaattt	tttttttttt	tttttggagt	agagagtgac	agatggcggc	gggtccgggg	1740
	ggagccggct	ctcccccagt	gcagacgcac	gcaaatccac	gtctctcatg	tgatctctgc	1800
	tgcccgtagc	gtgcacagcc	catatggcct	ggcatagagg	ctgggtacccc	gctctgtaga	1860
20	gatgcacac	tgctctccgg	gttcgcacgg	cgctctgaag	acgcggcgcc	cccgccggct	1920
	gaggagccgc	tgccccccgc	ccctgaagat	ggggggaaca	tgaataaagc	gagaagatcat	1980
	ctctctctcc	ccctctctct	cttgcgccct	cccccctccc	ccctccctcc	ctccttgact	2040
	ccctctccag	gtaagtgtgc	cgaaaggagg	cgagatctga	cccgccgggt	gggagagggg	2100
	cgccgagctt	cgccgcacag	gagggtctct	aaatacctcc	ttcctggagt	gatcgccccc	2160
25	tcattgggtg	ggcatctggg	gggccccagg	ttctctctcc	cttagggggt	gcagcccaagg	2220
	gggtctcgag	ggaggtctct	gtgcctcgga	tgggctcggt	ggggggggaa	cgaggagatac	2280
	cgaggagggg	tatgcggaag	ggccgagacg	gaggacccct	ccatggttgt	cccaaaaagc	2340
	ctgcacactt	tcocccacac	cgaaaaaagg	gaagcaaaac	aaacaaattg	gatttttccc	2400
	ccattcaatc	caaaatacaa	cgagatctga	agagccctgt	gggagggag	cagcttgaag	2460
	gggggaaggg	gtccctcgag	cgagaggggg	cgactgtctc	tgctctctc	cagctctctc	2520
30	ccaccgcccc	gctgctctag	tcctcgccgc	ccagagccgg	ctccgggagc	tgggggacgc	2580
	tcggcttagag	gagacgatcc	tcocccctct	ggaattgggg	gtcggggggg	ggggggccag	2640
	caaggggcgg	cgccgcagca	agttgcacaa	tggattaggg	agcgtggggg	tgagagccac	2700
	gggggggggt	agggagcttg	gcgggggggg	ccggggccgc	agagcccgga	cgggggcgac	2760
	tgtcccccac	ggcgcccgag	cagcctctct	ccaccgcacg	gagagaaocg	gctttcaggg	2820
35	cgagcgccgc	gcctccctgt	gcacaaat	ctggtcccta	aaacccccc	ccggtccctg	2880
	ccctgacact	gagaagaagc	agggcggggg	agcagccccc	caatcaagcg	agggggggag	2940
	ccggggccca	gcgcggggga	gagggccttg	gcgcagatcc	caggccggga	gcggggtagg	3000
	gctggggcgc	ctctggggcg	ggcaggccgc	ggaggtgggc	atccagggtg	ctcaggccag	3060
	gagcccgccg	gagactcggt	gggtggaggag	ggttgtgggg	ggggctcggt	acccagacgc	3120
40	gcocctcaat	ttgtgctgtc	tgtctccctc	tcoccccgcc	ggggcgccct	gagccaacat	3180
	gctgacccgc	ctgtctcagc	agcccgccct	tcctctcgag	gtgcccaaat	tcggcagctg	3240
	ggcgagccgc	gaagacgagc	agccgaggag	cgacaagggc	gcagcccgcc	ccacccacac	3300
	gcctcgccac	gggcacgggg	ctccggggcc	agcccgggcg	gccacagccg	tcctctctcg	3360
	tggaagaagc	gggcgcggag	ccacgttgcc	cgagctcaag	gaggaaggcg	agctggggggg	3420
	ggcaggagg	gaggaagagg	aggaaggaga	aggaactggc	gagggcgagg	gcgagcgccc	3480
45	caagagagcg	gggcaccaag	agcgcaagat	gaccacggcg	cgcttgagcg	ctcccaagct	3540
	tcggcgccag	aaggccgaag	cgccgggagc	caaccccgat	cacgacccga	accagccctc	3600
	ggacacactc	cgcaacgttg	ctagcccaag	gtccactgta	ctccaagagc	cagaagctgt	3660
	cgctgctggc	ctagctgtct	acgtgcagac	ggcgctctcg	gagatctctg	gtcccgccaa	3720
	goggccagac	ctagctgtct	tgacgtcaaa	ctctgtcaac	ggctgtgctg	agcccaacac	3780
50	caa tctgggt	gcggctgtgt	acggctctgg	ctctgtcaac	ttctctcagg	agcaaggcgc	3840
	cgacggtgcc	ggccgctctc	acggctctgg	cgcccgcttc	gccaatgccc	ccctaacgctc	3900
	cccgctgctc	cgctctggcg	ggccacagtg	ccaggcgccc	ggcgccctgg	ggcgccggcg	3960
	ggcgccagcc	ctcgccgaac	acggctactg	cgcccgctac	gagacgctgt	atgctccggc	4020
	agggcggtgc	gcgcgagacg	cggaactcaa	cagctctcgc	taacagggcg	gcgcgagccc	4080
	cccgctctgt	ctcaa tggca	acttctcaat	caagcaaggc	tcctcgcccg	accaagagaa	4140
55	aagctacac	taactatgac	actactccgc	gctgcocggt	tcgcggccag	gcaacgggct	4200

	agtcttcagcg	tgcgtggcgt	tgcgcggggg	cgccactcgt	gagaaatctct	tgtcttaccga	4260
	tatgcacactt	caccacgacc	ggggcccccac	gtacgaggag	ctcaatgcgt	tttttcaata	4320
	ctgagaccttc	gocgcgggtc	cctctctttt	cttttgccct	tgcgcgcgcgc	cctgtcccca	4380
5	gcccacagca	gocccaggta	caaccccaac	ctaccccgcc	tgcggggcgcg	gggagcgggc	4440
	cacgcgtctct	gcgcgtctct	tggggcgagcg	cagtccttgt	accgttggtg	ggcctgtccc	4500
	aggggcctcgt	cttcccccag	gggactgcgc	ttctctctcc	ccaagggtgt	ccctctctct	4560
	ctctcccaacg	gagtgctctc	ccagggaacct	ctctccgggg	ggtccctgga	ggcaccctcc	4620
	cccatctccc	aatactctcg	ctgaggtttc	ctctccccc	tcctccctgc	agggcccaagg	4680
10	cggttgtaag	ggggcgagctg	agcaatggaa	cgctttccc	ccctccatta	ttattttaaa	4740
	aacagacacc	cagctgcgcgc	ggcaaaaaagg	agccaggcgc	tcctcttttc	ttgaagaggg	4800
	tagtattttg	ggcgccggag	cccgggcctcg	gaaocgcctc	accgcgaacc	tcagctctcc	4860
	gcgtttttg	atttcaattt	tggcgggagg	ggaagtggat	tgagaggaaa	gagagagccc	4920
	aagacaattt	agactagaaa	tcogtttttc	ccctttctct	tttttaaaaa	aacaaacata	4980
	aaaaaaataa	aaaaaaataa	aaaaaaataa	aagctaaag	gcgaacgaa	ccgaacgaa	5040
15	agtcggatc	ggagagaaaa	cgcagtaagg	acttttagaa	gcaataaaag	gcacacacata	5100
	caaaaaacaa	aaaaaacacaa	aaaaaaaac	ccctactacc	ataatacaaa	gcacacacata	5160
	ctctatgcgc	gaggtctccac	tgagcctcgc	ggcccgccgc	ggcccgccgc	tgcccgccgc	5220
	ggcctgcggg	ccgcgcgcgc	cgagcgcgga	tcgtgcact	ttggtgaagt	ggggggccgc	5280
	gcgcgcgcgc	cccccctccc	aggttcttac	aatacgtgac	tcggagaatt	ggggcccgag	5340
20	tgccactgc	ctcccccgc	ccgtcccgct	tgctgcact	gcgtgttttt	aaaaaacctg	5400
	ttccaaattt	gtaggaatg	gcacaaactgt	gggggggtgg	tttggtggag	gaggggttgc	5460
	atgaagagca	caocgacac	acacgcgcgc	ccacagcagg	cccgccgcgc	gcgtccgggg	5520
	ggcagaagga	ggtgagctcg	ccgcctctct	ctcccgcgcg	ccattctgtc	ccctctgggg	5580
	gtgaggggtg	gggtgagaga	ccgtggggca	gccccccccc	tgcccgagat	gtgcctcggt	5640
	gggtgcaccc	tgccgatttc	ccgtgtctgg	agagagtatt	ttttgggtcca	aggaagtctc	5700
25	ttggctttga	ctggtgggtg	ggcggggaga	ggtctgaggg	ctcctactgg	aggtttccccc	5760
	aaaaagggcg	aaagaggagac	ccctgcgcgc	ccggagcgag	gggtacagac	atccaaatcc	5820
	acgatgcaca	aatgcacatc	caacaggcgac	acacccacac	actcccccac	acacacagca	5880
	ttttaccttc	ctctgttagc	gaagatgaaa	ctcccgctgg	acaccccgag	tgacatctcg	5940
	gtttctgttc	agtttaatag	cgatttaata	attattatgt	aaatgagatg	caaagcccgga	6000
30	ccggtttctc	acggtggcct	catttcattg	aggggggaga	gaaggtttga	gctggggctg	6060
	gggtgagtaa	gcagagtgct	caagtgtactg	tgccagggcc	aaacagaggg	acttccacgc	6120
	aaaaagcact	g					6180
	<210>	7					
35	<211>	2020					
	<212>	DNA					
40	<213>	Homo sapiens					
	<400>	7					
	gactactgag	cgcgcggagc	ggactgcggt	tgggcgggga	agagccgggg	ccgtggctga	60
	catggagcag	ccctctgtct	gaggccgcgc	ctcccccgc	ctgaggttgg	ggcccccagc	120
45	gatgagcaag	gcgcgcagg	agctgacccg	agacttggag	ccgagcctgc	ccgccttgcg	180
	ctccctgggg	tcctcactgt	cccacagcca	gagcctctcc	tcgcacactc	tcgcgcgcgc	240
	tgagaagcga	agggccatct	ctgatgtccg	cgcacacttc	tgctctctcg	tcacactcga	300
	cctgtctctc	atctccctgc	tcctggatcat	cgaactgaat	acccaacacg	gcactcgtga	360
	gaacttggag	caggagatca	tcagtcacaa	ctttaaaaat	tcctctctct	acacttttgt	420
50	ccctggcttc	cttgcgtctc	ctggactgct	ccctaggctat	gcgcgtctgc	agctccggca	480
	ctgggtgggtg	attgcggtca	cgaacgtggt	gtccagtgca	ttccctactg	tcacaggtcat	540
	ccctcttgag	ctgctccagca	aaggggcaatt	tggtcactcg	ctcccccctg	tcctcttttgt	600
	ccgcgcctgg	ttggagacac	ggttctctga	cttcaaatgc	ctaccccagg	aagctgagga	660
	ggatgcagtg	tatcttgccg	cccaggttgc	tggtgcgcct	ggacccctgc	tggtctctcg	720
55	tgctctgtcc	gagggacagt	ctatttccac	cccagaatcc	tttgacgggt	ctgacaaatga	780
	atcagatgaa	gaagtgtcgt	ggaagaaaaa	ttctctctct	caggagcggg	agtcactccg	840

	ccagggggaag	gaggccoacgg	cagtgggtgga	ccagatctctg	gcccagggaag	agaactggaa	900
	gtttgagaag	aataatgaat	atggggacac	ogtgtacaco	attgaagtgc	cotttccacgg	960
	aaagacgtttt	atcctgaaga	cottcctgccc	ctgtcctgog	gagctogtgt	acccaggaggt	1020
5	gatcctcgag	cccgagagga	tggtgctgtg	gaacaagaca	gtgactcgct	gocagatcct	1080
	gcagcagagtg	gaagaaca	cctcctatctc	ctatgacgtg	tctgcagggg	ctgcggggcgg	1140
	ogtgggtctcc	cccgaggact	ctgtgaaatgt	ccggcgcatc	gagcggcgca	gggacogga	1200
	cttgtcaatca	gggagtgcca	cctcacaacag	tgccaagccc	ccgaagcaca	aatatgtcog	1260
	gggaggaat	ggcctctggg	gcttcaatgt	gctcaagtgc	gocagtaacc	ccogtgtttg	1320
	caocctttgtc	tggtattctta	atcacagatct	caaggggcgc	ctgcocccgt	acctcattcca	1380
10	ccagagctgc	gggcoaccca	tggttgaaatt	tgcttttca	gctcattcgc	gcatcagcga	1440
	gctggggggcc	cgggcgtgac	tggtgcccct	cccaacctgc	gggcoagggt	cctgtcgcca	1500
	ccactctccag	agccagaag	gggtgccagtt	gggtcgtgac	gtgcccactg	tgacotggcc	1560
	ccaggctgtc	acccctcaacc	gagccacgca	gtgctctggag	ttgactgact	gagcaggctg	1620
	tggggtggag	caotggaact	cggggcacca	ctggctggag	gaagtggggt	ctggcctgtt	1680
15	gatgtttaca	tgggcgcctg	cctcctggag	gaaccagattg	ctctgcccaca	ccttgccagg	1740
	gcagggtctg	ggctggggcac	ctgacttgcc	tggggaggac	caggggccctg	ggcaggggcag	1800
	ggcagcctgt	acccctgtgt	aagatgaagg	ggctcttccat	ctgcctgggc	ctctgtgggt	1860
	ttttttttaga	ttattgaag	agtctgggac	cctgtttggg	gagtggtggg	caggtgggggg	1920
20	tggtgctgtc	gcatgaatc	tgtcctcttc	ccaggctgtc	ccctccctcc	cagggccctcc	1980
	tgggggacct	ttgtattaa	ccaattaaaa	acatgaattt			2020
	<210>	8					
	<211>	1730					
25	<212>	DNA					
	<213>	Homo sapiens					
30	<400>	8					
	gtgggtgaggg	tgactggggg	ctaggcaacta	ggcctttgggt	gcaggggccct	gagggaactgg	60
	ttgcaactctc	cctctctggg	atatgcccct	gagccaccgg	agagggagagc	acagcccagg	120
	gcagggaactg	gcagcctctg	tacagagccc	agagggggga	tcagttctctg	ctgtctcctgc	180
35	ctctgtttaca	gacaaactgc	tgctcctcct	gcacaaaggga	gtgggtggggg	caggaggccaa	240
	ktgcacagggg	ggcacaaggg	tgggcatgtg	gctggcaatga	gaocgtgtct	gagttaattg	300
	aggcaactcgg	aggcattgac	ccagggaact	tggaacccag	acctctgacc	gtggggcagc	360
	cagcgtccag	gtaccccaac	cctgcctctg	ggtccggcgt	ccccccatta	gtgagtcttg	420
	gctctactta	tagcatctga	caacagaggg	gcgaaaaata	gccccctggag	aaggggggagg	480
40	agggggcctat	ctgaacactgt	tgggaggggga	gagagaatga	ggagtgtatca	tggtacacct	540
	agagctgagc	tgogagggtg	cgaggaggaa	ctgtgagcgc	cgggaggccct	ctctgggcaga	600
	atggaagagat	ctgacactgt	ccacacggcc	cgaggagggg	tgagtgtggg	ctctgtagag	660
	tcctctgctc	tgctggcccca	gagcacccct	actgagccat	gaggccagag	catgaagccc	720
	tggaagaatt	ctctgggggtg	ggggcaggaa	gaatgcccaca	tggggagagc	aaagggggaac	780
	caocctctcct	ggcccccagg	ccagcagccc	cagggggagcc	ccccacccag	cctgtgcccac	840
45	gagagcaact	gctccacagg	gctcactgac	cctccctctc	ccccagctgc	tcctgtcatg	900
	aggagggacac	ccagagacatg	gagacctaac	acccagcagg	cgactgtccag	ctgtcgggtgc	960
	agcctgcctca	ctggctcgtg	atgggtagtg	gcatcctcgg	cogtgggctg	gagagtaacc	1020
	agctgcctca	ccagggggtg	ctgcctctgc	ccatcttccac	cctgcacaa	atggggccca	1080
	ccaggaggga	gogtgaggac	gaacccatgc	aggttcagga	aggtggctga	ctggagacag	1140
50	ccactgtgtg	tgctgagcag	gaacccaggg	ctgtgtggcg	gactgtggct	ccactgtccc	1200
	ctgtgtgtgac	gttcagcaag	ccgggtgccc	ttogtgctct	gagtcacaa	ccactgtccc	1260
	aggaagacaca	gctccacagg	gagggactgt	gacttggtgt	ccgtgtggcc	cgcccccctgg	1320
	gctggggccct	tcctggctag	gaacctgtga	ggggcagctc	gctggcccat	ggctgctttg	1380
	tagtttgccc	agagtgtggg	gctaggggag	gggggagcca	gagggccagc	tgccctgagcc	1440
	ccctgagttc	ccaaaggagg	ggtggcagag	acagtgggca	ctaaagggtg	agagttgggg	1500
55	gccaagccag	ctgagggaacc	tcaggcccag	gagaaggggac	aaaagggtact	ggtgagggca	1560

agagggtgcct	gggaggagtg	gccctgatcc	aggaaaaatgt	gaggggaatc	tggaacgctc	1620
tggcagagaag	aagctgtggag	ggagggggag	gtgaaaaagg	cagaggcaag	gatgggtggg	1680
cccccagcac	ccctctgttag	tgccgcacata	aatgtctcaat	catgtgcacag		1730

<210> 9

<211> 3799

<212> DNA

<213> Homo sapiens

<400> 9

ctggcactgg	gtggtaaacca	gcaagccagc	tggcatccgc	atccagggtt	tgtttcaatg	60
atgtctctctg	gagaatatgg	aggggctgtg	gccaggactg	tocttggctt	tgctctgggg	120
tgtgaacggg	gtcagtgacc	tctaaaacta	acctgcctct	cagttcttgaa	tcacagacga	180
atcaattctc	agctgtgtct	cgctccacac	ccctgcctcc	ggagcccgga	gaaggcttga	240
gggtctaggg	ggtctaggctc	ccctctgtga	ccctgcagc	tgttgggtgtg	acctcatgtcc	300
caactagatg	cgctctccca	aggagacttt	ccctctggga	aaagggggag	gaatggcatg	360
gaggagggccc	acatacaagc	gggccaaggaa	cccaagggtg	caggagctgg	gctggtgacc	420
tacccagggc	agaaggccc	gggactcaatc	cagaggggaa	ggaggggttc	ttcaggaaga	480
ccaagagatg	gccacaggca	gaattgtgctt	cccatctggg	agatagggtg	ggagaccctg	540
gcattttgac	gcccagaacc	tggggtgtgtg	agcagaaatc	tcatgtcctg	ccctggccgcc	600
ttggaggagg	agctggagg	ttgggtgtgga	gaggagtggtg	gtcagagccc	ctcaatctcgc	660
aggaccacca	atcggtctgg	ccccaaggcc	cggactgggc	tcctcgtctg	ccctggccgcc	720
ctctccggaa	tggttctcgt	ccctccctctg	cccaagccct	ctgcccctac	ccgggtctcgg	780
cgccgtccccc	gaagtggcgg	gaaccaaccc	aaacccgaac	ttctgtctctc	gggagccccc	840
ctgaataagc	ctgggaacccc	cgggggccccc	cagggggaggc	ccgggtctctc	cgcccgctaa	900
ctgtcatatgc	acagctccac	tcctccatctg	cgctgtccat	cggaagggca	gtgcgcgcgc	960
ctgtctctggg	gcgccccagg	cgaccaacgc	ggaggccggga	acggactgtc	ctttctgggg	1020
cggggtgtggg	aggggtgtgt	gctggagggc	ccggttggcat	agcaacggac	gagagaggcc	1080
tgaggagagg	ggggggagg	ggagtgtgtg	ggcagttcta	agggaaagggt	gggtgtctggg	1140
acgggtgtcc	gggaggagg	ggagcctggc	gggtgtgtgg	gcctgtctgc	ggagggccgt	1200
gcgaggggga	aaactgtggga	agggcctaata	tcoccatctc	ccactctgaa	tcaggaaga	1260
gaagggggcg	gctgtctggc	aaaagaggtg	aatgtctgtc	gggggtctgga	gaagagagat	1320
gggagggggcc	ggcggggcg	ggtgaggggg	ttctaaagatt	gtgggggtga	ggaaactgagg	1380
ctggggggcg	cccagaggcg	ggactcgggg	cggggcaggc	gaggggagg	gcgagggctg	1440
cgggagcaag	taaggagccg	ggggtgtggg	ggaagatgtc	cgctgcagcc	gcgcccccac	1500
tcacctccgg	tgtgtctgca	gcccggacac	taaggagagat	ggatgaatgg	gtggggaggga	1560
cgggagccgc	atggccccc	cgggctctggc	ggtcagctgc	cgcccccaca	cggaacccgt	1620
cgggggcggg	gtcggggcgt	agaaaaaagg	gcgcgcaggc	gagcggggca	ctggggcgac	1680
cgccggcgga	cgatgagcgt	cgacagccgt	agccccaatg	cgggcgccgc	ccctgactcg	1740
gcocccggcc	agcgcccggt	ggctctcggc	taccagcgct	togagcccg	cgccctacctc	1800
cgcaacaaac	acgcgccccc	tcggggggac	ctgtgtcaac	cgaaocggct	cgggccgtgg	1860
aaagctcgct	gcttggcgca	gacctctcgc	accggtgagc	gggggaaact	gaggccagag	1920
gggacagagg	tcgtctgggga	gtgaaagcag	gcgcagggaa	ataaaaagaa	ggaaaggagg	1980
acagacaggc	cgctcaacag	atgggggaaca	agaacaagaa	gatagctgag	aggtgcaaac	2040
agaagagaaa	aaaggagcaac	atcccttagg	agagggcgag	aggagagaga	ggtggagaga	2100
ggggggcgag	agtgtctcaga	attgagagct	aaagtgtggg	atgcaggaca	gactgaggtg	2160
gagatgtcta	ggaggaaatg	gaggcagatg	tgggacaggg	tttagaaact	ccaggatttc	2220
ctcgtctgagc	ctggctgtgtg	ggatagattg	ttttctttat	ttttctttat	tttattttca	2280
ttttatttact	tattttttatt	ttttatttgg	tttgagacgt	agttttctgc	ttgttgcoca	2340
ggctggagta	caatggcgcc	atctcggctc	actgcacact	ccgcctccccc	gggtctcaagc	2400
gattctcttg	ccctcagcttc	ccctagtagct	gggattacag	gcattgcggcc	ccatgctctcg	2460
ttttattttat	tgtatttttta	gtagagacgg	gaattctcca	tgttggctcca	gtgtgtctcg	2520
aaactccaac	cttaggatcc	acccaccccg	gcctcccaaa	gtgctgggat	taacaggtgtg	2580
agccacttgg	cccgccaggt	aggtatagtc	ttctagatgt	gaacactgag	ttctcagagc	2640

	gtgaagtcc	cttccgaagg	gcagcccatg	tggagctgg	gttcagctcta	actctggggc	2700
	caatgctttt	tccagatgga	gacacatttg	cagaggagaa	ggagaacta	gagagaggca	2760
	gggagatgca	ggggagggaa	gggttaaggag	gcaggggctg	cctgggctgg	ctggcaaccg	2820
5	gacccctctc	ctctgcccctg	cccaggtgaa	gtgtccggac	gcacccctat	cgacattggt	2880
	taaggcccca	cogtgtcaoca	gctgtccagt	gcoctgcagcc	actttgaggga	catcaccatg	2940
	acagatttcc	tggaggctcaa	ccgcacaggag	ctggggcgctc	ggctgcaggga	ggagccggggg	3000
	gocctccaact	ggagacatgta	oagccaaacat	gocctgoccca	ctgaggggcaa	ggggtaaggga	3060
	ctgggggggtg	agggttggggg	aggaggcttcc	ccatagagtg	gctgtgtggg	gcaacagagg	3120
10	cotgagcgta	gaacagccctt	gagccctgcc	ttgtgctctcc	tgccacaggga	atgctggcgag	3180
	gataaaggagc	gcacagctgcg	agccacgggtg	aaaocggctcc	agtcaccaagc	cgctgcacagc	3240
	ccccagccccc	tgggtgtcggg	gagcccaagct	ccocctgocctg	ctgacgcocct	ggctctctggc	3300
	ttctgtcttgg	aggctgtgag	cccagatctct	gocagcttctc	agocggccctc	ggacccaatc	3360
	acccaogctgc	tgaggccctgg	ggggccacccc	ctccctcatcg	gggcccctgga	ggagtctgtgg	3420
	taacctggctg	gggaggccag	gctgacgggtg	gtgccagtgct	ctgaggaggga	ggtagggagg	3480
15	gcccctggctg	gtagtggctca	caagggtccgg	gacccctcgca	ccctatatcat	gocctcccac	3540
	cttcagacag	gcgtagatgta	tgtcaaggggc	gtctctctctg	cctggggctga	gaagggttggg	3600
	ctgtgaggggc	tgttaacctgtg	gcccctgtggc	cccccccccac	ctggattcccc	tgctcttttga	3660
	agtggtcacct	aataaagaaa	taataccctg	ccgctgggggt	cagtgctgtg	ttgtggctctc	3720
	ctgggaagcca	gcgaaggccc	agagatctga	gtgtccgggt	agggggagaca	tccacccctag	3780
20	gcttttttttc	cagaagcttt					3799
	<210> 10						
	<211> 4530						
25	<212> DNA						
	<213> Homo sapiens						
30	<400> 10						
	aattctctgag	ctcgtcgacc	ggctgaagag	ctcaggggtc	gacgagctcg	agggcgccgg	60
	cccgccccc	acccctcgca	gcaccccggc	cccgccggcc	tcacagccgg	gtccagccgg	120
	agccatgggg	coggagccgc	agtgagcacc	atggagctgg	agcccttgtg	ccgctggggg	180
35	ctccctctgc	ccctctgtcc	cccccggagcc	gagagccccc	aagtgtgca	cgccacagac	240
	atgaagctgc	ggctccctgc	cagtcocag	acccacctgg	acatgtctcg	ccacctctac	300
	cagggtctgc	agggtgtgca	gggaaacctg	gaatccaccc	acctgcccac	caatgcacg	360
	ctgtctctcc	tgccaggtat	ccaggagggtg	cagggtctacg	tgctctatgc	tcacacccaa	420
	gtgaggcagg	tcocactgca	gaggctggcg	attgtgcgag	gcacccagct	ctttgaggac	480
	aactatgcgc	tggccgtgct	agacaaatgga	gaocccgtga	acaaatccac	ccctgtccaa	540
40	ggggccctccc	caggagccct	goggggagctg	cagctctgaa	gocctcacaga	gatcttgaaa	600
	ggaggggtct	tgaatccagc	gaaccccagc	ctctgtctacc	aggacacgat	tttgttgaaa	660
	gacatctctcc	acaagaacaa	ccagctggct	ctccacactga	tagacaccaa	ccgctctcgg	720
	gocctgccacc	ccgtgtctctc	gatgtgttaag	ggctcccgct	gctggggaga	gagttcttgag	780
	gaattgtcaga	gcgtgacgcg	cactgtctgt	gcoggtggct	gtgcoccgctg	caaggggcca	840
	ctgcocactgc	actgtgtccca	tgagcagtg	gtgcocggct	gcacggggccc	caagccactct	900
45	gactgcctgg	ccctgocctca	cttcaacccac	agtgccatct	gtgagctgca	ctgcccagcc	960
	ctgtgtcacct	acaaacacaga	caogttttgag	tcacatcccca	atccocagggg	ccggtataca	1020
	tgcggcgcca	ctgtgtgtgca	tgccctgtccc	tacaaactaac	ttctacaggga	cgtagggaac	1080
	ttgcccctgc	ctctgcccctc	gcacaaaccaa	gaggtgacag	cagaggatgg	aaacacaggg	1140
	tgtagaagtg	gcagcagccg	ctgtgcccga	gtgtgtctatg	gtctggggcat	ggagcaactg	1200
50	cgagaggtga	ggcgagcttac	cagtgccaat	atccaggagtg	ttgtctggctg	caaggaagtc	1260
	tttggggagc	tggcattctct	gocgggagagc	tttgtagggg	acccagctctc	caaacctggcc	1320
	ccgctccagc	ccagcagccg	ccaaagtgttt	gagactctgg	aaagatacc	aggttaacct	1380
	tacatctcag	catggccgga	caocctgocct	gacctccagcg	tcttccagaa	ccctgcaagta	1440
	atccgggggac	gaattctgtca	caatggcgcc	tactcgctga	ccctgcaagg	ctgtgggaatc	1500
55	agctggctgg	ggctgcgctc	actgagggga	ctggggcagtg	gactggccct	catccacatc	1560
	aaaccccacc	ctctgtctgt	gcacacgggtg	ccctggggacc	agctctcttg	gaaccccgac	1620

	caagctctgc	tcacactgc	caacogggcca	gaggacagtg	gtgtggggcga	gggcoctggcc	1680
	tgccacacgc	tgttgogccg	agggcaactgc	tggggtccag	ggccacaccca	gtgtgtcaac	1740
	tgacagccagt	tccttggggg	ccaggagtgct	gtggagggaat	gcgcgagtgct	gcagggggctc	1800
5	ccacggggagt	atgtgaatgc	caggcaactgt	ttgcogctgoc	acccctgagtg	tcagcccccag	1860
	aatggctcag	tgacccgttt	tggaccggag	gctgaccagtg	gtgtggcctg	tgccccactat	1920
	aaggacccctc	cccttctgct	ggccogctgc	ccacgcgggtg	tgaaaactga	ccctctctac	1980
	atgcccactct	ggaagtttcc	agatgaggag	ggcgcatgccc	agcccttgccc	catcaactgc	2040
	accccactctc	gtgtgggaact	ggatgacaag	ggctgcccog	ccgacagag	agccagccct	2100
	ctgacgttcca	tcgtctctgc	gggtgttggc	attctgtctgg	tcgtgtgtctt	gggggtgtgtc	2160
10	tttgggatacc	tcactcaagcg	acggcagccag	aagatccggg	agttacacgat	gcggagactg	2220
	ctgcacggaaa	cgagcctggt	ggagccogctg	acacccatagc	gagcgatgcc	caacacggcg	2280
	caga tgccag	tcctgaaaga	gaocggagctg	aggaaggtga	aggtgctgtg	attctggcgt	2340
	tttggccag	tcctcaacggg	catctggatc	tcctgatgggg	agaaatgtgaa	aattccagtg	2400
	gcca tcaagg	tgtttgaggga	aaacacatcc	cccaaaagcca	gcctctctggg	cttagacgaa	2460
	gcatacgtga	tggctggtgt	gggtctcccca	tatgtctccc	gcctctctggg	catctgctgt	2520
15	acatccacgg	tgccagctggt	gacacagctt	atgcccctatg	gctgcctctt	agaacatgtc	2580
	cgggaaaaacc	ggggacccct	gggtctccag	accctgtgta	actgtgtgtat	gcagattgoc	2640
	aaggggatga	gctacccctg	ggatgtggcg	ctogtaacaa	gggactctggc	ccgtctgggaa	2700
	gtgctgtcca	agagtcccaa	ccatgtccaa	attacagact	tcgggctgtg	tcgtgtgtgt	2760
20	gacattgaag	agacagagta	ccatgcagat	gggggcaagg	tgccccatca	gtggatgtggc	2820
	ctggagatcca	tcctcogccg	gcgggttccac	caccagagtg	atgtgtggag	ttatgtgtgt	2880
	actgtgtggg	agctgatgac	ttttggggcc	aaaccttaag	atgggattccc	agccccgggag	2940
	atccctgacc	ctgtggaaaa	gggggagcgg	ctgcccagcc	cccccatctg	caccattgat	3000
	gtctacatga	tcattggtcaa	atgttggatg	attgactctg	aatgtoggcc	aagattccgg	3060
	gagttgtgtg	ctgaattctc	cgcctatggc	agggaccccc	agccgttttgt	ggtcatccag	3120
25	aatgaggact	tgggccacgc	cagtcocctg	gacagcaact	tcacacgctc	actgctggag	3180
	gacgatgaca	tgggggaccc	ggtggatgct	gaggagatca	tggtacccca	gcaggagcttc	3240
	ttctgtccag	acccctgccc	ggcgctggg	ggcatgtgtc	accacaggca	ccgcagctca	3300
	tcacacagga	gtgcggctgg	ggacccagca	ctagggtctg	agccctctga	agaggagccc	3360
	ccacaggtctc	cactggccac	ctccgaaggg	gctggctccg	atgtatttga	tgtgtgacctg	3420
30	ggaatggggg	cagcccaagg	gctgcaaaag	ctcccccac	atgaccccag	ccctctacag	3480
	cggtacagtg	agggacccccc	agtaacccctg	ccctctgaga	ctgatggcta	cggttgcoccc	3540
	ctgacctgca	ggccccagcc	tgaatatgtg	aaacccccag	atgtctggcc	ccagccccct	3600
	tgcgcccgag	agggccctct	ccctgtgtgc	cgacctgtg	gtgcccactct	ggaaaggccc	3660
	aagaatctct	ccccagggaa	gaatgggttc	gtccaaagag	tttttgccct	tgggggtgoc	3720
	gtggagaccc	ccaggtactc	gacaccccag	ggaggagctg	cccccagcc	ccacccctct	3780
35	ccctgootcca	gcccagcctt	cgacaaactc	tattactggg	accaggaccc	accagagcgg	3840
	ggggctccac	ccagacactc	caaaagggaca	ccctacggcg	agaccccag	tcacctgggt	3900
	ctggacgtgc	cagtggtgaac	cagaaggccca	agtcocgaga	agccctgatg	tgctccacgg	3960
	gagcagggaac	ggcctgactt	ctgctggcat	caagaggtgg	gaggggccctc	cgacccactc	4020
	caggggaaacc	tgccatgcca	ggaaacctgtc	ctaaggaaac	ttccctctctg	cttgaggttcc	4080
40	ccatgggtctg	gaaggaggga	agccctgttg	gaagagggaac	agctctctgtg	agctctctgtg	4140
	gattctggag	ccctgcccac	tgagactcta	gggtccagtg	gatgcccacg	ccagctgtgg	4200
	ccctttccctt	ccagatgaaa	ggtactgaaa	gccttagggga	agctggcctg	agaggggggaag	4260
	ggcccccata	ggagtgtcta	agacacaaaag	cgacccatcc	agagactgtc	ccctgaaacct	4320
	agtcactccc	cccatgagga	aggaacagca	atggtgtccg	tatcccagct	ttgtcacagag	4380
	tgctttttctg	ttttagttttt	actttttttg	ttttgttttt	ttaaagaaga	aataaagacc	4440
45	caggggagaa	tgggtgttgt	atggggaggg	aagtgtgggg	ggtccctctc	caacccacac	4500
	tgttcacatt	gcaaatatat	tttggaaaaa				4530

<210> 11

<211> 2205

<212> DNA

<213> Homo sapiens

[illegible]

	cacagtcacc	ttggaggaca	tcaaggagca	cgtgaacacg	ctagagaaaag	cgggttcagg	420
	caaggagcgc	agattcgtgc	tgccggccct	gcggatgctg	ccttccacat	cacgcgcct	480
	caaccocat	gtttcgtata	aggctgtgca	gggctctctc	acttcaata	atgccactcg	540
5	agaacttttg	ctccocctcc	tggaagagcc	catggacaca	gaggctgatt	tacagttccg	600
	tcocccagcg	ggaanaagctg	cgtagacacc	cctcctgctc	gaagtggaa	cctatctcca	660
	actcctcgtg	gtcatcttca	tgatgaacag	caagcgctac	aaagaggcac	agaagatctc	720
	tgatgatctg	atgcagaaga	tcagtactca	gaaccgcg	gcccagacc	ttgtagccgc	780
	aaagtgttao	tattatcaac	cccggttcta	tgagttcctg	gacaagctgg	atgtgggtgg	840
	cagctctctg	catgctcggc	tcogggaacg	taogcttogg	catgaocgag	acgggcaggc	900
10	caocctgttg	aaactcctgc	tgccggaatta	cctacactac	agcttgttac	accaggctga	960
	gaagctgttg	tccaagctctg	tggttccaga	gcaggccaac	aaacaatgagt	gggcccaggt	1020
	cctctactac	acagggcgaa	tcaaaagcaat	ccagctggag	tactcagagg	cccggaagac	1080
	gatgaccaac	gccccttgca	aggccoccca	gcacacagct	gtcggcttca	aacagacggt	1140
	gcacagcttt	ctcatcgtgg	tggaagctgtt	gctgggggag	atccctgacc	ggctgcagtt	1200
15	ccgcacagctt	tcocccaagc	gctcactcat	gcccatttta	cttctgactc	aagctgtcac	1260
	gcacagagctt	ctagccaagt	tcaaccaggt	cctggatcag	tttggggaga	agtttcaagc	1320
	gcacaggaac	ctacagcaag	ttatccggct	gcggcacaac	gtgattaaga	caggtgttac	1380
	agatgggaac	ctagccactaa	cccggaatctc	cttggctgac	atcgccagga	agctgcagtt	1440
	catgatcagc	ctctcctatt	cccggaatctc	cttggctgac	atcgccagga	agctgcagtt	1500
20	ggatgacccc	gaagatgcag	agttcattgt	tgccaaagcc	gagatgatgt	gctgcatact	1560
	ggccagcact	aaaccaagga	agggctatgt	ccaatccaa	gagatgatgt	gctgcatact	1620
	caaccagagag	ccccaagctag	ccttccaccca	gcgcactctc	gagatgatgt	gctgcatact	1680
	catgtctgtc	aaggccatga	ggtttcctcc	caaatogtac	aaacaaggact	tggaagttgc	1740
	agaggaacgc	cgtgagcgag	aaacagcagga	cttggagttt	gccaaggaga	tggaagttgc	1800
	tgatgatgac	agcttccctt	gagctggggg	gctgggggag	ggtaggggga	atggggagac	1860
	gctctctctc	ccttggggct	ccctgcacca	gggcaactgt	ccacttttcc	caacaacagc	1920
25	tcatactgtc	cattcgtgca	gggggtgggg	gtgctgggag	ccagccaccc	tgacccctcc	1980
	cagggtccct	ccccaagctg	tgacttaact	tacagcagcc	aggaggggtg	gagggcaaac	2040
	tcocccggca	gggtcctggc	cagcagctgt	ggagcaggga	gggaaggata	gtctgtgtga	2100
	ctccttttag	gagtggggga	ctagaactgg	gatgtcttgg	cttgtatgtg	ttttgaagct	2160
	tgattatga	tttttaacaa	ataaaaagtt	ctcccaaaaa	aaaaaanaaa	aaaaaanaaa	2220
30	aaagcgccgc	cgaaattc					2280
	<210>	13					
	<211>	2960					
35	<212>	DNA					
	<213>	Homo sapiens					
40	<400>	13					
	ctgcagcttc	caggcgtcta	tcagcggctc	agccttctgt	cagctgttct	gttcaaacac	60
	ctcggggcca	ttcaggcctg	ggtagggcag	gggaggaag	ggagtttgag	gggggcaagg	120
	cgagctcaaa	ggagatcag	agattccaca	atttcacaaa	actttcgcaa	acagcttttt	180
45	gttccaccca	ccctgcattg	tcttggaac	caaatctgca	taaatcctgg	gaagttatta	240
	ctaaagcttc	tgctggcccc	caggtaattt	cctccagcta	ctccatgggg	ttatgtataa	300
	agggccccc	agagctgggc	cccaaaaagc	cccgagcctc	gcagccacgc	ccacccagga	360
	cccatgtcgt	gcactgcacc	ccagagcccc	atgaagctga	tggtgtgagt	tcttgcccca	420
	ggatgggaga	gcgcctgccc	ctggcatggg	agggaggctg	gtgtgcagga	ggggctgggg	480
	atcccccgtc	tggtgaatgg	gattaaaggg	acccagtgct	cccgagaggg	ctcaggtggg	540
50	taggggaacg	cagtlctctc	gagcccgctc	tgctccacgc	cctgcacgtg	ctgctgtggc	600
	acagtgcaat	ctgggacagt	tgaggaagct	cccccctggg	cctgcacgtg	ctgctgtggc	660
	agagcttctc	gtccaagctg	ttagagcaag	tgaggaagat	ccagggcgat	ggccagcgcc	720
	tcacaggaaa	gctgttgagt	gaggtgggtg	agagggctgc	ggagggagc	cgggtgggga	780
	gagctacagg	gcagggccaa	catcctctgg	agggggagac	aggggacat	ggaggaatatt	840
	aggagcagtg	gagctgggga	agcgtgggaa	gggagctggg	gagggagacc	ttgtgtggga	900
55	cagtgtctgg	gagggctggc	tggtgatggga	gtggagcact	cacactcagg	agaaagggca	960

	agggccocctg	tgagatcaga	gagtgggggt	gcagggcaga	gaggaaactga	acagocctggc	1020
	aggacatgga	gggaggggaa	agaccaagaga	gtcggggagg	accogggaag	gagcggcgac	1080
5	ccggccocagg	cgaggtctcac	toagca toct	tcacatccca	gtgtgccacc	tacaagctgt	1140
	gcacccocga	ggagctgggt	ctgtctggac	actctctggg	catccocctgg	gctccocctga	1200
	gcagctggcc	cagccaggcc	ctgcagctgg	tgagtgctcag	gaaaggataa	ggctaattag	1260
	gaggggggaag	gagaggagga	acacccatgg	gctcccccac	gtctccagggt	tccaagctgg	1320
	gggocctgaag	tatctcaggc	agcaccocct	aactctctccg	ctctgtctca	caggcaggct	1380
	gcttgagcca	actccatagc	ggccttttcc	tcacacaggg	gctcctgcag	gcccctggag	1440
10	ggatctccccc	cgagttgggt	cccaccttgg	acacactgca	gctggagctc	gcgcactttg	1500
	ccacccacat	ctggcagcag	gtgagccttg	ttgggcaggg	tgcccaagggt	cgctctggca	1560
	ttctggccac	cacagccggg	cctgtgtatg	ggccctgtcc	atgtgtgcag	ccccccagcat	1620
	ttctccatctt	gtaataacgc	ccactcagaa	gggcccaccc	actgatccca	gctttccccc	1680
	acagatggaa	gaactgggaa	tggccocctgc	oocgcagccc	accaggggtg	ccatgcocggc	1740
15	cttgcocctct	gctttccacg	gccgggcagg	aggggtcctg	gttgccctcc	atctgcagag	1800
	cttctcggag	gtgtctgtacc	gogttctacg	ccacocctgcc	cagccocctag	ccaagccctc	1860
	cccatcccat	gtattttatct	ctattttaata	tttatgtctc	tttaagctcc	atatttaaag	1920
	acaggggaaga	gcagaaocga	gccccaggcc	tcctgtctct	tcctgtcatt	tcctgagttc	1980
	attctctcgc	ctgtagcagt	gagaaaaagc	tcctgtctcc	ccatccocctg	gactggggagg	2040
20	tagataggta	aataccaaagt	atttattact	atgactgctc	ccagccocctg	gctctgcaat	2100
	gggcactggg	gtgagccocct	gtgagccocct	ggctcctgag	gtcccccact	ggggccocctg	2160
	agagtaacag	gtctcccccag	tgggagacaa	gaaatccctg	tttaatatctt	aaacagccagt	2220
	gttcccccac	tgggtccttg	cacccctcac	tcctggccca	gcgcactgca	cagcggccccc	2280
	tgcactccct	tgggtctgtag	gcccctggac	aagcagaggt	ggccagagct	ggggagccatg	2340
25	gcccctgggt	cccacgaatt	tgcctgggaa	tcctgttttt	cttcttaaga	cttttgggac	2400
	atgggttgac	cccacgaact	cacogacgtg	tcctgttttt	ttctgggtgg	cctcggggaca	2460
	cctgcocctg	cccacagagg	gtcaggactg	tgaactctttt	tagggccagg	caggtgocctg	2520
	gacatttgcc	ttgtctggatg	gggactgggg	atgtggggagg	gagcagacag	gagggaatcat	2580
	gtcaggctcg	tgtgtgaaag	gaagctccac	tgtcacccctc	caactcttca	ccccccactc	2640
	accagtgctc	cctccactgt	cacattgtaa	ctgaacttca	ggataataaa	gtgttttgct	2700
30	ccagtcacgt	cctctcctct	tcttgagtc	agctgggtcc	tgcccaagggt	ctggggagggt	2760
	ggctgaaggg	tgggagaggc	cagagggagg	tcggggaggga	ggtctggggga	ggagggtccag	2820
	ggaggaggag	gaaagtctct	aagttctctc	gacattccatt	cogtttagcac	atatattatc	2880
	gagcacctac	tcctgtgcaga	cgtcgggcta	agtgctgggg	acacagcagg	gaacaaggca	2940
	gacatggaat	ctgcacccga					2960
35	<210>	14					
	<211>	850					
	<212>	DNA					
40	<213>	Homo sapiens					
	<220>						
45	<221>	misc_feature					
	<222>	(3) .. (4)					
50	<223>	n=a, c, g or t					
55							

<220>

<221> misc_feature

<222> (9) .. (9)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (11) .. (11)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (18) .. (18)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (202) .. (202)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (205) .. (205)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (273) .. (273)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (327) .. (327)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (367) .. (367)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (581) .. (581)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (599) .. (599)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (628) .. (628)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (673) .. (673)

<223> n=a, c, g or t

<220>
<221> misc_feature
<222> (675) .. (675)
<223> n=a, c, g or t

<220>
<221> misc_feature
<222> (682) .. (682)
<223> n=a, c, g or t

<220>
<221> misc_feature
<222> (693) .. (693)
<223> n=a, c, g or t

<220>
<221> misc_feature
<222> (698) .. (698)
<223> n=a, c, g or t

<220>
<221> misc_feature
<222> (700) .. (700)
<223> n=a, c, g or t

<220>
<221> misc_feature
<222> (720) .. (720)
<223> n=a, c, g or t

<220>
<221> misc_feature
<222> (730)..(730)
<223> n=a, c, g or t

<220>
<221> misc_feature
<222> (734)..(734)
<223> n=a, c, g or t

<220>
<221> misc_feature
<222> (742)..(743)
<223> n=a, c, g or t

<220>
<221> misc_feature
<222> (746)..(746)
<223> n=a, c, g or t

<220>
<221> misc_feature
<222> (748)..(748)
<223> n=a, c, g or t

<220>
<221> misc_feature
<222> (752)..(752)
<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (762)..(762)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (767)..(767)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (777)..(777)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (783)..(784)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (789)..(789)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (794)..(794)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (797) .. (798)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (803) .. (805)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (810) .. (810)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (817) .. (817)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (826) .. (827)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (831) .. (832)

<223> n=a, c, g or t

```

<220>
<221> misc_feature
5 <222> (834)..(834)
<223> n=a, c, g or t

10 <220>
<221> misc_feature
<222> (837)..(838)
15 <223> n=a, c, g or t

<220>
20 <221> misc_feature
<222> (840)..(840)
<223> n=a, c, g or t

25 <220>
<221> misc_feature
30 <222> (844)..(844)
<223> n=a, c, g or t

35 <220>
<221> misc_feature
<222> (846)..(848)
40 <223> n=a, c, g or t

<400> 14
45 tttnctttnt ngccatgncc agttcaactc agcctctcag ttccacacgg acaacatgag 60
ggacccctctg aaccgagtcg tggcccaacct gtctctgctc atctctctcca tctctggggtc 120
tcgcaacocgt ggccccccaca cccagtttgt gcagttggtc atggaggagtg gtgtgggactg 180
cctggagcag ggtggcogtg gnagngtctc gcagttcatg cccttcacca cagtgtogga 240
actggtgaag gtgtcagcca canggtggtt ctggccatca cggacocag 300
cctgcocctg ggccgcacag tggctgntaa agccattgct gcactctgag gggcctggca 360
tggccgnagt gggggctggg gactggcgca cactgactca gcccacaggc octccaaggg aagcagtgag 420
gaaagatgag gcatctgtgc toacatcoct tocacatggt goaagagcct ctagoggctt 480
ccagttccccc gctcctgact cctgactcca ggaagtctcc cgtttcttc ttttcaaaat 540
tttctctccc atcttctggt caactgagga gagtggacag nctggaccac aagccacng 600
ggtcacocct gtgttgccc cgcocagncc agggagtagt ttacotcttg aggaactttc 660
ttggatggaa agnnggtttt tntgtgtgt gtntgtgnan gtgtttttog ggggtttttt 720
55 gggcaatatn ttangggaa cnncoctncc cncatttttt cnttagagct ccccgngnga 780

```

	aanntcttna	tcenctnnct	ttnnnctecn	tcacctnct	tcttntntct	nnntntnnn	840
	tcenennnc						850
5	<210>	15					
	<211>	2309					
	<212>	DNA					
10	<213>	Homo sapiens					
15	<400>	15					
	ccocgggggc	aggaggcggg	gggcccggcc	ccacccggcc	cccatggaag	cccccagcac	60
	ggggcgctga	gaccccccgg	tcgctgcccc	gcccggtcgg	gogcgccacg	ccagggatct	120
	ctgggacagg	caagactcgg	aagctactcc	ccagccacac	agcccgggac	ccacaaaacc	180
	agcttgcccc	cagcctccgc	acctgcacac	acctggcccc	ccccaccgcc	cgccccccct	240
20	ggggcgccgg	gcattggtgt	aaaggccaa	tgctgaggcg	ggtatcatgg	gtgctgtgac	300
	ctaggctcgg	gggtggcagg	gggtgggtgg	ctgtgggtgt	gcgggggggg	ccagtggtgc	360
	cacccacgtc	tcttgccgtg	ctggagggca	tcttggtatg	aattggaagt	aatggaacag	420
	aagcccaaga	aggtggagtg	tggtgcagac	ccagaggaga	acagtgccag	gtccaccagat	480
	ggaaagcgaa	aaagaaagaa	oggcccaatg	tcctgaaaaa	ccagcatgtc	aggggtatct	540
	actatgcctc	gtatcaactt	cgagcagtg	gtcgtgtgtg	gggacaaagg	aactcggtat	600
25	aacctccact	ccacactatc	ctgcaaatat	aagggtctct	ttcgccggac	aatccagaag	660
	cgcaatcagt	gcccagctgt	ccgctccaag	aagtgcctcg	ccgtgggcat	ggccatggac	720
	ttggttttag	atgactgtga	gcgggtggcc	aagcgttaag	tgattgagca	gaaccgggag	780
	cgggcgggga	aggaggagat	gatccagatc	ctgcagcagg	gacccagacc	caactcgtga	840
30	gagtgggatc	tgatccacat	tgccacagag	gcccacatgc	gacccagacc	caactcgtga	900
	cattggaaac	agaggcgga	atctcctgcc	gatgacattg	gacccagacc	caactcgtga	960
	atgcggagcg	gagacaagg	ggacatttgc	gcccacatgc	gacccagacc	caactcgtga	1020
	ccggccatca	cccggtgtgt	octgaagggg	gcccacatgc	gacccagacc	caactcgtga	1080
	tgccgaagac	agatcatcct	agagcgacac	tgctgcctgg	agatcatgtc	ccgtcgggcg	1140
35	gctgtccgct	acgaacctga	cgccctgggc	ctgacgctga	gtggggagat	ggctgtccag	1200
	aggtagcagg	tcagaaatgg	ccgctcgggc	gtagctcccg	agccacatct	tgaaactggg	1260
	aagtcactct	ctgcctttaa	ccgtggtgac	acgggaagtg	ctctgctgca	ggctgtgtgt	1320
	ctaatgtcaa	cagacccgtc	ggccctgctg	tgctgtgaca	agatccagaa	gagtcaggag	1380
	gcgtacctgc	tgccgttcca	gcactacgtc	aaccacccga	aacacacat	tcgcactctc	1440
	tgccccaagg	tgctgtgtaa	ggagagagaa	gtgcagagtt	cgattctgtg	caagggggca	1500
40	ggcgccgaag	gcggcgccgg	gggttcactg	ggcgtccacc	cggaagagca	ccagcttctc	1560
	ggaaatgcat	ttgttccagg	tcgcagagtg	ccgcagcttg	agcagcagct	tggtgaaagg	1620
	ggaggtctcc	aaaggccggg	tcttcaggca	ccagcccgca	agcagcagct	tggtgaaagg	1680
	cgtagctgct	tcacccagag	gggaattctc	ccagcccgca	agcagcagct	tggtgaaagg	1740
	agcagtgagg	gggaactccgc	gagctctctc	ccagcccgca	agcagcagct	tggtgaaagg	1800
45	gcaggcgaat	cagcctctcc	ctgaaagccc	ccagcccgca	agcagcagct	tggtgaaagg	1860
	gcacatacct	ctccacaggc	ctgtccccaa	gagcagggag	tgccatgaag	ctgggagcgt	1920
	gggtctgaca	gggtgtgtca	ccctccatcc	cgtaagagca	ccctccatcc	ctgagcagct	1980
	caaacatgtg	cagactccct	tgcttttttg	tgctgtagtc	ccctccatcc	ggatgcctct	2040
	ccccctttct	ctgcctggca	acatcttaac	tgctgtagtc	ggcccccacg	caagtgtaac	2100
	ctcctccccc	agctccccc	ggcagaaata	gttgctgtgt	cttcccttgt	tcactgtctc	2160
50	actgtgacac	ttatctcact	gttttataat	tagtcgggca	tgactctggt	tcocacagca	2220
	gactgtgtct	gaatcatgtc	tgatcccg				2280
							2309

<210> 16

<211> 2355

<212> DNA

<213> Homo sapiens

<400> 16

ccgttgccctc	aacgtccaac	cctttctgag	ggctgcagtc	cggtccacccc	aagacctgct	60
tgcaaggtgc	ttcggtacct	gatgtgagtc	cggtgggtcc	actcccccgc	cttagccagt	120
ccccaggggg	caacagcggc	gatcgcaaac	tttagtttga	gtcaagggtc	agtttgaaatg	180
acgcctctca	gctgggtgaag	acatgaacca	cctggaactc	aacaacacaa	caggtggggt	240
atcacacctac	attggctcca	gtggctcctc	ccccagccgc	accagccctg	aatccctcta	300
tagtgacaac	tccaatggca	gcttccagtc	cctgaaccaa	ggctgtccca	cctacttccc	360
acaccccccc	actggctccc	tcaacccaaga	cccggtctgc	tcctttggga	gcattccacc	420
cagcctgagt	gatgaaggct	ccctctcttc	ctcaattccc	tgctgttcac	cctccctccc	480
cttctataat	gggagccccc	ctgggagctc	acaagtggcc	atggaggaca	gcagccagat	540
gtccccccag	aagagaccca	gcaacatcac	caagctgaat	ggcattggtg	tactgtgtaa	600
agtgctgtgg	gacgttgctc	cggtcttcca	ctacggtgtg	ctcgccctgc	agggctgcac	660
gggtcttttc	ogtggagcca	tccagcagaa	catccagtac	aaaaggtgtc	tgaagaatga	720
gaattgtccc	atgtctcgca	tcaatcgcaa	ccgtgtccag	caatgtcgct	tcaagaagtg	780
ttctctctgt	ggcatgtctc	gagacgctgt	ggcttttggg	cgcatcccca	aacgagagaa	840
gcagcggatg	cttgctgaga	tgcaagagtc	catgaacctg	gccaacaccc	agttgagcag	900
ccagtcgccc	cttgagagctt	caacccacca	gcaacccacc	ccaggccccc	tggtgcccctc	960
gccaccccc	gctccgggtcc	ccctaccccc	gggtgggctc	tcacagtttc	cacaacagct	1020
cagccctccc	agatccccaa	gcccctgagc	cacagtgagg	gatgtgatat	cccaggtggc	1080
ccggggcccat	agagagatct	tcacctacgc	ccatgacacg	ctgggagctg	cacctggcaa	1140
cttcaatggc	aaacatgcct	caggtagccc	tcacgccaac	accccacatc	gctgggaaaa	1200
tcaggggtgc	ccacctgcgc	ccaatgacaa	caacacccct	gctgcccagc	gtcataaoga	1260
ggccctaaat	ggctctgccc	aggtccctcc	ctcctacccc	cccaactggc	ctcctggccc	1320
tgccacccac	agctgccacc	agtcaccaag	caacggggcc	gcttatggcc	ccacccagct	1380
gtatgcagcc	ccagaaggca	agggacctgc	caacagtcac	cggtggggag	tggttagagt	1440
tggtctgctg	gcattgtcct	tgaaacatgt	cccgcatgga	cgcaagtgcc	gaaoggtgca	1500
ggagatctgg	gaggattctc	ccatgagctt	ccagcccgct	gtgcccagct	ccctgcttaa	1560
tgccaaacac	atcccgggct	tcogtgacct	ttctcagcat	gaccaaagtc	ccctgcttaa	1620
ggctggccac	tttgagggtg	tgatggtgag	ctttgctctg	ttgttcaagc	tgaaggacaa	1680
cagactgatg	tttcaatagc	ggacccaact	cagcctgcag	gagcttggtg	ccatggggatc	1740
gggagacctg	ctcagtgcca	tgcttcagct	cagcagagag	ctcaactccc	tgccgcttac	1800
cgagggaggag	ctggccctct	tcacccgggt	ggctgtctgc	ctgcagagac	gctcggggat	1860
ggagaaatcc	gcttgggtgg	agcagctcca	ggagacgctg	ctcgggctct	ctcgggctct	1920
gggtgtgaag	aaacggccct	tggaagcttc	ccgcttcacc	aaagctgtgc	tcgaagctgc	1980
ggacgtggcg	acccctgaaca	acatgcattc	cggaagctgc	ctgtctctcc	gggtggaagc	2040
ccagtgaccc	gcccggccgg	cctttctgcg	ctgcacccct	gtacagaaat	gaactctgca	2100
cttctctctc	cttcaagaga	cgaaaaagaa	aagcaaacca	gaactctatt	tatttggtta	2160
taaaatattc	caagatgagc	ctctggcccc	ctgagccctc	ttgtataaac	ctgcccctcc	2220
cccccatcac	cgaaactccc	ctctccctcc	atttaaacca	ctctgtctcc	cccccaaccc	2280
tcacctggcc	ctctgatttg	ttctgttctc	gtctcaaatc	caatagttca	cagctaaaaa	2340
aaaaaaaaaa	aaaaa					2355

<210> 17

<211> 4119

<212> DNA

<213> Homo sapiens

<400> 17

gaattcogtt	gctgtogcag	acacacacac	acacacacac	acacccccac	acacacacac	60
acacccccac	acacacacac	acacacacac	acacacacac	acacacacac	acacagcggg	120
atggcogcag	gcgcgacogc	tagcacgcgc	ggactagcta	tcacgootcc	cagcagootc	180
tgogcggggc	goggttgcta	agtaoctcgc	oggttggtgc	ogttctcogt	aagatggcgg	240
acogggcgcc	gcagcgcgct	tcgcaagaca	cogaggacga	ggaatctggt	gcttcgggct	300
cgcagacggg	cggtctcccg	ttgcggggag	cggggagctg	cagcggtagc	gcgcggaggc	360
gcgcgacggg	ctctctgctc	tcacagcgcg	gaggcgcgaac	cggggocctt	catctcggcg	420
gggttgagag	ogggggcgcc	aagagtgcgt	aggagtgcga	gtgtgagagt	gaagatggca	480
ttgaagtgta	tgctgtcttc	toggatttatg	aaagtgcaga	agactcggaa	gggtgaagaa	540
tgtaatacga	tgaaaggagaa	aactccaaag	tgagctgtaa	atcagaagct	aatgatcgtc	600
ttaattcttc	aacaaaagaa	gagaagggag	aagaaaagcc	tgacacccaa	agcactgtga	660
ctggagagag	gcaaaatggg	gaogggacag	agagcacaag	gcctgtggag	aacaaaatgg	720
gtaaaaagg	ccctaagcat	ttggatgatg	atgaagatcg	gaagaatcca	gcatacatat	780
ctoggaaagg	gctctctctt	gagcatgctc	ttcaggggca	aactcaggag	gaggaaagtc	840
gaoccaaagg	gcgtcagcga	aagctatgga	aggatggagg	tcgctggggg	catgacaagt	900
tcogggaaga	tgagcaggcc	ccaaagtccc	gacaggagct	catgtctctt	tatgtttatg	960
acattcgtct	ccgtcctaag	ccctga tgaca	tcacaaactc	aagaatcccg	aaaccccgat	1020
atggggatcc	tcacacaaag	gatccaaaat	ggaaoggtga	goggcctaaac	aagtctcctc	1080
gcacacaaagg	tccttggggg	acccctacac	caaggacatt	tattaaacag	aatgctcgag	1140
gtacccggcg	tatgtctgca	ccacaggaa	atctctgcac	tggggggctc	aaggaaagtg	1200
gtgctgtgtt	taggcctgtg	gaagctgggt	ggcagcatgg	tgccgggtct	ggtgagactg	1260
ttaaagcatga	gattagttac	cggtccacgc	gcctagagca	gaactctgtg	agggatccat	1320
ctccagaagc	agatgtctca	gtgcttgga	gtcctgagaa	ggaaagggca	gcctcagaga	1380
caccagctgc	tgctcctgat	gctgcaccac	cacccocctg	tagggccatt	gagaaagaa	1440
ccattctccg	ggcagaagaa	actcgaaaca	aagtgtggga	tgcaatcaag	cttgcaagag	1500
agggtgcccc	tcctcctgaa	ggactgatcc	cagcaactcc	agtcocagaa	acacccccaa	1560
ctccacactac	taagactggg	acctggggag	ctccggtgga	ttctagtaca	agtggaactg	1620
agcaagatgt	ggcacaacta	aataagcag	aacagaattg	gagtcggggg	cagcctctct	1680
tcctgcaccc	acgggaactt	cgagggtatg	ccacacatat	acacatggga	gcaggacctc	1740
cacctcagtt	taacoggatg	gaagaaatgg	gtgtccaggg	tggtcgagac	aaacgcctat	1800
catccacagc	gcacaaagct	gtgcacagc	cccccgcgcc	tcacgtgcac	atcagtatca	1860
tgaggaggaca	ttacttatgat	ccactgcagt	tcacagggac	aatctatacc	catgttgaca	1920
gcocctgcgcc	gtctgcctcca	cagggcctgc	ttgtgcagcc	aggaaatgaa	cttcccaccc	1980
cagggtttaca	tcocccacag	acaccagctc	ctctgcocaa	tcacaggoot	tatccccccc	2040
cagttgcatt	gtctcagga	cagccacac	ctcagcagtt	gcttgctctc	ctactctttt	2100
ctgctccagg	ggctcatagg	tttggttaac	ccagttaccc	ttatgctcca	ggggcactgt	2160
ctcccccacc	acogcctcca	ctgta tccca	atacacaggc	ccactcacag	gtatatggag	2220
gagtgaccta	ctataacccc	gccagcagc	agggtcacgc	aaagocctcc	ccaccccctc	2280
ggactccocca	gcacgtccac	atccagcccc	ctccacacga	ggtctgtaag	aggggttcca	2340
gttaatacaa	gtttctgaa	atttttaatt	ttaacatcat	ataaaaagca	gcagagggtga	2400
gaaotcagaa	gagaaataca	gctggctatc	tactacagga	aggggttcaa	agatataggg	2460
ttgggtctct	acacgcaaac	agctgaagaa	ggaggacccc	tgcccttcct	tgaggacagg	2520
ctctagagag	agggagaaac	aagtggacat	ogtcccatct	tcactcttca	cttgatgttg	2580
ctgtgtctgg	gggagcagag	agagccagac	agocccaaag	ttctgagctt	agatacagaa	2640
gcocactgct	tcctgtgttc	ttcactctct	ggaaattgaa	gtgtctctct	ttcccaaggaa	2700
agctccttcc	tggtttgttt	gttttttaag	atgttcaatt	ttaaagcctt	gctctctatc	2760
cttaataatta	tttttaattt	ttctctttgt	ttctgtttct	tgctctctct	ccctgccttt	2820
aaatgaacaa	agttcagttc	ttctgttttc	tagccctcat	ggattccctt	ctctctcttc	2880

	ogtgoatccc	agataatgga	gaatgtatca	gccagoccttc	ccacccaagt	ctaaaaagac	2940
	ctggoccttc	acttttagtt	ggcatttgtt	atcctcttgt	atacttgtat	tcocottaact	3000
	ctaaccctgt	ggaagcatgg	ctgtctgcac	agaggggtccc	attgtgga	aaagctocaga	3060
5	tacggttgat	aggagcocc	ctctttgact	taggttttta	ggagctctgag	catocataca	3120
	gatggttgat	atgatgggct	ctgtctctct	gctgaggccc	aatccocctac	tgtgtgggaga	3180
	gatggccac	cagatgcttt	tgtgagaaag	ggatgtgtga	gtgagagoc	tgtoccttag	3240
	gggtgtgtat	tcacatagtc	ctcaggggctc	agtcttttga	ggttaagtga	attagagggc	3300
	cttgctctct	tctcttccat	tctctctgct	acaccccttc	tcocagttgt	gtggaccac	3360
10	gcactctctt	aaaggccaaat	attatocagc	aagcagctta	ccctgtcctt	tgcactgtct	3420
	ctctctccag	tctctctgct	tacaaagtgt	ttagatgtta	ctacccattt	tccocogaat	3480
	tctatttttg	tcctgtcaga	cagaataata	aaactccctg	gcttaaggcc	taaggaaagcc	3540
	agtoacocct	tgggcaagg	ctcctatctt	tccctccatc	ccatggcaact	aaacccactc	3600
	ctctgctgct	ctgtggaaga	gattcctatt	actgcaagtac	atacgtctgc	caggggtaac	3660
	ctggccactg	tcocctgtct	tctacagaac	ctgaggggcaa	agatgtgtgc	tgtgtctctc	3720
15	cccggttaag	tcactgtttt	tattcctctc	atctagcagc	tggcctaact	actctgagtc	3780
	acaggtgtgt	gatgtgagag	ggggagaggc	acttaactgt	taacccocaa	ggaggaatac	3840
	actaagagat	tctctagagg	gtagctgggt	gttgtgctct	tgttaggtct	tccoccttgc	3900
	cttaaacctg	aagatgtctc	ctcaagccctg	tgggcaagcat	gcccagatct	ccagcaoccta	3960
	agacacctgt	agagctgtct	ctgtgtgtct	actgtgttta	gttgcaagg	tttctccatg	4020
20	tgtggtgtgt	ttttttgtta	ctgttttaaa	gggtgcccac	tgtgtgacg	catgtgact	4080
	tggagataat	aaaatttaga	ctataaacct	gaaaaaaa			4119

<210> 18

<211> 2653

<212> DNA

<213> Homo sapiens

<400> 18

	gagcgggct	ggagtttgtct	gctgcogctg	tgcagtttgt	tcagggggct	gtggcgggtga	60
	gtccagagag	ctgogtgtga	gagaogtgag	aaggatccgt	caactgaggag	gtggaagaa	120
	gaggattgct	cgaggaggcc	tgggggtctgt	gagacagogg	agctgggtga	aggctgcggg	180
35	ttccggcgag	gootgagctg	tgtgtgtctc	atgcctccaa	cccgatccca	ggcaccggct	240
	acaaactagt	ttccaaaaag	gaagctgtct	ogggcattga	acaaagctaa	aaactccagt	300
	gatgccaaac	tagaaccac	aaatgtccaa	acogtaacct	gttctcctcg	tgtaaaaagcc	360
	ctgcctctca	gcccacggaa	acgtctgggc	gatgacaccc	tatgcaaac	tcocccattta	420
	oocctctgtt	ctccacccaa	gcaaggccag	aaagagaatg	gtcccccctca	ctcacatata	480
40	cttaaggggac	gaagatttgt	atttgacaat	cagctgacaa	ttaagtctcc	tagcaaaaaga	540
	gaactagccca	aagtctccca	aaacaaaata	ctttctctcag	ttagaaaaag	tcaggagatc	600
	acacaaaatt	ctgagcagag	atgtccactg	aagaaagaat	ctgactgtgt	gagactatct	660
	aagcaagaag	gcactgtcta	ccagccagcca	aagctggccc	tgaacacagc	tgtccagagc	720
	cggtctgctg	ccaggggaaag	ggagatggat	gtccatcagg	attctttagg	ggaaacacac	780
45	tgtgggaaaa	aagctgggaag	ccctttacctt	tctgggtgct	ctggaaactgg	aaaaactgoc	840
	tgtcttaagcc	ggattctgca	agaccccaag	aaggaaactga	agggctttta	ccactctgat	900
	ctgaacttgc	tgtccttagg	gactgcccag	gctgtattcc	cagctattgc	tcaggagatt	960
	tgtcagggaag	aggtatccag	gccagctggg	aaggacatga	tgaggaaatt	ggaaaaaacat	1020
	atgactgcag	agaaggggccc	catgatgtgt	tgtgtattgg	acgagatgga	tcacactggac	1080
	agcaaaaggcc	aggatgtatt	gtacacogcta	tttgaattgc	actaggttaag	caattctccac	1140
50	tgtgtgtctga	tgtgtattgc	taataccctg	gatctccacg	ttagaattct	actatggctct	1200
	caagcttagag	aaaaatgtaa	gcccacagctg	ttgaacttcc	caacttatcc	cagaaataacg	1260
	atgtcaacta	ttttgcaga	togaacttaat	caggtatctc	gagatccagc	tcctggacaat	1320
	gctgcagttc	aattctgtgc	cogcaaaagtc	tctgctgttt	caggagatgt	tcgcacaaagc	1380
	ctggatgtgtt	cgaggagagc	tattgaaatt	ttagagtcag	atgtccaaaag	ccagactatt	1440
55	ctcaaacacc	tgtctgaagt	taaatccact	tctgagccct	tgtatcccaa	gagggttggt	1500
	cttattccaa	tatcccaagt	catctcagaa	gttgatgtga	acaggatgac	cttgagccaa	1560

	gagggagcagc	aagattcoctt	ccctcttcag	cagaagatct	tggtttgtct	tttgatgtct	1620
	ttgatcaggc	agttgaaat	caaagaggtc	actctgggga	agttatatga	agcctacagt	1680
	aaagctctgc	gcaaacagca	ggtggcggtc	gtggaccagt	cagagtgttt	gtcactttca	1740
5	gggctcttgg	aagcagggg	caatttagga	ttaaagagaa	acaaggaaac	ccgtttgaca	1800
	aaggtgtttt	tcaagattga	agagaaagaa	atagaacatg	ctctgaagaa	taaaagttta	1860
	attggaata	tcttagctac	tggattgcoct	taaattctto	tottacaacc	caacogaaag	1920
	tattcagctg	gcatttagag	agctacagtc	ttoatttttag	tgctttacac	attcggcgct	1980
	gaatacaaat	atgaactttt	ttacttgaag	ccaatgaatt	ttactctata	gattctttaa	2040
10	tattagcaca	gaataatato	tttgggtott	actatttttta	ccactataag	tgaccaggta	2100
	gaacottttt	aattacaatto	actacttota	ccaacttgtt	aatctatgoc	aatctgtctg	2160
	caagttgtaca	gtctgtgtga	gaggaaatgt	tgatatattta	ccctctogtt	tgctcaaaac	2220
	tgagtgggtt	tttttttgtt	tgtttttttt	gttgttgttg	tttttgaggc	gcgtctaac	2280
	ctgttgccca	ggctggagtg	caaaggcgog	ttctctgtct	actacagcac	ccgcttccca	2340
15	ggttgaagtg	attctcttgc	ctcagcctcc	cgagtatgtc	ggattacagg	tgcccaacc	2400
	cgcccccagc	taattttttt	atttttagta	gagacaggg	tttaccatgt	tggccaggct	2460
	ggtcttgaac	tcttgaccc	caagtgatct	gcccaccttg	gcctccctaa	tgtctgggag	2520
	tatagtgatg	agcccaaatg	ctcagccatt	aaggtatttt	gttaagaact	ttaaagttat	2580
	ggttaagaaga	atgaataatg	tccagaaaaa	tgaacgaag	tccacatgga	gattttggag	2640
	acactggtta	aag					2653
20	<210> 19						
	<211> 2907						
	<212> DNA						
25	<213> Homo sapiens						
30	<400> 19						
	gcatctctggg	cccaggcccc	atgccccgag	gaggggtggg	ctgaagccca	ccagagcccc	60
	ctgcagcagct	gtctgtctcc	ctctgtactg	tgcccgcttg	gcattggccag	caacagcagc	120
	tcctgcgccga	ccactggccg	cgggcaacct	aatgggtacc	ccgttgccctc	ctaagccttc	180
	ttcttccccc	ctatgtctgg	tggaactctc	ccgcagcgog	ctctgaccac	tctccagcac	240
35	cagotctccag	ttagtggata	tagcacacaa	tcocccagcca	ccattgagac	ccagagcaag	300
	agttctgaag	agatagtcc	cagccctccc	tcgccacccc	ctctaccccg	catctacaag	360
	octtgctttg	tctgtcagga	caagtctctc	ggctaccact	atggggctag	cgctgttag	420
	ggctgcaagg	gcttctctcc	cgcagcctc	cagaagaaca	tggtgtacac	gtgtccacgg	480
	gacaaagaact	gcattcatca	caaaggtagc	cggaacgcct	gcagtagctg	ccagctgcag	540
	aagtgctttg	aagtgggcat	gtccaaaggag	tctgtagaaa	acgcacgaaa	caagaagaag	600
40	aaggaggtgc	ccaaagccga	gtgctctgag	agctacacgc	tgacgcggga	gggtggggag	660
	ctcattgaga	aggtgcgcga	agcgccaccg	gaaacctctc	ctgcctctct	ccagctgggg	720
	aaatacacta	cgaaacaaac	ctcagaaaca	ctgtctctct	tggaatttga	cctctgggac	780
	aagttcagtg	aaactctcac	caagtgcatc	attaagactg	tggaatttgc	caagcagctg	840
	ccgggcttca	ccacccctcc	catcgccgac	cagatcaccc	tctcctcagg	tgccctgctg	900
45	gacatctctg	tctctgggag	ctgcagccgg	tacacgcccc	agcagggcac	catgaccttc	960
	tgccagccgc	tgacccctgaa	cgggaccocag	atgcaacaag	ctggctctgg	cccctccacc	1020
	gaacctgtctg	ttgcctctgc	caacccagctg	ctgcocctgg	agatggatga	tgccggagag	1080
	gggtctgctca	ggccatctctg	ccctcatctg	ggagacgcgc	aggaactgga	gcagccggag	1140
	cggttgagaca	tgctgcagga	gcgcctctgc	atgtcaatga	agattactga	cctctgggac	1200
	agggccagcc	gcccccaatg	gtctcccaag	acgtgtgata	tggaatctcc	gggtctccatg	1260
50	atcagcgcca	agggcgctga	gcgggtgata	tcagagggccc	tggaatctcc	gggtctccatg	1320
	ccgcctctca	tcagggaaat	gttgaggagac	ctggcccccc	ccgcagctct	gggtctccatg	1380
	ccgggggggtg	ggggcgggga	cggggggtgg	ccggcccccc	ccgcagctct	gggtctccatg	1440
	agcctcagcc	ccagctccaa	cagaagcagc	ccggcccccc	ccgcagctct	gggtctccatg	1500
	ccacatggac	ccagcctctg	ccctccgccc	ccggctttct	ccgcagctct	gggtctccatg	1560
	tgaccccgca	ccagcctctg	ccctccgccc	ccggctttct	ccgcagctct	gggtctccatg	1620
55	ggggacgggg	agggagggag	cagcgactcc	ttggacagag	ccctggggcc	ccgtgtggact	1680


```

5      gctgctccc acagcctggg ctgacgtcag aggcogaggc caggaaactga gtgaggcccc 1740
      tggctctggg tctcaggatg ggtcctgggg gctcgtgttt catcaagaca cccctctgcc 1800
      cagctccaca catcttctac accagcaaac gccaggactt ggctccccc tctctcagaac 1860
      tcacaaggca ttgctcccca gctgggggaa cctcaacctcc cccctgccc gggtgggtgac 1920
      agagggggtg ggcacggggg ggggggttcc cctgtatcat acocctgocat accaaccocca 1980
      ggttattaatt ctgctgtggt ttgtttttat ttttaatttt ttgttttgat ttttttaata 2040
      agaattttca ttttaagcac atttatactg aaggaatttg tgctgtgtat tggggggagc 2100
      tggatccaga gctggagggg gtgggtccgg gggaggaggt ggctoggaag gggocccac 2160
      tctcctttca tgtccctgtg ccccccagtt ctctctcca gocttttct cctcagtttt 2220
      ctcttataaa ctgtgaagta ctaactttcc aaggcctgcc ttccctccc tccactgga 2280
      gaagccgcca gcccctttct cctctgccc gaccactggg tgtggaoggt gtggggcagc 2340
      cctgaaggga caggctcctg gccctggcac ttgcttgac ccacactgag catggagca 2400
      gggcagagca agggcccccg gacagagttt tccagacct ggctcctgg cagagctgcc 2460
      tccctgcagg gcccaatcca totaggctcc ccagccccc ctgtgaaggg gctggccagg 2520
      ggcccgagct gcccocaccc cgggctcag ccaccagcac ccccaatagg ccccccagca 2580
      ccacacacat gctgctgccc acacacacaa acacacacac actggacagt agatggggcc 2640
      acacacactt ggcocaggtt cctccatttc cctggcctgc ccccccaccc caacctgtcc 2700
      cccccctgt cccctccttt accccgcagg accgggctac aggggggtct cccctcacc 2760
      ctgcaccccc agctggggga gtggctctg ccccgacct cttoaccag gggtggggcc 2820
      ccttccctgt aagccctgt gtgcacctg tactgttggt ctttccactg agatctactg 2880
      gataaagaat aaagtctct tttttct 2907

<210> 20

<211> 2096

25 <212> DNA

<213> Homo sapiens

30 <220>

<221> misc_feature

35 <222> (23)..(23)

<223> n=a, c, g or t

40 <220>

<221> misc_feature

<222> (27)..(27)

45 <223> n=a, c, g or t

<220>

50 <221> misc_feature

<222> (80)..(80)

<223> n=a, c, g or t

55
```

<220>

<221> misc_feature

<222> (120)..(120)

<223> n=a, c, g or t

<400> 20

agatgtttaa	aaataactttg	atnctcngtt	tccacctctc	ttaaattgtc	tttccctatg	60
ttaaatatac	agtcatacnc	ttgctgaaaa	aagttogcaa	tgagaacaat	catctaaaaa	120
tggtgttaac	taggtcaggc	gcggttgctc	atgcoctgtaa	tcccaccact	ttggggaggcc	180
gaggcaattg	gatcaactga	ggtcaggatt	ttgagaccag	cttgaccaac	atgggtggaat	240
cccatctctc	ctaaaaatcc	aaaaaattag	cggggtgtgg	tggcacaccc	ctgttaatccc	300
acctaactcg	gaggctgagg	caggaaaatc	cottgaaccc	aggaggcaaa	ggttgcatctg	360
acgcgaata	acaccactgc	actccagcct	ggacgataga	gtgagacccc	atctcaaaaa	420
aaagacagct	gtgacaaatg	cctgtattga	attgcaaggto	agtcttccac	ctccactaac	480
gtgtgcacaaa	aaagggctgc	ccccaaaagg	actaaaaggg	atccagcttt	gaattctgggt	540
gtctctcctc	agcctgtacc	tgccaaaacc	aagaatgcgc	gcacaaaggaa	gccatccact	600
ctctgatgatt	ctgactctaa	ttttgaaaaa	attgttttga	aagcagtcac	aagcaagggtg	660
agtggttgatc	ctagtcaatg	cttttgctgt	agatgtttctg	aaacacgttaa	ctaagccatt	720
gtctctctaaa	atttgccata	ctcttaagaa	aattaactct	catattctgt	tagcttttac	780
gttaacattt	tagtttttaac	aaagtttaaat	atgccaactta	tttgcccaat	ggaagagttg	840
gccttagatc	tgctctctat	tacttggttag	aaaatagaaa	actccttgaa	tatagtgtct	900
tgatacattt	ttttacatta	caattatggt	gtcagattta	caatgttgcaa	gttaactggg	960
ctttctctct	ttagaaatcc	aaagggggaga	gtgatgaact	ccatgtggac	tttgactcaag	1020
ctgtggtctc	togggcaaaa	ctctgtacgg	caagaaaacc	tataaggtac	ctgggaagagt	1080
cagatgaaga	tgatctgttt	taaaatgtga	ggcgattatt	ttaagttaatt	atcttaccac	1140
gccccagact	ggtttttaag	ttaactgaag	ctcttaactt	ctcccccctc	gaatttagtt	1200
tggggaagggt	gttttttagta	caagacatca	aagtgaagta	aagccccagt	gtcttttagc	1260
tttttataat	actgtataaa	tagtgaccat	ctcatgggca	ttgttttctt	ctctgtcttg	1320
ctctgtgttt	gagctgtctt	cttttgctt	taaaacctga	ttttttaagtt	ctctggaact	1380
gtgaaaatag	ctatctgato	acttcacgct	aaagcagtg	gtttatttaac	catccactaa	1440
gctaaaaacta	gagcagtttg	atttaaaagt	gtcaactctc	ctccttttct	aacttcaagta	1500
gatgtgagat	agagcaataat	tatctgtttt	atcttagttt	tatacataat	ttaccactcaag	1560
atagaaacttt	atggtttctag	tacagatact	ctactacact	cagcctcttta	tggtcccaagt	1620
ttttcttttaa	gcaattgaaa	attgctcatg	ttcttcatct	ctctaaatca	tcagaggccg	1680
aaagaaaaaca	ctttggctgt	gtctataact	tgacacagtc	aatagaatga	agaaaattag	1740
agtagttattg	tgatttatct	agctcttgac	ctgtccctct	tggtgtgctc	tgagcttgaa	1800
ctctcccaact	agagaaaacca	attctotaaga	ggactgtggt	gcagaaagact	cggggacacac	1860
atttgatcca	agattcttaaa	tgttatatgt	ataacatgc	tcagcaatga	ctgatttagat	1920
tcattttggg	aaatctccat	aatttcaatt	tgtaaacctt	gttaagacct	gtctacattg	1980
ttatatgtgt	gtgactgtgag	taagtgttat	aacgtttttg	taaatattta	ctatgttttt	2040
ctattagcta	aattccaaca	attttgtact	tttaataaat	gtcttaaaaa	ttgaaa	2096

<210> 21

<211> 2160

<212> DNA

<213> Homo sapiens

<400> 21
 agacgccttcg cccctgcgcgc ccccccgcgc ctgccttgggc cgggcgcgagg atgcgcgcga 60
 ggcgcctcggc ggccaggcttc cgcctccctcc ggccagccttc ctacactccg ccgcctacgtc 120
 ccgccttcggc ccgcgcgcgcg cgcctctatc ccgcgcgcctc cgggtccgggc tcctccaggac 180
 cggccaggccc gtgcgcgcgct gtgcctcccg ccgcctgcgc cgcgcgcgcgc cgctcccccgc 240
 ctgcgcgcag ccccccgcgc cgcctgcctgc ggccctgcgc gcttgggcga cgaagccatc 300
 ctgttgggcg cactgcctgc cctgcctcga cctgcctcgc cgcgcctcgc gcccccgcgt gggtcgcgag 420
 cactgcgcgc gaggaggcgg tgcacacccc ccgttgccgc ggccagggcg ccttgggcga tgcgcgggg 480
 ccctgcgggg gtggagaagc cccctgcacac cctgatgcac cctgcagggc tctgcaaggc 540
 atcgagagcca tccaggaaag cgcctctgtag tgggggcagc atgaaggtca atggggcgcc 600
 aacagcttca gccctctgtg ggagacaccg ctctctccag agcagcgtgc cccggcgctc 660
 attcgagacc ggccgcgggg agagcgcgac ccacgaggac ctctacttca tcccactccc 720
 gccgcgcctg tgcgccaggg cgcgcgcgac caagcagtgt gaagcagctc tgggatggca 780
 cgcgcctcgc tggcacccgc gaocccagctc gggttgccag ggggccttgc gcccaaggcg 840
 gccgcgcctg tgcgccaggg cgcgcgcgac caagcagtgt tcttcagagt gaggcgtcgc 900
 cgcgcctcgc tggcacccgc gaocccagctc gggttgccag ggggccttgc gcccaaggcg 960
 tgcctgtgtg tggcacccgc gaocccagctc gggttgccag ggggccttgc gcccaaggcg 1020
 cgcctgtgag cgcgcgcgac ctctctctgc ctctctctgc ctctctctgc ctctctctgc 1080
 ggacatcagc tccctctcta tctgtctgtc tctgtctgtc tctgtctgtc tctgtctgtc 1140
 tctgtgagtc gagtctgtgc tctgtctgtc tctgtctgtc tctgtctgtc tctgtctgtc 1200
 cagctgtgct tgcctgtgct ctctgtgtgt ctctgtgtgt ctctgtgtgt ctctgtgtgt 1260
 gccgcagagc ggggtgttgt ctctgtgtgt ctctgtgtgt ctctgtgtgt ctctgtgtgt 1320
 tagcccaaga tgcctgcagc ctctgtgtgt ctctgtgtgt ctctgtgtgt ctctgtgtgt 1380
 gccgcagagc actcactcag cctctgtgtgt ctctgtgtgt ctctgtgtgt ctctgtgtgt 1440
 tctagtcttc agccttggga ggcagcttca agccttgggg tgcagcttcc atgttgagac 1500
 actcactata atgtacctg ccagtgagag cagtgggtag cctctctctc aggttgaggc 1560
 cctcaaggag cgttgggtag cctctctctc aggttgaggc cctctctctc aggttgaggc 1620
 agtcagagtc tgcagagct cctctctctc aggttgaggc cctctctctc aggttgaggc 1680
 gttgcacact cctgcttggt gcttgccagc acagaggcgt ctgaaatgbc atgttgaggc 1740
 ctcaaggaaa gacttgaagc caagagggtg tctccctacc aggttgaggc 1800
 cagggttgga agaaagaaat caggttggtg tctccctacc aggttgaggc 1860
 ctgaggagga gaggggcctt caggttggtg tctccctacc aggttgaggc 1920
 caagatcacg ggaatgaaat caggttggtg tctccctacc aggttgaggc 1980
 tccatctctc ctctccctac ctctccctac ctctccctac ctctccctac ctctccctac 2040
 tggcctgact ggaagggaag agacttagga aactacagat tggccatgat gtcctttctt 2100

<210> 22
 <211> 2215
 <212> DNA
 <213> Homo sapiens

<400> 22
 ctgcagggag ctcatgattgc accactgcac tccagccttg taactctagg gcaacagagt gagacatgt 60
 ctcaagaaaaga aaaaanaaga aagaaacac tgcatttcct taactctagg taactctagg gacacacaga 120
 agcacaccag ctcaactcgc ttgatttccc gcaacacaca aactctaaag cctggttctc cactttgttc 180
 caagatctgag gctccagaga ggaogctggg ggcactatgc aactcactgc cctgttcttc gacctatc 240
 gactcactca ttgtttctgt caatgcagat cctgtctgtg cgcacagtc cgcacagtc cgttgcttc 300
 tattcaaga ggcacagac caatgcagat cctgtctgtg cgcacagtc cgcacagtc cgttgcttc 360
 ataactctcc tcctgcaccc ctgtctgtgc cctgtctgtg cgcacagtc cgcacagtc cgttgcttc 420
 ggtgtccact ttgcagctc catctttgoc atctttgoc atctttgoc atctttgoc atctttgoc 480
 ctgtactctc ttgtcactg actttgcagc actttgcagc actttgcagc actttgcagc actttgcagc 540

	cacccgccacc	gtgcccgcgt	ccttctcacc	agcaagctgt	cctgtgtggg	catctggata	660
	ctagccacag	tgctctccat	cccagagctc	ctgtacagtg	acotccagag	gagcagcagt	720
5	gagcaagcag	tgogatgtct	tctcatcaca	gagcatgtgg	aggcctttat	caccatccag	780
	gtggcccaaga	tggtgtatgg	cttctgtgtc	ccctgtgtgg	ccatgagcgt	ctgtttacott	840
	gtcatcatcc	gcacccctgt	ccaggccacg	aactttgagc	gcaccaaggc	catcaagggtg	900
	atcatcogct	tggtctgtgt	cttcatagtc	ttccagctgc	octacaaatg	ggtgtgtcgtg	960
	gcccagacgg	tgcccaactt	caacatcacc	agtagcaoct	gtgagctcag	taagcaactc	1020
10	aacatcgocct	acgacgtcac	ctacagocgt	gocctgocct	gctgctgocgt	caaccccttc	1080
	ttgtacgocct	toatcggcgt	caagttccgc	aaogatctct	tcaagtcctt	caaggacctg	1140
	ggctgcctcca	gcccaggagca	gctccggcag	tggtcttctc	gtccggacat	coggocgtcc	1200
	tccatgagtg	tggaaggccga	gaocccaccc	acctctcccc	cataggcgac	tcttctgocct	1260
	ggactagagg	gacctctccc	agggtccctg	gggtggggat	agggagcaga	tgcaatgaot	1320
15	caggacatcc	cccccgcata	agctgctcag	ggaaaaagag	ctctccccc	agagtgcagg	1380
	ccctgctcca	gaagttagct	tcaccccaat	cccagctacc	tcaaccaaag	cogaaaaaga	1440
	cagggtctgat	aagctaacac	cagacagaca	acactgggaa	acagaggcta	ttgtccccc	1500
	aacccaaaac	tgaaagttaa	agtcacagaa	ctgttccac	ctgctggagt	gaagggggcca	1560
	aggagggctga	gtgcaagggt	cgtgggagtg	gocctgaagag	tccctctgaat	gaacccctctg	1620
	gocctccaca	gactccaaatg	ctcagaccag	ctctccogaa	aacccaggcct	tatctccaaag	1680
20	accagagata	gtggggagac	ttcttggcct	ggtgagggaa	aggggacatc	agctgttcaa	1740
	acaaactctc	tgaacccctc	ccctccatgt	ttcttccact	gtcctccaaag	ccagcggggaa	1800
	tggtcagctgc	caocgcgcgc	taaaagcaca	ctcatccoct	caactgcgcg	gtgcgccctcc	1860
	caggctctcca	caagggggaga	gtgtggtgtt	tcctgcaggc	caagccagctc	gcctccgcgt	1920
	gatcaaaagcc	acactctggg	ctccagagtg	gggatgacat	gcactcaagct	cttggctcca	1980
	ctgggatggg	aggagggacc	aagggaatg	tcaggggggg	taagggtgac	agtgggcgcc	2040
25	caaggccagc	agcttgttgt	ttgttctttg	tcacaggggc	tgaaaaacctc	tccctaatgtt	2100
	ctgctcttga	ttcgtttaaga	gagcaacatt	ttacccacac	acagataaag	cttccctctg	2160
	aggaacaac	agctttaaaa	gaaaaaagaa	aaaaaaagct	tggtgaagta	agtag	2215

<210> 23

<211> 958

<212> DNA

<213> Homo sapiens

<400> 23

	ggggccggac	gogagggggg	ggggagaggg	gggacaaagg	gaagogaagg	oggagctggg	60
40	gggccttttt	ctgcccgcgg	tgctccagat	tcattctttaa	ggaactgaga	acttaacttt	120
	ccaaatgtct	ccaaaagaca	tcttatgcct	acatccccac	ccagctccct	gcacacacaa	180
	tgccagcac	accagggttt	gtgggataca	atccatcacg	tcatctcgcc	tacaaacaat	240
	acaggctggg	agggaaccgc	agcaccacaa	ccggggtcc	ggcatccctc	ggatcacaga	300
	ttccaaaacc	cccaagtgca	ccagataagg	cgtgtatgcc	ctcatatgag	tacagacaga	360
45	aggctgtggg	gcttccaaac	gttccaaacc	ctgacctaaa	ctatcataat	attggcaaga	420
	ttattgtgtg	catgtggoga	gatctccactg	atgaagaaaa	acaagaatat	ttaaacgaat	480
	acgaagacga	aaagatagag	tacaaatgaat	ctatgaaggc	ctatcataat	tcoccccgct	540
	accttgctta	cataaatgca	aaaagtctgtg	cagaagctgc	tttagaggaa	gaagtgctac	600
	agagacaact	tgccatggag	aaaggagaaac	cgtacatgag	cattcagcct	gctgagatcc	660
50	cagatgatta	tgatgatggc	ttttccatga	agcatcacgc	cacccgocctg	ttccagagaa	720
	accacccgct	catcagtgaa	attcttatgtg	agagtgtggt	gocagacgtt	cggttcaggtg	780
	tcacacacag	tagaatgcag	gtctccaaac	gcacaggtcca	gtccctaatg	gttccatcgc	840
	gaaaactaga	agctgaactt	cttcaaatag	aggaacgcaca	ccaggagaaag	aagaggaat	900
55	tcttggaag	cacagattcca	tttaacaaatg	aacttaaaag	gtgtgctgggt	ctgaaagt	958

<210> 24

<211> 6483

<212> DNA

<213> Homo sapiens

<400> 24

aagcttctaa	ttgcagttca	accacactgt	acatatcttc	aggaaaaaat	cacacacctct	60
caacttcaac	ttctctctct	ataaattaga	aataacaata	accacacctg	taacccccagc	120
actttgggag	gcoagggcag	gcagatcaag	aggtagaggag	attgagacaa	tcctgggctaa	180
catgtatgaa	ccctgtctct	accaaaaaga	caaaaaatta	gccaggtatg	gtggcacaca	240
ccgttagctc	cagctactcg	ggaggctgag	gcaggagaat	ggcgtgaacc	cgggagggtgg	300
agcttgacgt	gagccagat	ggcgccactg	cactccagcc	tgggcgacag	agcaagcctc	360
cgtctaaaaa	aaaaaaaaga	aagaaaagaa	gaaagaaaag	aaagaaaata	taataaccac	420
cattctcttc	tcacacagctt	gtcttagaaa	tttttaaacg	acagatacac	aaacagcact	480
acataatgtg	aaaaatctga	tgaatatata	catccaaaca	acagcaatgt	catagocctat	540
gggtagatat	aatctttatc	aatgtaccac	aatcccaatt	tacttcaact	gacaaaatgtg	600
tataccaaat	tcgttacaca	gtatatccaa	gaaaatctgt	tgtttttatt	gagaaaatga	660
acctagcttg	ggaacacatg	tgccacgtct	agttcaatat	atttggtgca	agtatacttc	720
tctaatatag	atttcaactt	ttgcaagcaa	atttttactt	gcaatcgtta	catatccaaa	780
ttttcccttt	ttactcactc	agaaacttagt	ttaaagtact	acaagttagt	tcttggtatt	840
tcagtctcag	aaaaaatatg	agattttctg	cattattatg	gtcttccacg	aaacottaac	900
tatgataaat	ttaaaagtgc	aaaaaatatc	aggataaact	tatgattcca	cattttttaa	960
tgtaaaaaat	aatgccatca	ttaatagaaa	aatctaaaaa	tcattacttc	caactttotta	1020
ggcaaaatat	caatatactc	tcatttgcca	aaataattaa	aaagatctct	acaaacacaa	1080
tcctctaaat	tggtggttta	tggtcttaat	gttttatctg	tggaacatgt	tgaatctagt	1140
taaaatttta	gaaactcttt	ctttttgatt	ccctacagtt	gtctacaaga	acctattgtg	1200
agcatgatcc	tgccagactt	tatactatct	gttgctccaa	ttaaaacctg	ttaaaacatg	1260
aatttgaaaa	atctttatct	aacataaatt	ttgtagctga	aaactttttt	tctaaacttt	1320
gcaaacatcc	tatgcaacct	gaattagtgc	tgagaaaaat	ggatcttaat	ggttgctcaa	1380
tggttctcaa	caggtgaaaa	gcataataaa	acatgctcat	ctgaaatcca	cccattttca	1440
atttcaacat	agcatacctc	gtgtttatct	ttagggcaaa	ttcaaaattg	tacatattag	1500
gattggttat	tactgaagat	aatttatgca	atcataagcc	aaagatgcta	agttgggcaaa	1560
aagaaaaaca	tgtaagttaag	caaacctotaa	cacatgtgga	ccacacctct	cagtataata	1620
aggcttgtca	ctgtccttgg	tagcaggcac	tccttggtgt	aaacagcatc	acactgtctg	1680
ttogatacac	ctcagcgaag	caactactct	ccctccgcag	tgaggagga	ggaggaggag	1740
gaggatgttg	aggaggagga	ggagtgtcat	ccctaaagat	ttctagcagc	aaaggctccc	1800
ttgtgtggag	atttagctca	ggggggttca	gtgtgtgctc	ttttagccgt	gggagctctg	1860
gtgggggatg	ctttgggggc	tcatacaggt	gctatggagg	atttagaggt	tttgggtggg	1920
gtactcttca	tggaagctat	ggaagtacga	gctttgtgtg	gagttatgga	ggcagctttg	1980
gagggggcaa	tttoggaggt	ggcagctctg	gtgggggcag	ctttgtgtga	ggcggctttg	2040
gtggaggggc	ctttggagga	ggctttgggt	gtggatttgg	aggagatggt	ccctctctct	2100
ctggaaatga	aaagtatacc	atgcagaatc	tgaatgaccc	ccctggcttc	taacttgga	2160
aagtctgggg	tcctgggaaga	aatgtagaaa	agctgggaag	agctgttacc	gagtggtatg	2220
aaaagcatgg	caactcacat	caggggggagc	ctcgtgacta	cagcaaatat	tacaaaacaa	2280
tcgtagacct	taaaaatccag	gtaagaggta	tttttaactc	agcttttaag	ctcttcttcc	2340
atgtaatcca	gcagatgtga	tccttaaatc	agcaacaatg	ggctgttcc	tatgttatcc	2400
catgttactt	ttctctctca	aaaaataccc	agtcctcaac	aagataaaca	atggggagatt	2460
taaggctatg	gcaatctcca	tcacagcaagt	gtctctactc	ttttaagagg	atgaaatgtt	2520
tactaagcac	ttttgaggtt	ttaatgagca	ctgaaaaaaa	aaatcaatgt	acaaaatgtt	2580
gctattttca	aatgtagaaa	atattoagta	ccaaactggc	aagaaaaata	aatcttccag	2640
tatgaaatg	ggacactgct	tgttcttaag	gaattctcca	attgttaact	ttgtgttaaca	2700
gtccagagtg	tattttttta	ttagtattta	tcattgagta	aaacaaatga	ggtgtgagtc	2760
agccaaagca	tggtctgaat	acatggaaat	cacatagctc	aaaagaggag	ggcacactta	2820
						2880

	caggaataaca	tctatataat	tcocagttagt	tttcagaaaag	gaataattg	tgtacagaaa	2940
	tacaagaatg	gagaatctoc	aagagaaaca	ataattcaaa	gttaagtata	tgggttaagoc	3000
	tgcaattattt	catatttcaaa	ataaaaaatt	ttococaaag	tttgtaagag	aacacacataa	3060
5	aagtgcgaag	tgcatattatg	tcactacaaa	agocatatot	gcototgaoc	totttoca	3120
	taactgtgoc	ttctocctoca	gattotocaac	ctaaocaactg	ataatgcocaa	catctotgott	3180
	oagatogaca	atgcocaggct	ggcagctgat	gaottcaggc	tgaagttaagt	taagtgaatog	3240
	ttgtataata	ctatocaacac	gaatacotaac	gtgggttttta	acaatgacttt	gggagtgocot	3300
10	caatacaactt	tacatttttco	tgaattocaac	caaatgttaa	taagtattgga	gttatctgag	3360
	aaatttttcca	tgctaatggtt	acottttttgg	caatatataaa	ggaagaaat	gcatattaaa	3420
	gtaactgctca	aggttttttco	cattaaaoca	ctattacttc	taagagaact	gtacatgaca	3480
	aatatgtgta	ttacatgaga	toaaactatgt	agttgottttt	taaatagtct	ctgcocagat	3540
	acatctocccc	tatatataagt	ataaocagta	ttgatatcat	ggcttgogta	ggtagtagaa	3600
	tgagggtagct	ctgcgccaga	gogtggaggc	tgacatcaac	ggcctgogta	gggtgctgga	3660
15	tgagctgaccc	ctgacccaagg	ctgacctgga	gatgcaaat	gagagocgtga	ctgaagagot	3720
	ggcctatctg	aagaagaaccc	acgagggaggt	gacacaaaag	cttatactttt	cccgccacaa	3780
	agagagttca	ttatgttctoc	ogtgtagcca	ataaatctttt	ctgttctoca	aacaggaat	3840
	gaagaccctt	gcaagtgtgt	ccactgggtga	tgtgaatgtg	gaaaatgaat	ctgcocccggg	3900
	ttgtgtatctg	actocaacttc	tgaataacat	gagaagccaa	taagaacac	ttgtctgaaca	3960
20	aaacocgaaa	gatgcctgaag	ccgtgttcaa	tgaaaaggtta	aagtaatctt	ccctaatagt	4020
	aaactocagg	aggtttttctc	atttcagaaat	ttctocacccc	ttttcctgtt	tttttaactat	4080
	ctagagcaagg	gaactgacta	cagaaattga	taataacatt	gaacagata	ccagctataa	4140
	atctgagagtt	actgaatttga	gaogtaattgt	acaagctctg	gagatagaac	taacgtcccca	4200
	actogctgtg	ttatgtttaac	ttctactgaaa	tgaactcaac	tttctacac	aaagtctcat	4260
	gctcaactaa	gaataatgcaa	tgcacacaaa	aaatgcagag	ttggaggtaa	gaagagagaa	4320
25	accaaagtgaa	gtctactgtta	atggaggaaa	agtaactacta	gtgttgatct	aaaagtgtgt	4380
	aaaactgaant	gggtgcacata	aacatacaac	aaattctgttt	caatttcttta	ttctctcata	4440
	taaTgctotta	ctaaaatact	aaataagcgt	caacatactc	taacgtcaact	ggaagtaact	4500
	aagococaaa	aaaaatocggt	ctcagacagt	taococccaga	ctcctctggc	agaacacagaa	4560
	gggtgcgtact	gtgtgcagct	aaocgagtg	cagaaatctg	taococctgt	ggaagacacag	4620
30	ttgcaacaga	ttogagctga	tgaatttcaa	acctaactgt	aatcaacaca	actcctggat	4680
	athtaagatoc	gaactggagaa	agttatccca	gcttcttttta	gocctgtaga	aggagagggga	4740
	aggtaaatata	taacatgaaa	agttatccca	acctaactgt	ttcaatattc	cagatagcaa	4800
	ggttatctata	aaocccaaga	agatgccaaga	gaatgagagg	aaggggagg	agaagggata	4860
	gtacagaaaa	aggagtacgc	aaocgcaatc	toactttctc	atgaatttgg	cccaaaatga	4920
	ttcttaagag	ttctgtgaac	ttcaacttgt	tttcaaggga	tggtgtttta	aatataatac	4980
35	tgccagggtt	ttatttttttc	aaacacgtttt	gottatttttc	taaatatgaag	gcaactggaa	5040
	agctacccac	ogtttttccaa	ogtttagagat	aaocgaatgt	gaactaacoc	ogtttagtct	5100
	cggagcgccg	ggaocgcccg	gogggaagttt	cggogccggc	taocggcgccg	gaagctccag	5160
	ogggcggaag	tcocgcccgg	gctacggccg	ogggccacggc	ggcagttcgc	gocggccgta	5220
	cggagcgga	agctocggcg	gogggaagctc	cggocggccg	taocggggcg	gaagctccag	5280
	cggocggccac	ggocggggaa	gctocagcgg	cggccacggc	ggcagttcca	gocggccgta	5340
40	cgggtgtggc	agttccggcg	gocggcgccg	cggctacggc	ggocggcagct	cggcgccggc	5400
	caocagctoc	ggocggcgat	acggcgccgg	cagctocagc	ggagggccaca	agctctctct	5460
	ttcggggtoc	gtggcgagct	cttcaactaa	gggaccaaag	tcagcagaaa	ctagctgggg	5520
	taactatagaa	ttagtttttaa	cttctcgtga	tggtttttttt	gocgtttaag	ctctagagtt	5580
	gttttaaaaa	attaaaaaat	tttagagaog	ttcogttttg	attttgtcac	aaactactct	5640
45	taaacocccg	cgtgaaaaat	ggcatgatac	aaatgtcata	ctttaagcat	ttttttggcg	5700
	ttcaaatctg	aaagttgaaa	tttctctctt	tttcaataat	ttttgttcta	attcataagg	5760
	atcctacag	actgttttaa	attttttttc	catcaatcac	acagatacta	acaaaacocg	5820
	agtaatccag	acaaattattg	aagaggtggc	goccgacggt	agagttcttt	catctatgg	5880
	tgaatcagaa	acaaagaac	actactatta	aactgcatac	agaggaagaa	gtctcccttc	5940
50	acacagaccca	ttattttacag	atgcattgaa	sacaaagctc	ccaagaaac	actctgtct	6000
	tgtgtgtcta	tggaataatga	acttgaaaaa	aaggtgtctc	caaggtgttt	tggtgtttct	6060
	gtattttcttc	ttttctactt	accacaaagt	gtttcttaat	ggaaagacta	tcocaaattg	6120
	gtttctacttt	actaatgaat	ttcaataaac	ttttctactt	atgcaaaacta	tcocaaattg	6180
	tcagaatttta	gttttactta	agtaataaat	actottttaa	attaaagatt	agtaacocaa	6240
	ogcgaattgaa	cgtttgagta	tcacagaatt	acgaagacag	aaactatgtc	atgcottttc	6300
55	taagttttttt	aaactgttat	gttcagaccca	cogtcagtaa	attcaactgag	taagttgtgt	6360
	aaatccoccaa	tattactctt	taagatacac	aatatgtgga	aggctccocag	ctctctggct	6420

	ttaaattatt	tcaatctgg	aaattctgga	atatctcaaa	tataaccccc	aaaataataa	6480
	taa						6483
5	<210>	25					
	<211>	1871					
10	<212>	DNA					
	<213>	Homo sapiens					
15	<400>	25					
	agttgtggcg	accttccccc	ggccatggat	ctctccaaca	acacccatgtc	actctcagtg	60
	cgcaaccccc	gactgtcccg	goggtctctc	tgcagagtg	tgataggcag	accacggggc	120
	atgtctgctt	ccagtgcttg	aagtggttat	gggggaagtg	ootttggtct	tggagccagc	180
	tgtggggag	gtctttctgc	tgcttccatg	tttggttcta	gttcoggtct	tgggggtggc	240
20	tcoggaagtt	ccatggcagg	aggactgggt	gctggttatg	ggagagccct	gggtggaggt	300
	agctttggag	gggtggggat	gggatttggg	ggcagccag	gaggtggctc	tctaggtatt	360
	ctctogggaa	atgatggagg	ootttcttct	ggatcagaaa	aagaaactat	gcacaaatctt	420
	aatgatagat	tagcttctca	octggataag	gtgcgagctc	tagaagagggc	taatacttgag	480
	ctagaataata	aaattcgaga	atggtatgaa	acaagagaa	ctgggactgc	agatgcttca	540
	cagagcaggt	acagcaata	ttatccactg	attgaagacc	tcaggaaata	gatcatttca	600
25	gcacgcatgt	gaaatgcccc	gctctctctg	cagattgaca	atggcgagact	agctgcttgag	660
	gaactccagg	tgaagtatga	gaatgaactg	gccttgogcc	agggcgtaga	ggccgacatc	720
	aatggcctgc	gcgggtgtct	ggaagagctg	acootgacca	ggacccagct	ggagatgoag	780
	atogagagcc	tgaacagagg	gctggcctac	atgaagaaga	acacagagga	tgagctccaa	840
	agcttcoggg	tggggcgccc	agggcaggtc	agcgtagaaa	tggacgctgc	ccocggagtg	900
30	gacctccaca	ggctctccaa	tgatctgggg	gogcagtatg	aaacccctgc	tgagcagaat	960
	oggaaggagc	ctgaagcctg	gttcattgaa	aagagcgggg	agctccgtaa	ggagattago	1020
	acacaccccg	agcagcttca	gtccagcaag	agcgaggtca	cgcacctgog	togcgctttt	1080
	cagaaactgg	agatcgagct	acagtcacag	ctcgccatga	agaaatccct	ggaggactcc	1140
	ttggcogaag	ccgaggcgga	ttactgogcg	cagctgtccc	aggtgcagca	gctcatcagc	1200
35	aacctggagg	cacagctgct	ccaggtgcgc	goggaogcag	agcgccagaa	cgtggaccc	1260
	cagcggctgc	tgaatgtcaa	ggcccgccctg	gagctggaga	ttgagaccta	ccgcgccttg	1320
	ctggaocggg	agggcccaag	tgatggtttg	gaggaaagtt	tatttgtgac	agactccaaa	1380
	tcacaaagcac	agtcacactga	ttctctctaa	gaccccaacca	aaacccgaaa	aatacaagaa	1440
	gtttgtgcagg	agatgggtgaa	tggtgaggtg	gtctcatctc	aagtccagga	aattgaagaa	1500
40	ctaattgtaaa	atttccaaag	atctgcccc	tgattggttc	cttaggaaca	agaaatttca	1560
	aagtgaagaat	tattcttctc	agagtaacat	gctgtattac	tcaatccctt	attttgtct	1620
	gttccatttt	ctttggatcc	ccatctccca	ttgaatcttt	tttgcccttc	tgaacacata	1680
	ttcagtcaca	agtcattttg	gtcatgttgg	tctttgtaac	aaaaccaaat	taacttatat	1740
	ccctctggac	aacctggagta	gtcttttaac	gaactcttct	ctggtaaccc	ggaaattttt	1800
	cttaactcata	gagctttact	caagtagtat	tgttttaata	gagttaattg	taataaaaga	1860
45	tgaatggtaa	a					1871
	<210>	26					
	<211>	1447					
50	<212>	DNA					
	<213>	Homo sapiens					

<400> 26
 5 ctgcaactgg tctctgagag gctccttcaa tggcagcagag aaggagaacta tgcagtctcct 60
 gaacgaacgc ctggccagct aacctgagaa ggtgcgtcac gtggagcggg acaacgcgga 120
 gctgggaacac ctcatocggg agcgggtctca gcagcaggag cccctgtcgt gccacagcta 180
 ccagtctaac ttcgaagaca ttgaggagct ccagcagaag atcctgttgc gccagtctga 240
 gaattgccag ctggtgtgtgc agatogacaa tgcacagctg cctgcagatg acttcagaac 300
 caagtaccag acggagcagt cccctgogcca gctggtggag tcogacatca acagcctgcg 360
 10 caggattctg gatgagctga cccctgtgcag gtctgacotg gaggococaga tggagctcct 420
 gaaggagagg ctgctgttcc tcaagcagaa ccatgacag caagtcaaca ccttgogctg 480
 ccagcttgga gacgcctcaa acgtggaggt ggacgctgct cccgctgtgg accctgaacca 540
 ggtcctgaac gagacacagg atcagtatga ggccctggtg gaacccaacc gcagggaagt 600
 ggagcaatgg ttccgccacg agacogagga gctgaacaag cagggtggtat ccagctcogga 660
 gcagctgcag tctaccaggc oggagatoat cgagctgaga cgcacagta atgcctcogga 720
 15 gatogagctg caggccacgc acaacctcgc atactctctg gaaacacacg tgcacagagag 780
 cgaggccccc tacagctccc agctgttcca ggtgcagagc ctgatcaaca acgtggagtc 840
 ccagctggcg gagatoccaa gtgacctgga ggggcagaa caggagatct aggtgctgct 900
 ggacgtggcg gcgcggctgg agtgtgagat caacacatac cggagcctgc tggagagcga 960
 ggacgtcaag ctgccttcaa acccctgcgc ccccaacaaat gcatgtgaaa agccacttgg 1020
 atcctgtgtc accaaactct gtggttccgtg ttcccgctgt gggccttgca acacacttgg 1080
 20 gtactagata agcagaagta tagcatgaag caacagactac catcggtggg 1140
 ccagttctgc ctctctgaca accatcagcc acgggaacccc accccagggc atcaacacaa 1200
 atcattgtct ggaaggagaa caaatgcoca cggtttgggt ctgactctga gcatagggt 1260
 actgatctct ctaccccagc gtccctctcc tctagtctcgt ctgagttctg atggtcagag 1320
 gttgtgagctg tgacagtggt atcagaggtg ttttcttctc totgtcgtct ctacctttat 1380
 25 tgcagttccc caaatgcct aataaacctt cctcttgcaa agcagacaaa aaaaaaaaaa 1440
 aaaaaaa 1447

<210> 27

<211> 261

<212> PRT

<213> Homo sapiens

<400> 27
 40 Met Asn Pro Asn Cys Ala Arg Cys Gly Lys Ile Val Tyr Pro Thr Glu
 1 5 10 15
 Lys Val Asn Cys Leu Asp Lys Phe Trp His Lys Ala Cys Phe His Cys
 20 25 30
 Glu Thr Cys Lys Met Thr Leu Asn Met Lys Asn Tyr Lys Gly Thr Glu
 35 40 45
 Lys Lys Pro Tyr Cys Asn Ala His Tyr Pro Lys Gln Ser Phe Thr Met
 50 55 60
 Val Ala Asp Thr Pro Glu Asn Leu Arg Leu Lys Gln Gln Ser Glu Leu
 65 70 75 80
 Gln Ser Gln Val Arg Tyr Lys Glu Glu Phe Glu Lys Asn Lys Gly Lys
 85 90 95
 Gly Phe Ser Val Val Ala Asp Thr Pro Glu Leu Gln Arg Ile Lys Lys
 100 105 110
 Thr Gln Asp Gln Ile Ser Asn Ile Lys Tyr His Glu Glu Phe Glu Lys
 115 120 125
 Ser Arg Met Gly Pro Ser Gly Gly Glu Gly Met Glu Pro Glu Arg Arg
 130 135 140
 55 Asp Ser Gln Asp Gly Ser Ser Tyr Arg Arg Pro Leu Glu Gln Gln Gln
 145 150 155 160

Pro His His Ile Pro Thr Ser Ala Pro Val Tyr Gln Gln Pro Gln Gln
 165 170 175
 Gln Pro Val Ala Gln Ser Tyr Gly Gly Tyr Lys Glu Pro Ala Ala Pro
 180 185 190
 Val Ser Ile Gln Arg Ser Ala Pro Gly Gly Gly Lys Arg Tyr Arg
 195 200 205
 Ala Val Tyr Asp Tyr Ser Ala Ala Asp Glu Asp Glu Val Ser Phe Gln
 210 215 220
 Asp Gly Asp Thr Ile Val Asn Val Gln Gln Ile Asp Asp Gly Trp Met
 225 230 235 240
 Tyr Gly Thr Val Glu Arg Thr Gly Asp Thr Gly Met Leu Pro Ala Asn
 245 250 255
 Tyr Val Glu Ala Ile
 260

<210> 28

<211> 478

<212> PRT

<213> Homo sapiens

<400> 28

Met Val Gln Lys Thr Ser Met Ser Arg Gly Pro Tyr Pro Pro Ser Gln
 1 5 10 15
 Glu Ile Pro Met Glu Val Phe Asp Pro Ser Pro Gln Gly Lys Tyr Ser
 20 25 30
 Lys Arg Lys Gly Arg Phe Lys Arg Ser Asp Gly Ser Thr Ser Ser Asp
 35 40 45
 Thr Thr Ser Asn Ser Phe Val Arg Gln Gly Ser Ala Glu Ser Tyr Thr
 50 55 60
 Ser Arg Pro Ser Asp Ser Asp Val Ser Leu Glu Glu Asp Arg Glu Ala
 65 70 75 80
 Leu Arg Lys Glu Ala Glu Arg Gln Ala Leu Ala Gln Leu Glu Lys Ala
 85 90 95
 Lys Thr Lys Pro Val Ala Phe Ala Val Arg Thr Asn Val Gly Tyr Asn
 100 105 110
 Pro Ser Pro Gly Asp Glu Val Pro Val Gln Gly Val Ala Ile Thr Phe
 115 120 125
 Glu Pro Lys Asp Phe Leu His Ile Lys Glu Lys Tyr Asn Asn Asp Trp
 130 135 140
 Trp Ile Gly Arg Leu Val Lys Glu Gly Cys Glu Val Gly Phe Ile Pro
 145 150 155 160
 Ser Pro Val Lys Leu Asp Ser Leu Arg Leu Leu Gln Glu Gln Lys Leu
 165 170 175
 Arg Gln Asn Arg Leu Gly Ser Ser Lys Ser Gly Asp Asn Ser Ser Ser
 180 185 190
 Ser Leu Gly Asp Val Val Thr Gly Thr Arg Arg Pro Thr Pro Pro Ala
 195 200 205
 Ser Ala Lys Gln Lys Gln Lys Ser Thr Glu His Val Pro Pro Tyr Asp
 210 215 220
 Val Val Pro Ser Met Arg Pro Ile Ile Leu Val Gly Pro Ser Leu Lys
 225 230 235 240
 Gly Tyr Glu Val Thr Asp Met Met Gln Lys Ala Leu Phe Asp Phe Leu
 245 250 255

	Lys	His	Arg	Phe	Asp	Gly	Arg	Ile	Ser	Ile	Thr	Arg	Val	Thr	Ala	Asp
				260					265						270	
5	Ile	Ser	Leu	Ala	Lys	Arg	Ser	Val	Leu	Asn	Asn	Pro	Ser	Lys	His	Ile
			275					280					285			
	Ile	Ile	Glu	Arg	Ser	Asn	Thr	Arg	Ser	Ser	Leu	Ala	Glu	Val	Gln	Ser
			290				295						300			
	Glu	Ile	Glu	Arg	Ile	Phe	Glu	Leu	Ala	Arg	Thr	Ile	Gln	Leu	Val	Ala
	305					310					315					320
10	Leu	Asp	Ala	Asp	Thr	Ile	Asn	His	Pro	Ala	Gln	Leu	Ser	Lys	Thr	Ser
				325						330					335	
	Leu	Ala	Pro	Ile	Ile	Val	Tyr	Ile	Lys	Ile	Thr	Ser	Pro	Lys	Val	Leu
				340					345					350		
	Gln	Arg	Leu	Ile	Lys	Ser	Arg	Gly	Lys	Ser	Gln	Ser	Lys	His	Leu	Asn
			355					360					365			
15	Val	Gln	Ile	Ala	Ala	Ser	Glu	Lys	Leu	Ala	Gln	Cys	Pro	Pro	Glu	Met
			370				375						380			
	Phe	Asp	Ile	Ile	Leu	Asp	Glu	Asn	Gln	Leu	Glu	Asp	Ala	Cys	Glu	His
	385					390					395					400
20	Leu	Ala	Glu	Tyr	Ile	Glu	Ala	Tyr	Trp	Lys	Ala	Thr	His	Pro	Pro	Ser
				405						410					415	
	Ser	Thr	Pro	Pro	Asn	Pro	Leu	Leu	Asn	Arg	Thr	Met	Ala	Thr	Ala	Ala
				420					425					430		
	Leu	Arg	Arg	Ser	Pro	Ala	Pro	Val	Ser	Asn	Leu	Gln	Val	Gln	Val	Leu
				435				440					445			
25	Thr	Ser	Leu	Arg	Arg	Asn	Leu	Gly	Phe	Trp	Gly	Gly	Leu	Glu	Ser	Ser
			450				455					460				
	Gln	Arg	Gly	Ser	Val	Val	Pro	Gln	Glu	Gln	Glu	His	Ala	Met		
	465					470						475				
	<210>	29														
30	<211>	196														
	<212>	PRT														
35	<213>	Homo sapiens														
	<400>	29														
40	Met	Ser	Met	Leu	Arg	Leu	Gln	Lys	Arg	Leu	Ala	Ser	Ser	Val	Leu	Arg
	1			5						10					15	
	Cys	Gly	Lys	Lys	Lys	Val	Trp	Leu	Asp	Pro	Asn	Glu	Thr	Asn	Glu	Ile
				20					25					30		
	Ala	Asn	Ala	Asn	Ser	Arg	Gln	Gln	Ile	Arg	Lys	Leu	Ile	Lys	Asp	Gly
			35				40					45				
45	Leu	Ile	Ile	Arg	Lys	Pro	Val	Thr	Val	His	Ser	Arg	Ala	Arg	Cys	Arg
			50				55					60				
	Lys	Asn	Thr	Leu	Ala	Arg	Arg	Lys	Gly	Arg	His	Met	Gly	Ile	Gly	Lys
	65					70										

Leu Lys Ala Asp Lys Ala Arg Lys Lys Leu Leu Ala Asp Gln Ala Glu
 145 150 155 160
 Ala Arg Arg Ser Lys Thr Lys Glu Ala Arg Lys Arg Arg Glu Glu Arg
 165 170 175
 Leu Gln Ala Lys Lys Glu Glu Ile Ile Lys Thr Leu Ser Lys Glu Glu
 180 185 190
 Glu Thr Lys Lys
 195

<210> 30

<211> 1566

<212> PRT

<213> Homo sapiens

<400> 30

Met Ser Ser Leu Leu Glu Arg Leu His Ala Lys Phe Asn Gln Asn Arg
 1 5 10 15
 Pro Trp Ser Glu Thr Ile Lys Leu Val Arg Gln Val Met Glu Lys Arg
 20 25 30
 Val Val Met Ser Ser Gly Gly His Gln His Leu Val Ser Cys Leu Glu
 35 40 45
 Thr Leu Gln Lys Ala Leu Lys Val Thr Ser Leu Pro Ala Met Thr Asp
 50 55 60
 Arg Leu Glu Ser Ile Ala Gly Gln Asn Gly Leu Gly Ser His Leu Ser
 65 70 75 80
 Ala Ser Gly Thr Glu Cys Tyr Ile Thr Ser Asp Met Phe Tyr Val Glu
 85 90 95
 Val Gln Leu Asp Pro Ala Gly Gln Leu Cys Asp Val Lys Val Ala His
 100 105 110
 His Gly Glu Asn Pro Val Ser Cys Pro Glu Leu Val Gln Gln Leu Arg
 115 120 125
 Glu Lys Asn Ser Asp Glu Phe Ser Lys His Leu Lys Gly Leu Val Asn
 130 135 140
 Leu Tyr Asn Leu Pro Gly Asp Asn Lys Leu Lys Thr Lys Met Tyr Leu
 145 150 155 160
 Ala Leu Gln Ser Leu Glu Gln Asp Leu Ser Lys Met Ala Ile Met Tyr
 165 170 175
 Trp Lys Ala Thr Asn Ala Gly Pro Leu Asp Lys Ile Leu His Gly Ser
 180 185 190
 Val Gly Tyr Leu Thr Pro Arg Ser Gly Gly His Leu Met Asn Leu Lys
 195 200 205
 Tyr Tyr Val Ser Pro Ser Asp Leu Leu Asp Asp Lys Thr Ala Ser Pro
 210 215 220
 Ile Ile Leu His Glu Asn Asn Val Ser Arg Ser Leu Gly Met Asn Ala
 225 230 235 240
 Ser Val Thr Ile Glu Gly Thr Ser Ala Val Tyr Lys Leu Pro Ile Ala
 245 250 255
 Pro Leu Ile Met Gly Ser His Pro Val Asp Asn Lys Trp Thr Pro Ser
 260 265 270
 Phe Ser Ser Ile Thr Ser Ala Asn Ser Val Asp Leu Pro Ala Cys Phe
 275 280 285
 Phe Leu Lys Phe Pro Gln Pro Ile Pro Val Ser Arg Ala Phe Val Gln
 290 295 300

Lys Leu Gln Asn Cys Thr Gly Ile Pro Leu Phe Glu Thr Gln Pro Thr
 305 310 315 320
 Tyr Ala Pro Leu Tyr Glu Leu Ile Thr Gln Phe Glu Leu Ser Lys Asp
 325 330 335
 Pro Asp Pro Ile Pro Leu Asn His Asn Met Arg Phe Tyr Ala Ala Leu
 340 345 350
 Pro Gly Gln Gln His Cys Tyr Phe Leu Asn Lys Asp Ala Pro Leu Pro
 355 360 365
 Asp Gly Arg Ser Leu Gln Gly Thr Leu Val Ser Lys Ile Thr Phe Gln
 370 375 380
 His Pro Gly Arg Val Pro Leu Ile Leu Asn Leu Ile Arg His Gln Val
 385 390 395 400
 Ala Tyr Asn Thr Leu Ile Gly Ser Cys Val Lys Arg Thr Ile Leu Lys
 405 410 415

 Glu Asp Ser Pro Gly Leu Leu Gln Phe Glu Val Cys Pro Leu Ser Glu
 420 425 430
 Ser Arg Phe Ser Val Ser Phe Gln His Pro Val Asn Asp Ser Leu Val
 435 440 445
 Cys Val Val Met Asp Val Gln Gly Leu Thr His Val Ser Cys Lys Leu
 450 455 460
 Tyr Lys Gly Leu Ser Asp Ala Leu Ile Cys Thr Asp Asp Phe Ile Ala
 465 470 475 480
 Lys Val Val Gln Arg Cys Met Ser Ile Pro Val Thr Met Arg Ala Ile
 485 490 495
 Arg Arg Lys Ala Glu Thr Ile Gln Ala Asp Thr Pro Ala Leu Ser Leu
 500 505 510
 Ile Ala Glu Thr Val Glu Asp Met Val Lys Lys Asn Leu Pro Pro Ala
 515 520 525
 Ser Ser Pro Gly Tyr Gly Met Thr Thr Gly Asn Asn Pro Met Ser Gly
 530 535 540
 Thr Thr Thr Ser Thr Asn Thr Phe Pro Gly Gly Pro Ile Ala Thr Leu
 545 550 555 560
 Phe Asn Met Ser Met Ser Ile Lys Asp Arg His Glu Ser Val Gly His
 565 570 575
 Gly Glu Asp Phe Ser Lys Val Ser Gln Asn Pro Ile Leu Thr Ser Leu
 580 585 590
 Leu Gln Ile Thr Gly Asn Gly Gly Ser Thr Ile Gly Ser Ser Pro Thr
 595 600 605
 Pro Pro His His Thr Pro Pro Val Ser Ser Met Ala Gly Asn Thr
 610 615 620
 Lys Asn His Pro Met Leu Met Asn Leu Leu Lys Asp Asn Pro Ala Gln
 625 630 635 640
 Asp Phe Ser Thr Leu Tyr Gly Ser Ser Pro Leu Glu Arg Gln Asn Ser
 645 650 655
 Ser Ser Gly Ser Pro Arg Met Glu Ile Cys Ser Gly Ser Asn Lys Thr
 660 665 670
 Lys Lys Lys Lys Ser Ser Arg Leu Pro Pro Glu Lys Pro Lys His Gln
 675 680 685
 Thr Glu Asp Asp Phe Gln Arg Glu Leu Phe Ser Met Asp Val Asp Ser
 690 695 700
 Gln Asn Pro Ile Phe Asp Val Asn Met Thr Ala Asp Thr Leu Asp Thr
 705 710 715 720
 Pro His Ile Thr Pro Ala Pro Ser Gln Cys Ser Thr Pro Pro Thr Thr
 725 730 735
 Tyr Pro Gln Pro Val Pro His Pro Gln Pro Ser Ile Gln Arg Met Val
 740 745 750
 Arg Leu Ser Ser Ser Asp Ser Ile Gly Pro Asp Val Thr Asp Ile Leu
 755 760 765

Ser Asp Ile Ala Glu Glu Ala Ser Lys Leu Pro Ser Thr Ser Asp Asp
 770 775 780
 Cys Pro Ala Ile Gly Thr Pro Leu Arg Asp Ser Ser Ser Ser Gly His
 785 790 795 800
 Ser Gln Ser Thr Leu Phe Asp Ser Asp Val Phe Gln Thr Asn Asn Asn
 805 810 815
 Glu Asn Pro Tyr Thr Asp Pro Ala Asp Leu Ile Ala Asp Ala Ala Gly
 820 825 830
 Ser Pro Ser Ser Asp Ser Pro Thr Asn His Phe Phe His Asp Gly Val
 835 840 845
 Asp Phe Asn Pro Asp Leu Leu Asn Ser Gln Ser Gln Ser Gly Phe Gly
 850 855 860
 Glu Glu Tyr Phe Asp Glu Ser Ser Gln Ser Gly Asp Asn Asp Asp Phe
 865 870 875 880
 Lys Gly Phe Ala Ser Gln Ala Leu Asn Thr Leu Gly Val Pro Met Leu
 885 890 895
 Gly Gly Asp Asn Gly Glu Thr Lys Phe Lys Gly Asn Asn Gln Ala Asp
 900 905 910
 Thr Val Asp Phe Ser Ile Ile Ser Val Ala Gly Lys Ala Leu Ala Pro
 915 920 925
 Ala Asp Leu Met Glu His His Ser Gly Ser Gln Gly Pro Leu Leu Thr
 930 935 940
 Thr Gly Asp Leu Gly Lys Glu Lys Thr Gln Lys Arg Val Lys Glu Gly
 945 950 955 960
 Asn Gly Thr Ser Asn Ser Thr Leu Ser Gly Pro Gly Leu Asp Ser Lys
 965 970 975
 Pro Gly Lys Arg Ser Arg Thr Pro Ser Asn Asp Gly Lys Ser Lys Asp
 980 985 990
 Lys Pro Pro Lys Arg Lys Lys Ala Asp Thr Glu Gly Lys Ser Pro Ser
 995 1000 1005
 His Ser Ser Ser Asn Arg Pro Phe Thr Pro Pro Thr Ser Thr Gly
 1010 1015 1020
 Gly Ser Lys Ser Pro Gly Ser Ala Gly Arg Ser Gln Thr Pro Pro
 1025 1030 1035
 Gly Val Ala Thr Pro Pro Ile Pro Lys Ile Thr Ile Gln Ile Pro
 1040 1045 1050
 Lys Gly Thr Val Met Val Gly Lys Pro Ser Ser His Ser Gln Tyr
 1055 1060 1065
 Thr Ser Ser Gly Ser Val Ser Ser Ser Gly Ser Lys Ser His His
 1070 1075 1080
 Ser His Ser Ser Ser Ser Ser Ser Ser Ala Ser Thr Ser Gly Lys
 1085 1090 1095
 Met Lys Ser Ser Lys Ser Glu Gly Ser Ser Ser Ser Lys Leu Ser
 1100 1105 1110
 Ser Ser Met Tyr Ser Ser Gln Gly Ser Ser Gly Ser Ser Gln Ser
 1115 1120 1125
 Lys Asn Ser Ser Gln Ser Gly Gly Lys Pro Gly Ser Ser Pro Ile
 1130 1135 1140
 Thr Lys His Gly Leu Ser Ser Gly Ser Ser Ser Thr Lys Met Lys
 1145 1150 1155
 Pro Gln Gly Lys Pro Ser Ser Leu Met Asn Pro Ser Leu Ser Lys
 1160 1165 1170
 Pro Asn Ile Ser Pro Ser His Ser Arg Pro Pro Gly Gly Ser Asp
 1175 1180 1185
 Lys Leu Ala Ser Pro Met Lys Pro Val Pro Gly Thr Pro Pro Ser
 1190 1195 1200
 Ser Lys Ala Lys Ser Pro Ile Ser Ser Gly Ser Gly Gly Ser His
 1205 1210 1215

Met Ser Gly Thr Ser Ser Ser Ser Gly Met Lys Ser Ser Ser Gly
 1220 1225
 Leu Gly Ser Ser Gly Ser Leu Ser Gln Lys Thr Pro Pro Ser Ser
 1235 1240
 Asn Ser Cys Thr Ala Ser Ser Ser Ser Phe Ser Ser Ser Gly Ser
 1250 1255
 Ser Met Ser Ser Ser Gln Asn Gln His Gly Ser Ser Lys Gly Lys
 1265 1270
 Ser Pro Ser Arg Asn Lys Lys Pro Ser Leu Thr Ala Val Ile Asp
 1280 1285
 Lys Leu Lys His Gly Val Val Thr Ser Gly Pro Gly Gly Glu Asp
 1295 1300
 Pro Leu Asp Gly Gln Met Gly Val Ser Thr Asn Ser Ser Ser His
 1310 1315
 Pro Met Ser Ser Lys His Asn Met Ser Gly Gly Glu Phe Gln Gly
 1325 1330
 Lys Arg Glu Lys Ser Asp Lys Asp Lys Ser Lys Val Ser Thr Ser
 1340 1345
 Gly Ser Ser Val Asp Ser Ser Lys Lys Thr Ser Glu Ser Lys Asn
 1355 1360
 Val Gly Ser Thr Gly Val Ala Lys Ile Ile Ile Ser Lys His Asp
 1370 1375
 Gly Gly Ser Pro Ser Ile Lys Ala Lys Val Thr Leu Gln Lys Pro
 1385 1390
 Gly Glu Ser Ser Gly Glu Gly Leu Arg Pro Gln Met Ala Ser Ser
 1400 1405
 Lys Asn Tyr Gly Ser Pro Leu Ile Ser Gly Ser Thr Pro Lys His
 1415 1420
 Glu Arg Gly Ser Pro Ser His Ser Lys Ser Pro Ala Tyr Thr Pro
 1430 1435
 Gln Asn Leu Asp Ser Glu Ser Glu Ser Gly Ser Ser Ile Ala Glu
 1445 1450
 Lys Ser Tyr Gln Asn Ser Pro Ser Ser Asp Asp Gly Ile Arg Pro
 1460 1465
 Leu Pro Glu Tyr Ser Thr Glu Lys His Lys Lys His Lys Lys Glu
 1475 1480
 Lys Lys Lys Val Lys Asp Lys Asp Arg Asp Arg Asp Arg Asp Lys
 1490 1495
 Asp Arg Asp Lys Lys Lys Ser His Ser Ile Lys Pro Glu Ser Trp
 1505 1510
 Ser Lys Ser Pro Ile Ser Ser Asp Gln Ser Leu Ser Met Thr Ser
 1520 1525
 Asn Thr Ile Leu Ser Ala Asp Arg Pro Ser Arg Leu Ser Pro Asp
 1535 1540
 Phe Met Ile Gly Glu Glu Asp Asp Asp Leu Met Asp Val Ala Leu
 1550 1555
 Ile Gly Asn
 1565

<210> 31

<211> 1490

<212> PRT

<213> Homo sapiens

<400> 31
 Met Pro Asn Ser Glu Arg His Gly Gly Lys Lys Asp Gly Ser Gly Gly
 1 5 10 15
 Ala Ser Gly Thr Leu Gln Pro Ser Ser Gly Gly Gly Ser Ser Asn Ser
 20 25 30
 Arg Glu Arg His Arg Leu Val Ser Lys His Lys Arg His Lys Ser Lys
 35 40 45
 His Ser Lys Asp Met Gly Leu Val Thr Pro Glu Ala Ala Ser Leu Gly
 50 55 60
 Thr Val Ile Lys Pro Leu Val Glu Tyr Asp Asp Ile Ser Ser Asp Ser
 65 70 75 80
 Asp Thr Phe Ser Asp Asp Met Ala Phe Lys Leu Asp Arg Arg Gly Asn
 85 90 95
 Asp Glu Arg Arg Gly Ser Asp Arg Ser Asp Arg Leu His Lys His Arg
 100 105 110
 His His Gln His Arg Arg Ser Arg Asp Leu Leu Lys Ala Lys Gln Thr
 115 120 125
 Glu Lys Glu Lys Ser Gln Glu Val Ser Ser Lys Ser Gly Ser Met Lys
 130 135 140
 Asp Arg Ile Ser Gly Ser Ser Lys Arg Ser Asn Glu Glu Thr Asp Asp
 145 150 155 160
 Tyr Gly Lys Ala Gln Val Ala Lys Ser Ser Ser Lys Glu Ser Arg Ser
 165 170 175
 Ser Lys Leu His Lys Glu Lys Thr Arg Lys Glu Arg Glu Leu Lys Ser
 180 185 190
 Gly His Lys Asp Arg Ser Lys Ser His Arg Lys Arg Glu Thr Pro Lys
 195 200 205
 Ser Tyr Lys Thr Val Asp Ser Pro Lys Arg Arg Ser Arg Ser Pro His
 210 215 220
 Arg Lys Trp Ser Asp Ser Ser Lys Gln Asp Asp Ser Pro Ser Gly Ala
 225 230 235 240
 Ser Tyr Gly Gln Asp Tyr Asp Leu Ser Pro Ser Arg Ser His Thr Ser
 245 250 255
 Ser Asn Tyr Asp Ser Tyr Lys Lys Ser Pro Gly Ser Thr Ser Arg Arg
 260 265 270
 Gln Ser Val Ser Pro Pro Tyr Lys Lys Glu Pro Ser Ala Tyr Gln Ser Ser
 275 280 285
 Thr Arg Ser Pro Ser Pro Tyr Ser Arg Arg Gln Arg Ser Val Ser Pro
 290 295 300
 Tyr Ser Arg Arg Arg Ser Ser Ser Tyr Glu Arg Ser Gly Ser Tyr Ser
 305 310 315 320
 Gly Arg Ser Pro Ser Pro Tyr Gly Arg Arg Arg Ser Ser Ser Pro Phe
 325 330 335
 Leu Ser Lys Arg Ser Leu Ser Arg Ser Pro Leu Pro Ser Arg Lys Ser
 340 345 350
 Met Lys Ser Arg Ser Arg Ser Pro Ala Tyr Ser Arg His Ser Ser Ser
 355 360 365
 His Ser Lys Lys Lys Arg Ser Ser Ser Arg Ser Arg His Ser Ser Ile
 370 375 380
 Ser Pro Val Arg Leu Pro Leu Asn Ser Ser Gly Ala Glu Leu Ser
 385 390 395 400
 Arg Lys Lys Lys Glu Arg Ala Ala Ala Ala Ala Lys Met Asp
 405 410 415
 Gly Lys Glu Ser Lys Gly Ser Pro Val Phe Leu Pro Arg Lys Glu Asn
 420 425 430
 Ser Ser Val Glu Ala Lys Asp Ser Gly Leu Glu Ser Lys Lys Leu Pro
 435 440 445
 Arg Ser Val Lys Leu Glu Lys Ser Ala Pro Asp Thr Glu Leu Val Asn
 450 455 460

Val Thr His Leu Asn Thr Glu Val Lys Asn Ser Ser Asp Thr Gly Lys
 465 470 475 480
 Val Lys Leu Asp Glu Asn Ser Glu Lys His Leu Val Lys Asp Leu Lys
 485 490 495
 Ala Gln Gly Thr Arg Asp Ser Lys Pro Ile Ala Leu Lys Glu Glu Ile
 500 505 510
 Val Thr Pro Lys Glu Thr Glu Thr Ser Glu Lys Glu Thr Pro Pro Pro
 515 520 525
 Leu Pro Thr Ile Ala Ser Pro Pro Pro Leu Pro Thr Thr Thr Pro
 530 535 540
 Pro Pro Gln Thr Pro Pro Leu Pro Pro Ile Pro Ala Leu
 545 550 555 560
 Pro Gln Gln Pro Pro Leu Pro Pro Ser Gln Pro Ala Phe Ser Gln Val
 565 570 575
 Pro Ala Ser Ser Thr Ser Thr Leu Pro Ser Thr His Ser Lys Thr
 580 585 590 595
 Ser Ala Val Ser Ser Gln Ala Asn Ser Gln Pro Pro Val Gln Val Ser
 595 600 605
 Val Lys Thr Gln Val Ser Val Thr Ala Ala Ile Pro His Leu Lys Thr
 610 615 620
 Ser Thr Leu Pro Pro Leu Pro Leu Pro Pro Leu Leu Pro Gly Gly Asp
 625 630 635 640
 Asp Met Asp Ser Pro Lys Glu Thr Leu Pro Ser Lys Pro Val Lys Lys
 645 650 655
 Glu Lys Glu Gln Arg Thr Arg His Leu Leu Thr Asp Leu Pro Leu Pro
 660 665 670
 Pro Glu Leu Pro Gly Gly Asp Leu Ser Pro Pro Asp Ser Pro Glu Pro
 675 680 685
 Lys Ala Ile Thr Pro Pro Gln Gln Pro Tyr Lys Lys Arg Pro Lys Ile
 690 695 700
 Cys Cys Pro Arg Tyr Gly Glu Arg Arg Gln Glu Ser Asp Trp Gly
 705 710 715 720
 Lys Arg Cys Val Asp Lys Phe Asp Ile Ile Gly Ile Ile Gly Glu Gly
 725 730 735
 Thr Tyr Gly Gln Val Tyr Lys Ala Arg Asp Lys Asp Thr Gly Glu Leu
 740 745 750
 Val Ala Leu Lys Lys Val Arg Leu Asp Asn Glu Lys Glu Gly Phe Pro
 755 760 765
 Ile Thr Ala Ile Arg Glu Ile Lys Ile Leu Arg Gln Leu Ile His Arg
 770 775 780
 Ser Val Val Asn Met Lys Glu Ile Val Thr Asp Lys Gln Asp Ala Leu
 785 790 795 800
 Asp Phe Lys Lys Asp Lys Gly Ala Phe Tyr Leu Val Phe Glu Tyr Met
 805 810 815
 Asp His Asp Leu Met Gly Leu Leu Glu Ser Gly Leu Val His Phe Ser
 820 825 830
 Glu Asp His Ile Lys Ser Phe Met Lys Gln Leu Met Glu Gly Leu Glu
 835 840 845
 Tyr Cys His Lys Lys Asn Phe Leu His Arg Asp Ile Lys Cys Ser Asn
 850 855 860
 Ile Leu Leu Asn Asn Ser Gly Gln Ile Lys Leu Ala Asp Phe Gly Leu
 865 870 875 880
 Ala Arg Leu Tyr Asn Ser Glu Glu Ser Arg Pro Tyr Thr Asn Lys Val
 885 890 895
 Ile Thr Leu Trp Tyr Arg Pro Pro Glu Leu Leu Gly Glu Glu Arg
 900 905 910
 Tyr Thr Pro Ala Ile Asp Val Trp Ser Cys Gly Cys Ile Leu Gly Glu
 915 920 925

Leu Phe Thr Lys Lys Pro Ile Phe Gln Ala Asn Leu Glu Leu Ala Gln
 930 935 940
 Leu Glu Leu Ile Ser Arg Leu Cys Gly Ser Pro Cys Pro Ala Val Trp
 945 950 955 960
 Pro Asp Val Ile Lys Leu Pro Tyr Phe Asn Thr Met Lys Pro Lys Lys
 965 970 975
 Gln Tyr Arg Arg Arg Leu Arg Glu Glu Phe Ser Phe Ile Pro Ser Ala
 980 985 990
 Ala Leu Asp Leu Leu Asp His Met Leu Thr Leu Asp Pro Ser Lys Arg
 995 1000 1005
 Cys Thr Ala Glu Gln Thr Leu Gln Ser Asp Phe Leu Lys Asp Val
 1010 1015 1020
 Glu Leu Ser Lys Met Ala Pro Pro Asp Leu Pro His Trp Gln Asp
 1025 1030 1035
 Cys His Glu Leu Trp Ser Lys Lys Arg Arg Arg Gln Arg Gln Ser
 1040 1045 1050
 Gly Val Val Val Glu Glu Pro Pro Pro Ser Lys Thr Ser Arg Lys
 1055 1060 1065
 Glu Thr Thr Ser Gly Thr Ser Thr Glu Pro Val Lys Asn Ser Ser
 1070 1075 1080
 Pro Ala Pro Pro Gln Pro Ala Pro Gly Lys Val Glu Ser Gly Ala
 1085 1090 1095
 Gly Asp Ala Ile Gly Leu Ala Asp Ile Thr Gln Gln Leu Asn Gln
 1100 1105 1110
 Ser Glu Leu Ala Val Leu Leu Asn Leu Leu Gln Ser Gln Thr Asp
 1115 1120 1125
 Leu Ser Ile Pro Gln Met Ala Gln Leu Leu Asn Ile His Ser Asn
 1130 1135 1140
 Pro Glu Met Gln Gln Gln Leu Glu Ala Leu Asn Gln Ser Ile Ser
 1145 1150 1155
 Ala Leu Thr Glu Ala Thr Ser Gln Gln Gln Asp Ser Glu Thr Met
 1160 1165 1170
 Ala Pro Glu Glu Ser Leu Lys Glu Ala Pro Ser Ala Pro Val Ile
 1175 1180 1185
 Leu Pro Ser Ala Glu Gln Met Thr Leu Glu Ala Ser Ser Thr Pro
 1190 1195 1200
 Ala Asp Met Gln Asn Ile Leu Ala Val Leu Leu Ser Gln Leu Met
 1205 1210 1215
 Lys Thr Gln Glu Pro Ala Gly Ser Leu Glu Glu Asn Asn Ser Asp
 1220 1225 1230
 Lys Asn Ser Gly Pro Gln Gly Pro Arg Arg Thr Pro Thr Met Pro
 1235 1240 1245
 Gln Glu Glu Ala Ala Ala Cys Pro Pro His Ile Leu Pro Pro Glu
 1250 1255 1260
 Lys Arg Pro Pro Glu Pro Pro Gly Pro Pro Pro Pro Pro Pro Pro
 1265 1270 1275
 Pro Pro Leu Val Glu Gly Asp Leu Ser Ser Ala Pro Gln Glu Leu
 1280 1285 1290
 Asn Pro Ala Val Thr Ala Ala Leu Leu Gln Leu Leu Ser Gln Pro
 1295 1300 1305
 Glu Ala Glu Pro Pro Gly His Leu Pro His Glu His Gln Ala Leu
 1310 1315 1320
 Arg Pro Met Glu Tyr Ser Thr Arg Pro Arg Pro Asn Arg Thr Tyr
 1325 1330 1335
 Gly Asn Thr Asp Gly Pro Glu Thr Gly Phe Ser Ala Ile Asp Thr
 1340 1345 1350
 Asp Glu Arg Asn Ser Gly Pro Ala Leu Thr Glu Ser Leu Val Gln
 1355 1360 1365

Thr Leu Val Lys Asn Arg Thr Phe Ser Gly Ser Leu Ser His Leu
 1370 1375 1380
 Gly Glu Ser Ser Ser Tyr Gln Gly Thr Gly Ser Val Gln Phe Pro
 1385 1390 1395
 Gly Asp Gln Asp Leu Arg Phe Ala Arg Val Pro Leu Ala Leu His
 1400 1405 1410
 Pro Val Val Gly Gln Pro Phe Leu Lys Ala Glu Gly Ser Ser Asn
 1415 1420 1425
 Ser Val Val His Ala Glu Thr Lys Leu Gln Asn Tyr Gly Glu Leu
 1430 1435 1440
 Gly Pro Gly Thr Thr Gly Ala Ser Ser Ser Gly Ala Gly Leu His
 1445 1450 1455
 Trp Gly Gly Pro Thr Gln Ser Ser Ala Tyr Gly Lys Leu Tyr Arg
 1460 1465 1470
 Gly Pro Thr Arg Val Pro Pro Arg Gly Gly Arg Gly Arg Gly Val
 1475 1480 1485
 Pro Tyr
 1490

 <210> 32

 <211> 381

 <212> PRT

 <213> Homo sapiens

 <400> 32
 Met Leu Thr Arg Leu Phe Ser Glu Pro Gly Leu Leu Ser Asp Val Pro
 1 5 10 15
 Lys Phe Ala Ser Trp Gly Asp Gly Glu Asp Asp Glu Pro Arg Ser Asp
 20 25 30
 Lys Gly Asp Ala Pro Pro Pro Pro Ala Pro Gly Pro Gly Ala
 35 40 45
 Pro Gly Pro Ala Arg Ala Ala Lys Pro Val Pro Leu Arg Gly Glu Glu
 50 55 60
 Gly Thr Glu Ala Thr Leu Ala Glu Val Lys Glu Glu Gly Glu Leu Gly
 65 70 75 80
 Gly Glu Glu Glu Glu Glu Glu Glu Glu Gly Leu Asp Glu Ala
 85 90 95
 Glu Gly Glu Arg Pro Lys Lys Arg Gly Pro Lys Lys Arg Lys Met Thr
 100 105 110
 Lys Ala Arg Leu Glu Arg Ser Lys Leu Arg Arg Gln Lys Ala Asn Ala
 115 120 125
 Arg Glu Arg Asn Arg Met His Asp Leu Asn Ala Ala Leu Asp Asn Leu
 130 135 140
 Arg Lys Val Val Pro Cys Tyr Ser Lys Thr Gln Lys Leu Ser Lys Ile
 145 150 155 160
 Glu Thr Leu Arg Leu Ala Lys Asn Tyr Ile Trp Ala Leu Ser Glu Ile
 165 170 175
 Leu Arg Ser Gly Lys Arg Pro Asp Leu Val Ser Tyr Val Gln Thr Leu
 180 185 190
 Cys Lys Gly Leu Ser Gln Pro Thr Thr Asn Leu Val Ala Gly Cys Leu
 195 200 205
 Gln Leu Asn Ser Arg Asn Phe Leu Thr Glu Gln Gly Ala Asp Gly Ala
 210 215 220

Gly Arg Phe His Gly Ser Gly Gly Pro Phe Ala Met His Pro Tyr Pro
 225 230 235 240
 Tyr Pro Cys Ser Arg Leu Ala Gly Ala Gln Cys Gln Ala Ala Gly Gly
 245 250 255
 Leu Gly Gly Gly Ala Ala His Ala Leu Arg Thr His Gly Tyr Cys Ala
 260 265 270
 Ala Tyr Glu Thr Leu Tyr Ala Ala Gly Gly Gly Ala Ser Pro
 275 280 285
 Asp Tyr Asn Ser Ser Glu Tyr Glu Gly Pro Leu Ser Pro Pro Leu Cys
 290 295 300
 Leu Asn Gly Asn Phe Ser Leu Lys Gln Asp Ser Ser Pro Asp His Glu
 305 310 315 320
 Lys Ser Tyr His Tyr Ser Met His Tyr Ser Ala Leu Pro Gly Ser Arg
 325 330 335
 His Gly His Gly Leu Val Phe Gly Ser Ser Ala Val Arg Gly Gly Val
 340 345 350
 His Ser Glu Asn Leu Leu Ser Tyr Asp Met His Leu His His Asp Arg
 355 360 365
 Gly Pro Met Tyr Glu Glu Leu Asn Ala Phe Phe His Asn
 370 375 380

<210> 33

<211> 445

<212> PRT

<213> Homo sapiens

<400> 33

Met Ser Lys Leu Pro Arg Glu Leu Thr Arg Asp Leu Glu Arg Ser Leu
 1 5 10 15
 Pro Ala Val Ala Ser Leu Gly Ser Ser Leu Ser His Ser Gln Ser Leu
 20 25 30
 Ser Ser His Leu Leu Pro Pro Glu Lys Arg Arg Ala Ile Ser Asp
 35 40 45
 Val Arg Arg Thr Phe Cys Leu Phe Val Thr Phe Asp Leu Leu Phe Ile
 50 55 60
 Ser Leu Leu Trp Ile Ile Glu Leu Asn Thr Asn Thr Gly Ile Arg Lys
 65 70 75 80
 Asn Leu Glu Gln Glu Ile Ile Gln Tyr Asn Phe Lys Thr Ser Phe Phe
 85 90 95
 Asp Ile Phe Val Leu Ala Phe Phe Arg Phe Ser Gly Leu Leu Leu Gly
 100 105 110
 Tyr Ala Val Leu Gln Leu Arg His Trp Trp Val Ile Ala Val Thr Thr
 115 120 125
 Leu Val Ser Ser Ala Phe Leu Ile Val Lys Val Ile Leu Ser Glu Leu
 130 135 140
 Leu Ser Lys Gly Ala Phe Gly Tyr Leu Leu Pro Ile Val Ser Phe Val
 145 150 155 160
 Leu Ala Trp Leu Glu Thr Trp Phe Leu Asp Phe Lys Val Leu Pro Gln
 165 170 175
 Glu Ala Glu Glu Glu Arg Trp Tyr Leu Ala Ala Gln Val Ala Val Ala
 180 185 190
 Arg Gly Pro Leu Leu Phe Ser Gly Ala Leu Ser Glu Gly Gln Phe Tyr
 195 200 205

Ser Pro Pro Glu Ser Phe Ala Gly Ser Asp Asn Glu Ser Asp Glu Glu
 210 215 220
 Val Ala Gly Lys Lys Ser Phe Ser Ala Gln Glu Arg Glu Tyr Ile Arg
 225 230 235 240
 Gln Gly Lys Glu Ala Thr Ala Val Val Asp Gln Ile Leu Ala Gln Glu
 245 250 255
 Glu Asn Trp Lys Phe Glu Lys Asn Asn Glu Tyr Gly Asp Thr Val Tyr
 260 265 270
 Thr Ile Glu Val Pro Phe His Gly Lys Thr Phe Ile Leu Lys Thr Phe
 275 280 285
 Leu Pro Cys Pro Ala Glu Leu Val Tyr Gln Glu Val Ile Leu Gln Pro
 290 295 300
 Glu Arg Met Val Leu Trp Asn Lys Thr Val Thr Ala Cys Gln Ile Leu
 305 310 315 320
 Gln Arg Val Glu Asp Asn Thr Leu Ile Ser Tyr Asp Val Ser Ala Gly
 325 330 335
 Ala Ala Gly Gly Val Val Ser Pro Arg Asp Phe Val Asn Val Arg Arg
 340 345 350
 Ile Glu Arg Arg Arg Asp Arg Tyr Leu Ser Ser Gly Ile Ala Thr Ser
 355 360 365
 His Ser Ala Lys Pro Pro Thr His Lys Tyr Val Arg Gly Glu Asn Gly
 370 375 380
 Pro Gly Gly Phe Ile Val Leu Lys Ser Ala Ser Asn Pro Arg Val Cys
 385 390 395 400
 Thr Phe Val Trp Ile Leu Asn Thr Asp Leu Lys Gly Arg Leu Pro Arg
 405 410 415
 Tyr Leu Ile His Gln Ser Leu Ala Ala Thr Met Phe Glu Phe Ala Phe
 420 425 430
 His Leu Arg Gln Arg Ile Ser Glu Leu Gly Ala Arg Ala
 435 440 445

<210> 34

<211> 167

<212> PRT

<213> Homo sapiens

<400> 34

Met Ala Thr Ser Glu Leu Ser Cys Glu Val Ser Glu Glu Asn Cys Glu
 1 5 10 15
 Arg Arg Glu Ala Phe Trp Ala Glu Trp Lys Asp Leu Thr Leu Ser Thr
 20 25 30
 Arg Pro Glu Glu Gly Cys Ser Leu His Glu Glu Asp Thr Gln Arg His
 35 40 45
 Glu Thr Tyr His Gln Gln Gly Gln Cys Gln Val Leu Val Gln Arg Ser
 50 55 60
 Pro Trp Leu Met Met Arg Met Gly Ile Leu Gly Arg Gly Leu Gln Glu
 65 70 75 80
 Tyr Gln Leu Pro Tyr Gln Arg Val Leu Pro Leu Pro Ile Phe Thr Pro
 85 90 95
 Ala Lys Met Gly Ala Thr Lys Glu Glu Arg Glu Asp Thr Pro Ile Gln
 100 105 110
 Leu Gln Glu Leu Leu Ala Leu Glu Thr Ala Leu Gly Gly Gln Cys Val
 115 120 125

Asp Arg Gln Glu Val Ala Glu Ile Thr Lys Gln Leu Pro Pro Val Val
 130 135 140
 Pro Val Ser Lys Pro Gly Ala Leu Arg Arg Ser Leu Ser Arg Ser Met
 145 150 155 160
 Ser Gln Glu Ala Gln Arg Gly
 165

<210> 35

<211> 282

<212> PRT

<213> Homo sapiens

<400> 35

Met Ser Gly Ala Asp Arg Ser Pro Asn Ala Gly Ala Ala Pro Asp Ser
 1 5 10 15
 Ala Pro Gly Gln Ala Ala Val Ala Ser Ala Tyr Gln Arg Phe Glu Pro
 20 25 30
 Arg Ala Tyr Leu Arg Asn Asn Tyr Ala Pro Pro Arg Gly Asp Leu Cys
 35 40 45
 Asn Pro Asn Gly Val Gly Pro Trp Lys Leu Arg Cys Leu Ala Gln Thr
 50 55 60
 Phe Ala Thr Gly Glu Val Ser Gly Arg Thr Leu Ile Asp Ile Gly Ser
 65 70 75 80
 Gly Pro Thr Val Tyr Gln Leu Leu Ser Ala Cys Ser His Phe Glu Asp
 85 90 95
 Ile Thr Met Thr Asp Phe Leu Glu Val Asn Arg Gln Glu Leu Gly Arg
 100 105 110
 Trp Leu Gln Glu Glu Pro Gly Ala Phe Asn Trp Ser Met Tyr Ser Gln
 115 120 125
 His Ala Cys Leu Ile Glu Gly Lys Gly Glu Cys Trp Gln Asp Lys Glu
 130 135 140
 Arg Gln Leu Arg Ala Arg Val Lys Arg Val Leu Pro Ile Asp Val His
 145 150 155 160
 Gln Pro Gln Pro Leu Gly Ala Gly Ser Pro Ala Pro Leu Pro Ala Asp
 165 170 175
 Ala Leu Val Ser Ala Phe Cys Leu Glu Ala Val Ser Pro Asp Leu Ala
 180 185 190
 Ser Phe Gln Arg Ala Leu Asp His Ile Thr Thr Leu Leu Arg Pro Gly
 195 200 205
 Gly His Leu Leu Leu Ile Gly Ala Leu Glu Glu Ser Trp Tyr Leu Ala
 210 215 220
 Gly Glu Ala Arg Leu Thr Val Val Pro Val Ser Glu Glu Glu Val Arg
 225 230 235 240
 Glu Ala Leu Val Arg Ser Gly Tyr Lys Val Arg Asp Leu Arg Thr Tyr
 245 250 255
 Ile Met Pro Ala His Leu Gln Thr Gly Val Asp Asp Val Lys Gly Val
 260 265 270
 Phe Phe Ala Trp Ala Gln Lys Val Gly Leu
 275 280

<210> 36

<211> 1255

<212> PRT

<213> Homo sapiens

<400> 36

```

Met Glu Leu Ala Ala Leu Cys Arg Trp Gly Leu Leu Leu Ala Leu Leu
1      5      10      15
Pro Pro Gly Ala Ala Ser Thr Gln Val Cys Thr Gly Thr Asp Met Lys
15      20      25      30
Leu Arg Leu Pro Ala Ser Pro Glu Thr His Leu Asp Met Leu Arg His
35      40      45
Leu Tyr Gln Gly Cys Gln Val Val Gln Gly Asn Leu Glu Leu Thr Tyr
50      55      60
Leu Pro Thr Asn Ala Ser Leu Ser Phe Leu Gln Asp Ile Gln Glu Val
65      70      75      80
Gln Gly Tyr Val Leu Ile Ala His Asn Gln Val Arg Gln Val Pro Leu
85      90      95
Gln Arg Leu Arg Ile Val Arg Gly Thr Gln Leu Phe Glu Asp Asn Tyr
100      105      110
Ala Leu Ala Val Leu Asp Asn Gly Asp Pro Leu Asn Asn Thr Thr Pro
115      120      125
Val Thr Gly Ala Ser Pro Gly Gly Leu Arg Glu Leu Gln Leu Arg Ser
130      135      140
Leu Thr Glu Ile Leu Lys Gly Gly Val Leu Ile Gln Arg Asn Pro Gln
145      150      155      160
Leu Cys Tyr Gln Asp Thr Ile Leu Trp Lys Asp Ile Phe His Lys Asn
165      170      175
Asn Gln Leu Ala Leu Thr Leu Ile Asp Thr Asn Arg Ser Arg Ala Cys
180      185      190
His Pro Cys Ser Pro Met Cys Lys Gly Ser Arg Cys Trp Gly Glu Ser
195      200      205
Ser Glu Asp Cys Gln Ser Leu Thr Arg Thr Val Cys Ala Gly Gly Cys
210      215      220
Ala Arg Cys Lys Gly Pro Leu Pro Thr Asp Cys Cys His Glu Gln Cys
225      230      235      240
Ala Ala Gly Cys Thr Gly Pro Lys His Ser Asp Cys Leu Ala Cys Leu
245      250      255
His Phe Asn His Ser Gly Ile Cys Glu Leu His Cys Pro Ala Leu Val
260      265      270
Thr Tyr Asn Thr Asp Thr Phe Glu Ser Met Pro Asn Pro Glu Gly Arg
275      280      285
Tyr Thr Phe Gly Ala Ser Cys Val Thr Ala Cys Pro Tyr Asn Tyr Leu
290      295      300
Ser Thr Asp Val Gly Ser Cys Thr Leu Val Cys Pro Leu His Asn Gln
305      310      315      320
Glu Val Thr Ala Glu Asp Gly Thr Gln Arg Cys Glu Lys Cys Ser Lys
325      330      335
Pro Cys Ala Arg Val Cys Tyr Gly Leu Gly Met Glu His Leu Arg Glu
340      345      350
Val Arg Ala Val Thr Ser Ala Asn Ile Gln Glu Phe Ala Gly Cys Lys
355      360      365
Lys Ile Phe Gly Ser Leu Ala Phe Leu Pro Glu Ser Phe Asp Gly Asp
370      375      380

```

Pro Ala Ser Asn Thr Ala Pro Leu Gln Pro Glu Gln Leu Gln Val Phe
 385 435 390 395 400
 Glu Thr Leu Glu Glu Ile Thr Gly Tyr Leu Tyr Ile Ser Ala Trp Pro
 405 410 415
 Asp Ser Leu Pro Asp Leu Ser Val Phe Gln Asn Leu Gln Val Ile Arg
 420 425 430

 Gly Arg Ile Leu His Asn Gly Ala Tyr Ser Leu Thr Leu Gln Gly Leu
 435 440 445
 Gly Ile Ser Trp Leu Gly Leu Arg Ser Leu Arg Glu Leu Gly Ser Gly
 450 455 460
 Leu Ala Leu Ile His His Asn Thr His Leu Cys Phe Val His Thr Val
 465 470 480
 Pro Trp Asp Gln Leu Phe Arg Asn Pro His Gln Ala Leu Leu His Thr
 485 490 495
 Ala Asn Arg Pro Glu Asp Glu Cys Val Gly Glu Gly Leu Ala Cys His
 500 505 510
 Gln Leu Cys Ala Arg Gly His Cys Trp Gly Pro Gly Pro Thr Gln Cys
 515 520 525
 Val Asn Cys Ser Gln Phe Leu Arg Gly Gln Glu Cys Val Glu Glu Cys
 530 535 540
 Arg Val Leu Gln Gly Leu Pro Arg Glu Tyr Val Asn Ala Arg His Cys
 545 550 555 560
 Leu Pro Cys His Pro Glu Cys Gln Pro Gln Asn Gly Ser Val Thr Cys
 565 570 575
 Phe Gly Pro Glu Ala Asp Gln Cys Val Ala Cys Ala His Tyr Lys Asp
 580 585 590
 Pro Pro Phe Cys Val Ala Arg Cys Pro Ser Gly Val Lys Pro Asp Leu
 595 600 605
 Ser Tyr Met Pro Ile Trp Lys Phe Pro Asp Glu Glu Gly Ala Cys Gln
 610 615 620
 Pro Cys Pro Ile Asn Cys Thr His Ser Cys Val Asp Leu Asp Asp Lys
 625 630 635 640
 Gly Cys Pro Ala Glu Gln Arg Ala Ser Pro Leu Thr Ser Ile Val Ser
 645 650 655
 Ala Val Val Gly Ile Leu Leu Val Val Leu Gly Val Val Phe Gly
 660 665 670
 Ile Leu Ile Lys Arg Arg Gln Gln Lys Ile Arg Lys Tyr Thr Met Arg
 675 680 685
 Arg Leu Leu Gln Glu Thr Glu Leu Val Glu Pro Leu Thr Pro Ser Gly
 690 695 700
 Ala Met Pro Asn Gln Ala Gln Met Arg Ile Leu Lys Glu Thr Glu Leu
 705 710 715 720
 Arg Lys Val Lys Val Leu Gly Ser Gly Ala Phe Gly Thr Val Tyr Lys
 725 730 735
 Gly Ile Trp Ile Pro Asp Gly Glu Asn Val Lys Ile Pro Val Ala Ile
 740 745 750
 Lys Val Leu Arg Glu Asn Thr Ser Pro Lys Ala Asn Lys Glu Ile Leu
 755 760 765
 Asp Glu Ala Tyr Val Met Ala Gly Val Gly Ser Pro Tyr Val Ser Arg
 770 775 780
 Leu Leu Gly Ile Cys Leu Thr Ser Thr Val Gln Leu Val Thr Gln Leu
 785 790 795 800
 Met Pro Tyr Gly Cys Leu Leu Asp His Val Arg Glu Asn Arg Gly Arg
 805 810 815
 Leu Gly Ser Gln Asp Leu Leu Asn Trp Cys Met Gln Ile Ala Lys Gly
 820 825 830
 Met Ser Tyr Leu Glu Asp Val Arg Leu Val His Arg Asp Leu Ala Ala
 835 840 845

Arg Asn Val Leu Val Lys Ser Pro Asn His Val Lys Ile Thr Asp Phe
 850 855 860
 Gly Leu Ala Arg Leu Leu Asp Ile Asp Glu Thr Glu Tyr His Ala Asp
 865 870 875 880
 Gly Gly Lys Val Pro Ile Lys Trp Met Ala Leu Glu Ser Ile Leu Arg
 885 890 895
 Arg Arg Phe Thr His Gln Ser Asp Val Trp Ser Tyr Gly Val Thr Val
 900 905 910
 Trp Glu Leu Met Thr Phe Gly Ala Lys Pro Tyr Asp Gly Ile Pro Ala
 915 920 925
 Arg Glu Ile Pro Asp Leu Leu Glu Lys Gly Glu Arg Leu Pro Gln Pro
 930 935 940
 Pro Ile Cys Thr Ile Asp Val Tyr Met Ile Met Val Lys Cys Trp Met
 945 950 955 960
 Ile Asp Ser Glu Cys Arg Pro Arg Phe Arg Glu Leu Val Ser Glu Phe
 965 970 975
 Ser Arg Met Ala Arg Asp Pro Gln Arg Phe Val Val Ile Gln Asn Glu
 980 985 990
 Asp Leu Gly Pro Ala Ser Pro Leu Asp Ser Thr Phe Tyr Arg Ser Leu
 995 1000 1005
 Leu Glu Asp Asp Asp Met Gly Asp Leu Val Asp Ala Glu Glu Tyr
 1010 1015 1020
 Leu Val Pro Gln Gln Gly Phe Phe Cys Pro Asp Pro Ala Pro Gly
 1025 1030 1035
 Ala Gly Gly Met Val His His Arg His Arg Ser Ser Thr Arg
 1040 1045 1050
 Ser Gly Gly Gly Asp Leu Thr Leu Gly Leu Glu Pro Ser Glu Glu
 1055 1060 1065
 Glu Ala Pro Arg Ser Pro Leu Ala Pro Ser Glu Gly Ala Gly Ser
 1070 1075 1080
 Asp Val Phe Asp Gly Asp Leu Gly Met Gly Ala Ala Lys Gly Leu
 1085 1090 1095
 Gln Ser Leu Pro Thr His Asp Pro Ser Pro Leu Gln Arg Tyr Ser
 1100 1105 1110
 Glu Asp Pro Thr Val Pro Leu Pro Ser Glu Thr Asp Gly Tyr Val
 1115 1120 1125
 Ala Pro Leu Thr Cys Ser Pro Gln Pro Glu Tyr Val Asn Gln Pro
 1130 1135 1140
 Asp Val Arg Pro Gln Pro Pro Ser Pro Arg Glu Gly Pro Leu Pro
 1145 1150 1155
 Ala Ala Arg Pro Ala Gly Ala Thr Leu Glu Arg Ala Lys Thr Leu
 1160 1165 1170
 Ser Pro Gly Lys Asn Gly Val Val Lys Asp Val Phe Ala Phe Gly
 1175 1180 1185
 Gly Ala Val Glu Asn Pro Glu Tyr Leu Thr Pro Gln Gly Gly Ala
 1190 1195 1200
 Ala Pro Gln Pro His Pro Pro Ala Phe Ser Pro Ala Phe Asp
 1205 1210 1215
 Asn Leu Tyr Tyr Trp Asp Gln Asp Pro Pro Glu Arg Gly Ala Pro
 1220 1225 1230
 Pro Ser Thr Phe Lys Gly Thr Pro Thr Ala Glu Asn Pro Glu Tyr
 1235 1240 1245
 Leu Gly Leu Asp Val Pro Val
 1250 1255

<210> 37

<211> 532

<212> PRT

<213> Homo sapiens

<400> 37

```

Met Glu Leu Asp 5 Leu Ser Pro Pro His Leu Ser Ser Ser Pro Glu Asp
1 10 15
Leu Trp Pro Ala 20 Pro Gly Thr Pro Gly Thr Pro Arg Pro Pro Asp
Thr Pro Leu Pro Glu Glu Val Lys Arg Ser Gln Pro Leu Ile Pro
35 40 45
Thr Thr Gly Arg Lys Leu Arg Glu Glu Glu Arg Arg Ala Thr Ser Leu
50 55 60
Pro Ser Ile Pro Asn Pro Phe Pro Glu Leu Cys Ser Pro Pro Ser Gln
65 70 75 80
Ser Pro Ile Leu Gly Gly Pro Ser Ser Ala Arg Gly Leu Leu Pro Arg
85 90 95
Asp Ala Ser Arg 100 Pro His Val Val Lys Val Tyr Ser Glu Asp Gly Ala
110 115
Cys Arg Ser Val Glu Val Ala Ala Gly Ala Thr Ala Arg His Val Cys
120 125
Glu Met Leu Val Gln Arg Ala His Ala Leu Ser Asp Glu Thr Trp Gly
130 135 140
Leu Val Glu Cys His Pro His Leu Ala Leu Glu Arg Gly Leu Glu Asp
145 150 155 160
His Glu Ser Val Val Glu Val Gln Ala Ala Trp Pro Val Gly Gly Asp
165 170 175
Ser Arg Phe Val Phe Arg Lys Asn Phe Ala Lys Tyr Glu Leu Phe Lys
180 185 190
Ser Ser Pro His Ser Leu Phe Pro Glu Lys Met Val Ser Ser Cys Leu
195 200 205
Asp Ala His Thr Gly Ile Ser His Glu Asp Leu Ile Gln Asn Phe Leu
210 215 220
Asn Ala Gly Ser Phe Pro Glu Ile Gln Gly Phe Leu Gln Leu Arg Gly
225 230 235 240
Ser Gly Arg Lys Leu Trp Lys Arg Phe Phe Cys Phe Leu Arg Arg Ser
245 250 255
Gly Leu Tyr Tyr Ser Thr Lys Gly Thr Ser Lys Asp Pro Arg His Leu
260 265 270
Gln Tyr Val Ala Asp Val Asn Glu Ser Asn Val Tyr Val Val Thr Gln
275 280 285
Gly Arg Lys Leu Tyr Gly Met Pro Thr Asp Phe Gly Phe Cys Val Lys
290 295 300
Pro Asn Lys Leu Arg Asn Gly His Lys Gly Leu Arg Ile Phe Cys Ser
305 310 315 320
Glu Asp Glu Gln Ser Arg Thr Cys Trp Leu Ala Ala Phe Arg Leu Phe
325 330 335
Lys Tyr Gly Val Gln Leu Tyr Lys Asn Tyr Gln Gln Ala Gln Ser Arg
340 345 350
His Leu His Pro Ser Cys Leu Gly Ser Pro Pro Leu Arg Ser Ala Ser
355 360 365
Asp Asn Thr Leu Val Ala Met Asp Phe Ser Gly His Ala Gly Arg Val
370 375 380

```

Ile Glu Asn Pro Arg Glu Ala Leu Ser Val Ala Leu Glu Glu Ala Gln
 385 390 395 400
 Ala Trp Arg Lys Lys Thr Asn His Arg Leu Ser Leu Pro Met Pro Ala
 405 415
 Ser Gly Thr Ser Leu Ser Ala Ala Ile His Arg Thr Gln Leu Trp Phe
 420 430
 His Gly Arg Ile Ser Arg Glu Glu Ser Gln Arg Leu Ile Gly Gln Gln
 435 440 445
 Gly Leu Val Asp Gly Leu Phe Leu Val Arg Glu Ser Gln Arg Asn Pro
 450 455 460
 Gln Gly Phe Val Leu Ser Leu Cys His Leu Gln Lys Val Lys His Tyr
 465 470 475
 Leu Ile Leu Pro Ser Glu Glu Glu Gly Arg Leu Tyr Phe Ser Met Asp
 485 490 495
 Asp Gly Gln Thr Arg Phe Thr Asp Leu Leu Gln Leu Val Glu Phe His
 500 505 510
 Gln Leu Asn Arg Gly Ile Leu Pro Cys Leu Leu Arg His Cys Cys Thr
 515 520 525
 Arg Val Ala Leu
 530

<210> 38

<211> 534

<212> PRT

<213> Homo sapiens

<400> 38

Met Lys Gln Glu Gly Ser Ala Arg Arg Arg Gly Ala Asp Lys Ala Lys
 1 5 10 15
 Pro Pro Pro Gly Gly Glu Gln Glu Pro Pro Pro Pro Pro Ala Pro
 20 25 30
 Gln Asp Val Glu Met Lys Glu Glu Ala Ala Thr Gly Gly Gly Ser Thr
 35 40 45
 Gly Glu Ala Asp Gly Lys Thr Ala Ala Ala Val Glu His Ser Gln
 50 55 60
 Arg Glu Leu Asp Thr Val Thr Leu Glu Asp Ile Lys Glu His Val Lys
 65 70 75 80
 Gln Leu Glu Lys Ala Val Ser Gly Lys Glu Pro Arg Phe Val Leu Arg
 85 90 95
 Ala Leu Arg Met Leu Pro Ser Thr Ser Arg Arg Leu Asn His Tyr Val
 100 105 110
 Leu Tyr Lys Ala Val Gln Gly Phe Phe Thr Ser Asn Asn Ala Thr Arg
 115 120 125
 Asp Phe Leu Leu Pro Phe Leu Glu Glu Pro Met Asp Thr Glu Ala Asp
 130 135 140
 Leu Gln Phe Arg Pro Arg Thr Gly Lys Ala Ala Ser Thr Pro Leu Leu
 145 150 155 160
 Pro Glu Val Glu Ala Tyr Leu Gln Leu Leu Val Val Ile Phe Met Met
 165 170 175
 Asn Ser Lys Arg Tyr Lys Glu Ala Gln Lys Ile Ser Asp Asp Leu Met
 180 185 190
 Gln Lys Ile Ser Thr Gln Asn Arg Arg Ala Leu Asp Leu Val Ala Ala
 195 200 205

Lys Cys Tyr Tyr Tyr His Ala Arg Val Tyr Glu Phe Leu Asp Lys Leu
 210 215 220
 Asp Val Val Arg Ser Phe Leu His Ala Arg Leu Arg Thr Ala Thr Leu
 225 230 235
 Arg His Asp Ala Asp Gly Gln Ala Thr Leu Leu Asn Leu Leu Leu Arg
 245 250 255
 Asn Tyr Leu His Tyr Ser Leu Tyr Asp Gln Ala Glu Lys Leu Val Ser
 260 265 270
 Lys Ser Val Phe Pro Glu Gln Ala Asn Asn Glu Trp Ala Arg Tyr
 275 280 285
 Leu Tyr Tyr Thr Gly Arg Ile Lys Ala Ile Gln Leu Glu Tyr Ser Glu
 290 295 300
 Ala Arg Arg Thr Met Thr Asn Ala Leu Arg Lys Ala Pro Gln His Thr
 305 310 315 320
 Ala Val Gly Phe Lys Gln Thr Val His Lys Leu Leu Ile Val Val Glu
 325 330 335
 Leu Leu Leu Gly Glu Ile Pro Asp Arg Leu Gln Phe Arg Gln Pro Ser
 340 345 350
 Leu Lys Arg Ser Leu Met Pro Tyr Phe Leu Leu Thr Gln Ala Val Arg
 355 360 365
 Thr Gly Asn Leu Ala Lys Phe Asn Gln Val Leu Asp Gln Phe Gly Glu
 370 375 380
 Lys Phe Gln Ala Asp Gly Thr Tyr Thr Leu Ile Arg Leu Arg His
 385 390 395 400
 Asn Val Ile Lys Thr Gly Val Arg Met Ile Ser Leu Ser Tyr Ser Arg
 405 410 415
 Ile Ser Leu Ala Asp Ile Ala Gln Lys Leu Gln Leu Asp Ser Pro Glu
 420 425 430
 Asp Ala Glu Phe Ile Val Ala Lys Ala Ile Arg Asp Gly Val Ile Glu
 435 440 445
 Ala Ser Ile Asn His Glu Lys Gly Tyr Val Gln Ser Lys Glu Met Ile
 450 455 460
 Asp Ile Tyr Ser Thr Arg Glu Pro Gln Leu Ala Phe His Gln Arg Ile
 465 470 475 480
 Ser Phe Cys Leu Asp Ile His Asn Met Ser Val Lys Ala Met Arg Phe
 485 490 495
 Pro Pro Lys Ser Tyr Asn Lys Asp Leu Glu Ser Ala Glu Glu Arg Arg
 500 505 510
 Glu Arg Glu Gln Gln Asp Leu Glu Phe Ala Lys Glu Met Ala Glu Asp
 515 520 525
 Asp Asp Asp Ser Phe Pro
 530

<210> 39

<211> 207

<212> PRT

<213> Homo sapiens

<400> 39

Met Ala Gly Pro Ala Thr Gln Ser Pro Met Lys Leu Met Ala Leu Gln
 1 5 10 15
 Leu Leu Leu Trp His Ser Ala Leu Trp Thr Val Gln Glu Ala Thr Pro
 20 25 30

Leu Gly Pro Ala Ser Ser Leu Pro Gln Ser Phe Leu Leu Lys Cys Leu
 35 40 45
 5 Glu Gln Val Arg Lys Ile Gln Gly Asp Gly Ala Ala Leu Gln Glu Lys
 50 55 60
 Leu Val Ser Glu Cys Ala Thr Tyr Lys Leu Cys His Pro Glu Glu Leu
 65 70 75 80
 Val Leu Leu Gly His Ser Leu Gly Ile Pro Trp Ala Pro Leu Ser Ser
 85 90 95
 10 Cys Pro Ser Gln Ala Leu Gln Leu Ala Gly Cys Leu Ser Gln Leu His
 100 105 110
 Ser Gly Leu Phe Leu Tyr Gln Gly Leu Gln Ala Leu Glu Gly Ile
 115 120 125
 Ser Pro Glu Leu Gly Pro Thr Leu Asp Thr Leu Gln Leu Asp Val Ala
 130 135 140
 15 Asp Phe Ala Thr Thr Ile Trp Gln Gln Met Glu Glu Leu Gly Met Ala
 145 150 155 160
 Pro Ala Leu Gln Pro Thr Gln Gly Ala Met Pro Ala Phe Ala Ser Ala
 165 170 175
 20 Phe Gln Arg Arg Ala Gly Gly Val Leu Val Ala Ser His Leu Gln Ser
 180 185 190
 Phe Leu Glu Val Ser Tyr Arg Val Leu Arg His Leu Ala Gln Pro
 195 200 205

 <210> 40
 25 <211> 989
 <212> PRT
 30 <213> Homo sapiens

 <400> 40
 35 Met Lys Val Val Asn Leu Lys Gln Ala Ile Leu Gln Ala Trp Lys Glu
 1 5 10 15
 Arg Trp Ser Tyr Tyr Gln Trp Ala Ile Asn Met Lys Lys Phe Phe Pro
 20 25 30
 Lys Gly Ala Thr Trp Asp Ile Leu Asn Leu Ala Asp Ala Leu Leu Glu
 35 40 45
 40 Gln Ala Met Ile Gly Pro Ser Pro Asn Pro Leu Ile Leu Ser Tyr Leu
 50 55 60
 Lys Tyr Ala Ile Ser Ser Gln Met Val Ser Tyr Ser Ser Val Leu Thr
 65 70 75 80
 Ala Ile Ser Lys Phe Asp Asp Phe Ser Arg Asp Leu Cys Val Gln Ala
 85 90 95
 45 Leu Leu Asp Ile Met Asp Met Phe Cys Asp Arg Leu Ser Cys His Gly
 100 105 110
 Lys Ala Glu Glu Cys Ile Gly Leu Cys Arg Ala Leu Leu Ser Ala Leu
 115 120 125
 50 His Trp Leu Leu Arg Cys Thr Ala Ala Ser Ala Glu Arg Leu Arg Glu
 130 135 140
 Gly Leu Glu Ala Gly Thr Pro Ala Ala Gly Glu Lys Gln Leu Ala Met
 145 150 155 160
 Cys Leu Gln Arg Leu Glu Lys Thr Leu Ser Ser Thr Lys Asn Arg Ala
 165 170 175
 55 Leu Leu His Ile Ala Lys Leu Glu Glu Ala Ser Ser Trp Thr Ala Ile
 180 185 190

Glu His Ser Leu Leu Lys Leu Gly Glu Ile Leu Thr Asn Leu Ser Asn
 195 200 205
 5 Pro Gln Leu Arg Ser Gln Ala Glu Gln Cys Gly Thr Leu Ile Arg Ser
 210 215 220
 Ile Pro Thr Met Leu Ser Val His Ala Glu Gln Met His Lys Thr Gly
 225 230 235 240
 Phe Pro Thr Val His Ala Val Ile Leu Leu Glu Gly Thr Met Asn Leu
 245 250 255
 10 Thr Gly Glu Thr Gln Ser Leu Val Glu Gln Leu Thr Met Val Lys Arg
 260 265 270
 Met Gln His Ile Pro Thr Pro Leu Phe Val Leu Glu Ile Trp Lys Ala
 275 280 285
 Cys Phe Val Gly Leu Ile Glu Ser Pro Glu Gly Thr Glu Glu Leu Lys
 290 295 300
 15 Trp Thr Ala Phe Thr Phe Leu Lys Ile Pro Gln Val Leu Val Lys Leu
 305 310 315 320
 Lys Lys Tyr Ser His Gly Asp Lys Asp Phe Thr Glu Asp Val Asn Cys
 325 330 335
 Ala Phe Glu Phe Leu Leu Lys Leu Thr Pro Leu Leu Asp Lys Ala Asp
 340 345 350
 20 Gln Arg Cys Asn Cys Asp Cys Thr Asn Phe Leu Leu Gln Glu Cys Gly
 355 360 365
 Lys Gln Gly Leu Leu Ser Glu Ala Ser Val Asn Asn Leu Met Ala Lys
 370 375 380
 Arg Lys Ala Asp Arg Glu His Ala Pro Gln Gln Lys Ser Gly Glu Asn
 385 390 395 400
 25 Ala Asn Ile Gln Pro Asn Ile Gln Leu Ile Leu Arg Ala Glu Pro Thr
 405 410 415
 Val Thr Asn Ile Leu Lys Thr Met Asp Ala Asp His Ser Lys Ser Pro
 420 425 430
 30 Glu Gly Leu Leu Gly Val Leu Gly His Met Leu Ser Gly Lys Ser Leu
 435 440 445
 Asp Leu Leu Leu Ala Ala Ala Ala Thr Gly Lys Leu Lys Ser Phe
 450 455 460
 Ala Arg Lys Phe Ile Asn Leu Asn Glu Phe Thr Thr Tyr Gly Ser Glu
 465 470 475 480
 35 Glu Ser Thr Lys Pro Ala Ser Val Arg Ala Leu Leu Phe Asp Ile Ser
 485 490 495
 Phe Leu Met Leu Cys His Val Ala Gln Thr Tyr Gly Ser Glu Val Ile
 500 505 510
 Leu Ser Glu Ser Arg Thr Gly Ala Glu Val Pro Phe Phe Glu Thr Trp
 515 520 525
 40 Met Gln Thr Cys Met Pro Glu Glu Gly Lys Ile Leu Asn Pro Asp His
 530 535 540
 Pro Cys Phe Arg Pro Asp Ser Thr Lys Val Glu Ser Leu Val Ala Leu
 545 550 555 560
 Leu Asn Asn Ser Ser Glu Met Lys Leu Val Gln Met Lys Trp His Glu
 565 570 575
 45 Ala Cys Leu Ser Ile Ser Ala Ala Ile Leu Glu Ile Leu Asn Ala Trp
 580 585 590
 Glu Asn Gly Val Leu Ala Phe Glu Ser Ile Gln Lys Ile Thr Asp Asn
 595 600 605
 50 Ile Lys Gly Lys Val Cys Ser Leu Ala Val Cys Ala Val Ala Trp Leu
 610 615 620
 Val Ala His Val Arg Met Leu Gly Leu Asp Glu Arg Glu Lys Ser Leu
 625 630 635 640
 Gln Met Ile Arg Gln Leu Ala Gly Pro Leu Phe Ser Glu Asn Thr Leu
 645 650 655

5 Gln Phe Tyr Asn Glu Arg Val Val Ile Met Asn Ser Ile Leu Glu Arg
 660 665 670
 Met Cys Ala Asp Val Leu Gln Gln Thr Ala Thr Gln Ile Lys Phe Pro
 675 680 685
 Ser Thr Gly Val Asp Thr Met Pro Tyr Trp Asn Leu Leu Pro Pro Lys
 690 700
 Arg Pro Ile Lys Glu Val Leu Thr Asp Ile Phe Ala Lys Val Leu Glu
 705 710 715 720
 10 Lys Gly Trp Val Asp Ser Arg Ser Ile His Ile Phe Asp Thr Leu Leu
 725 730 735
 His Met Gly Gly Val Tyr Trp Phe Cys Asn Asn Leu Ile Lys Glu Leu
 740 745 750 755
 Leu Lys Glu Thr Arg Lys Glu His Thr Leu Arg Ala Val Glu Leu Leu
 760 765 770
 15 Tyr Ser Ile Phe Cys Leu Asp Met Gln Gln Val Thr Leu Val Leu Leu
 775 780 785
 Gly His Ile Leu Pro Gly Leu Leu Thr Asp Ser Ser Lys Trp His Ser
 790 795 800
 Leu Met Asp Pro Pro Gly Thr Ala Leu Ala Lys Leu Ala Val Trp Cys
 805 810 815
 20 Ala Leu Ser Ser Tyr Ser Ser His Lys Gly Gln Ala Ser Thr Arg Gln
 820 825 830
 Lys Lys Arg His Arg Glu Asp Ile Glu Asp Tyr Ile Ser Leu Phe Pro
 835 840 845
 25 Leu Asp Asp Val Gln Pro Ser Lys Leu Met Arg Leu Leu Ser Ser Asn
 850 855 860
 Glu Asp Asp Ala Asn Ile Leu Ser Ser Pro Thr Asp Arg Ser Met Ser
 865 870 875 880
 Ser Ser Leu Ser Ala Ser Gln Leu His Thr Val Asn Met Arg Asp Pro
 885 890 895
 30 Leu Asn Arg Val Leu Ala Asn Leu Phe Leu Leu Ile Ser Ser Ile Leu
 900 905 910
 Gly Ser Arg Thr Ala Gly Pro His Thr Gln Phe Val Gln Trp Phe Met
 915 920 925
 Glu Glu Cys Val Asp Cys Leu Glu Gln Gly Gly Arg Gly Ser Val Leu
 930 935 940
 35 Gln Phe Met Pro Phe Thr Thr Val Ser Glu Leu Val Lys Val Ser Ala
 945 950 955 960
 Met Ser Ser Pro Lys Val Val Leu Ala Ile Thr Asp Leu Ser Leu Pro
 965 970 975
 Leu Gly Arg Gln Val Ala Ala Lys Ala Ile Ala Ala Leu
 980 985
 40 <210> 41
 <211> 490
 45 <212> PRT
 <213> Homo sapiens
 50 <400> 41
 Met Glu Gln Lys Pro Ser Lys Val Glu Cys Gly Ser Asp Pro Glu Glu
 1 5 10 15
 Asn Ser Ala Arg Ser Pro Asp Gly Lys Arg Lys Asn Gly Gln
 20 25 30

Cys Ser Leu Lys Thr Ser Met Ser Gly Tyr Ile Pro Ser Tyr Leu Asp
 35 40
 Lys Asp Glu Gln Cys Val Val Cys Gly Asp Lys Ala Thr Gly Tyr His
 50 55
 Tyr Arg Cys Ile Thr Cys Glu Gly Cys Lys Gly Phe Phe Arg Arg Thr
 65 70 75
 Ile Gln Lys Asn Leu His Pro Thr Tyr Ser Cys Lys Tyr Asp Ser Cys
 85 90 95
 Cys Val Ile Asp Lys Ile Thr Arg Asn Gln Cys Gln Leu Cys Arg Phe
 100 105 110
 Lys Lys Cys Ile Ala Val Gly Met Ala Met Asp Leu Val Leu Asp Asp
 115 120 125
 Ser Lys Arg Val Ala Lys Arg Lys Leu Ile Glu Gln Asn Arg Glu Arg
 130 135 140
 Arg Arg Lys Glu Glu Met Ile Arg Ser Leu Gln Gln Arg Pro Glu Pro
 145 150 155
 Thr Pro Glu Glu Trp Asp Leu Ile His Ile Ala Thr Glu Ala His Arg
 165 170 175
 Ser Thr Asn Ala Gln Gly Ser His Trp Lys Gln Arg Arg Lys Phe Leu
 180 185 190
 Pro Asp Asp Ile Gly Gln Ser Pro Ile Val Ser Met Pro Asp Gly Asp
 195 200 205
 Lys Val Asp Leu Glu Ala Phe Ser Glu Phe Thr Lys Ile Ile Thr Pro
 210 215 220
 Ala Ile Thr Arg Val Val Asp Phe Ala Lys Lys Leu Pro Met Phe Ser
 225 230 235
 Glu Leu Pro Cys Glu Asp Gln Ile Ile Leu Leu Lys Gly Cys Cys Met
 245 250 255
 Glu Ile Met Ser Leu Arg Ala Ala Val Arg Tyr Asp Pro Glu Ser Asp
 260 265 270
 Thr Leu Thr Leu Ser Gly Glu Met Ala Val Lys Arg Glu Gln Leu Lys
 275 280 285
 Asn Gly Gly Leu Gly Val Val Ser Asp Ala Ile Phe Glu Leu Gly Lys
 290 295 300
 Ser Leu Ser Ala Phe Asn Leu Asp Asp Thr Glu Val Ala Leu Leu Gln
 305 310 315
 Ala Val Leu Leu Met Ser Thr Asp Arg Ser Gly Leu Leu Cys Val Asp
 325 330 335
 Lys Ile Glu Lys Ser Gln Glu Ala Tyr Leu Leu Ala Phe Glu His Tyr
 340 345 350
 Val Asn His Arg Lys His Asn Ile Pro His Phe Trp Pro Lys Leu Leu
 355 360 365
 Met Lys Glu Arg Glu Val Gln Ser Ser Ile Leu Tyr Lys Gly Ala Ala
 370 375 380
 Ala Glu Gly Arg Pro Gly Gly Ser Leu Gly Val His Pro Glu Gly Gln
 385 390 395
 Gln Leu Leu Gly Met His Val Val Gln Gly Pro Gln Val Arg Gln Leu
 405 410 415
 Glu Gln Gln Leu Gly Glu Ala Gly Ser Leu Gln Gly Pro Val Leu Gln
 420 425 430
 His Gln Ser Pro Lys Ser Pro Gln Gln Arg Leu Leu Glu Leu Leu His
 435 440 445
 Arg Ser Gly Ile Leu His Ala Arg Ala Val Cys Gly Glu Asp Asp Ser
 450 455 460
 Ser Glu Ala Asp Ser Pro Ser Ser Ser Glu Glu Pro Glu Val Cys
 465 470 475 480
 Glu Asp Leu Ala Gly Asn Ala Ala Ser Pro
 485 490

<210> 42

<211> 614

<212> PRT

<213> Homo sapiens

<400> 42

```

Met Thr Thr Leu Asp Ser Asn Asn Asn Thr Gly Gly Val Ile Thr Tyr
1      5      10      15
Ile Gly Ser Ser Gly Ser Ser Pro Ser Arg Thr Ser Pro Glu Ser Leu
15     20     25     30
Tyr Ser Asp Asn Ser Asn Gly Ser Phe Gln Ser Leu Thr Gln Gly Cys
35     40     45
Pro Thr Tyr Phe Pro Pro Ser Pro Thr Gly Ser Leu Thr Gln Asp Pro
50     55     60
Ala Arg Ser Phe Gly Ser Ile Pro Pro Ser Leu Ser Asp Asp Gly Ser
65     70     75     80
Pro Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Phe Tyr Asn
85     90     95
Gly Ser Pro Pro Gly Ser Leu Gln Val Ala Met Glu Asp Ser Ser Arg
100    105    110
Val Ser Pro Ser Lys Ser Thr Ser Asn Ile Thr Lys Leu Asn Gly Met
115    120    125
Val Leu Leu Cys Lys Val Cys Gly Asp Val Ala Ser Gly Phe His Tyr
130    135    140
Gly Val Leu Ala Cys Glu Gly Cys Lys Gly Phe Phe Arg Arg Ser Ile
145    150    155    160
Gln Gln Asn Ile Gln Tyr Lys Arg Cys Leu Lys Asn Glu Asn Cys Ser
165    170    175
Ile Val Arg Ile Asn Arg Asn Arg Cys Gln Gln Cys Arg Phe Lys Lys
180    185    190
Cys Leu Ser Val Gly Met Ser Arg Asp Ala Val Arg Phe Gly Arg Ile
195    200    205
Pro Lys Arg Glu Lys Gln Arg Met Leu Ala Glu Met Gln Ser Ala Met
210    215    220
Asn Leu Ala Asn Asn Gln Leu Ser Ser Gln Cys Pro Leu Glu Thr Ser
225    230    235    240
Pro Thr Gln His Pro Thr Pro Gly Pro Met Gly Pro Ser Pro Pro Pro
245    250    255
Ala Pro Val Pro Ser Pro Leu Val Gly Phe Ser Gln Phe Pro Gln Gln
260    265    270
Leu Thr Pro Pro Arg Ser Pro Ser Pro Glu Pro Thr Val Glu Asp Val
275    280    285
Ile Ser Gln Val Ala Arg Ala His Arg Glu Ile Phe Thr Tyr Ala His
290    295    300
Asp Lys Leu Gly Ser Ser Pro Gly Asn Phe Asn Ala Asn His Ala Ser
305    310    315    320
Gly Ser Pro Pro Ala Thr Thr Pro His Arg Trp Glu Asn Gln Gly Cys
325    330    335
Pro Pro Ala Pro Asn Asp Asn Asn Thr Leu Ala Ala Gln Arg His Asn
340    345    350
Glu Ala Leu Asn Gly Leu Arg Gln Ala Pro Ser Ser Tyr Pro Pro Thr
355    360    365

```


	Trp	Pro	Pro	Gly	Pro	Ala	His	His	Ser	Cys	His	Gln	Ser	Asn	Ser	Asn
	370						375					380				
5	Gly	His	Arg	Leu	Cys	Pro	Thr	His	Val	Tyr	Ala	Ala	Pro	Glu	Gly	Lys
	385					390					395					400
	Ala	Pro	Ala	Asn	Ser	Pro	Arg	Gln	Gly	Asn	Ser	Lys	Asn	Val	Leu	Leu
					405					410					415	
	Ala	Cys	Pro	Met	Asn	Met	Tyr	Pro	His	Gly	Arg	Ser	Gly	Arg	Thr	Val
				420						425				430		
10	Gln	Glu	Ile	Trp	Glu	Asp	Phe	Ser	Met	Ser	Phe	Thr	Pro	Ala	Val	Arg
				435				440					445			
	Glu	Val	Val	Glu	Phe	Ala	Lys	His	Ile	Pro	Gly	Phe	Arg	Asp	Leu	Ser
	450					455						460				
	Gln	His	Asp	Gln	Val	Thr	Leu	Lys	Ala	Gly	Thr	Phe	Glu	Val	Leu	Leu
	465					470					475				480	
15	Met	Val	Arg	Phe	Ala	Ser	Leu	Phe	Asn	Val	Lys	Asp	Gln	Thr	Val	Met
				485					490						495	
	Phe	Leu	Ser	Arg	Thr	Thr	Tyr	Ser	Leu	Gln	Glu	Leu	Gly	Ala	Met	Gly
				500				505						510		
20	Met	Gly	Asp	Leu	Leu	Ser	Ala	Met	Phe	Asp	Phe	Ser	Glu	Lys	Leu	Asn
			515					520					525			
	Ser	Leu	Ala	Leu	Thr	Glu	Glu	Glu	Leu	Gly	Leu	Phe	Thr	Ala	Val	Val
	530					535						540				
	Leu	Val	Ser	Ala	Asp	Arg	Ser	Gly	Met	Glu	Ala	Ser	Ser	Ala	Ser	Glu
	545					550					555				560	
25	Gln	Leu	Gln	Glu	Thr	Leu	Leu	Arg	Ala	Leu	Arg	Ala	Leu	Val	Leu	Lys
					565					570					575	
	Asn	Arg	Pro	Leu	Glu	Thr	Ser	Arg	Phe	Thr	Lys	Leu	Leu	Leu	Lys	Leu
				580					585					590		
	Pro	Asp	Leu	Arg	Thr	Leu	Asn	Asn	Met	His	Ser	Glu	Lys	Leu	Leu	Ser
				595				600					605			
30	Phe	Arg	Val	Asp	Ala	Gln										
				610												
	<210>						43									
	<211>						703									
35	<212>						PRT									
	<213>						Homo sapiens									
40																
	<400>						43									
	Met	Ala	Asp	Arg	Arg	Arg	Gln	Arg	Ala	Ser	Gln	Asp	Thr	Glu	Asp	Glu
	1				5					10					15	
45	Glu	Ser	Gly	Ala	Ser	Gly	Ser	Asp	Ser	Gly	Gly	Ser	Pro			

Val Glu Leu Lys Ser Glu Ala Asn Asp Ala Val Asn Ser Ser Thr Lys
 115 120 125
 Glu Glu Lys Gly Glu Glu Lys Pro Asp Thr Lys Ser Thr Val Thr Gly
 130 135 140
 Glu Arg Gln Ser Gly Asp Gly Gln Glu Ser Thr Glu Pro Val Glu Asn
 145 150 155 160
 Lys Val Gly Lys Lys Gly Pro Lys His Leu Asp Asp Asp Glu Asp Arg
 165 170 175
 Lys Asn Pro Ala Tyr Ile Pro Arg Lys Gly Leu Phe Phe Glu His Asp
 180 185 190
 Leu Arg Gly Gln Thr Gln Glu Glu Glu Val Arg Pro Lys Gly Arg Gln
 195 200 205
 Arg Lys Leu Trp Lys Asp Glu Gly Arg Trp Glu His Asp Lys Phe Arg
 210 215 220
 Glu Asp Glu Gln Ala Pro Lys Ser Arg Gln Glu Leu Ile Ala Leu Tyr
 225 230 235 240
 Gly Tyr Asp Ile Arg Ser Ala His Asn Pro Asp Asp Ile Lys Pro Arg
 245 250 255
 Arg Ile Arg Lys Pro Arg Tyr Gly Ser Pro Gln Arg Asp Pro Asn
 260 265 270
 Trp Asn Gly Glu Arg Leu Asn Lys Ser His Arg His Gln Gly Leu Gly
 275 280 285
 Gly Thr Leu Pro Pro Arg Thr Phe Ile Asn Arg Asn Ala Ala Gly Thr
 290 295 300
 Gly Arg Met Ser Ala Pro Arg Asn Tyr Ser Arg Ser Gly Gly Phe Lys
 305 310 315 320
 Glu Gly Arg Ala Gly Phe Arg Pro Val Glu Ala Gly Gly Gln His Gly
 325 330 335
 Gly Arg Ser Gly Glu Thr Val Lys His Glu Ile Ser Tyr Arg Ser Arg
 340 345 350
 Arg Leu Glu Gln Thr Ser Val Arg Asp Pro Ser Glu Ala Asp Ala
 355 360 365
 Pro Val Leu Gly Ser Pro Glu Lys Glu Glu Ala Ala Ser Glu Pro Pro
 370 375 380
 Ala Ala Ala Pro Asp Ala Ala Pro Pro Pro Pro Asp Arg Pro Ile Glu
 385 390 395 400
 Lys Lys Ser Tyr Ser Arg Ala Arg Arg Thr Arg Thr Lys Val Gly Asp
 405 410 415
 Ala Val Lys Leu Ala Glu Glu Val Pro Pro Pro Glu Gly Leu Ile
 420 425 430
 Pro Ala Pro Pro Val Pro Glu Thr Thr Pro Thr Pro Thr Lys Thr
 435 440 445
 Gly Thr Trp Glu Ala Pro Val Asp Ser Ser Thr Ser Gly Leu Glu Gln
 450 455 460
 Asp Val Ala Gln Leu Asn Ile Ala Glu Gln Asn Trp Ser Pro Gly Gln
 465 470 475 480
 Pro Ser Phe Leu Gln Pro Arg Glu Leu Arg Gly Met Pro Asn His Ile
 485 490 495
 His Met Gly Ala Gly Pro Pro Pro Gln Phe Asn Arg Met Glu Glu Met
 500 505 510
 Gly Val Gln Gly Gly Arg Ala Lys Arg Tyr Ser Ser Gln Arg Gln Arg
 515 520 525
 Pro Val Pro Glu Pro Pro Ala Pro Pro Val His Ile Ser Ile Met Glu
 530 535 540
 Gly His Tyr Tyr Asp Pro Leu Gln Phe Gln Gly Pro Ile Tyr Thr His
 545 550 555 560
 Gly Asp Ser Pro Ala Pro Leu Pro Pro Gln Gly Met Leu Val Gln Pro
 565 570 575

Gly Met Asn Leu Pro His Pro Gly Leu His Pro His Gln Thr Pro Ala
 580 585 590
 Pro Leu Pro Asn Pro Gly Leu Tyr Pro Pro Pro Val Ser Met Ser Pro
 595 600 605
 Gly Gln Pro Pro Pro Gln Gln Leu Leu Ala Pro Thr Tyr Phe Ser Ala
 610 615 620
 Pro Gly Val Met Asn Phe Gly Asn Pro Ser Tyr Pro Tyr Ala Pro Gly
 625 630 635 640
 Ala Leu Pro Pro Pro Pro Pro His Leu Tyr Pro Asn Thr Gln Ala
 645 650 655
 Pro Ser Gln Val Tyr Gly Gly Val Thr Tyr Tyr Asn Pro Ala Gln Gln
 660 665 670
 Gln Val Gln Pro Lys Pro Ser Pro Pro Arg Arg Thr Pro Gln Pro Val
 675 685
 Thr Ile Lys Pro Pro Pro Pro Glu Val Val Ser Arg Gly Ser Ser
 690 695 700

<210> 44

<211> 560

<212> PRT

<213> Homo sapiens

<400> 44

Met Pro Gln Thr Arg Ser Gln Ala Gln Ala Thr Ile Ser Phe Pro Lys
 1 5 10 15
 Arg Lys Leu Ser Arg Ala Leu Asn Lys Ala Lys Asn Ser Ser Asp Ala
 20 25 30
 Lys Leu Glu Pro Thr Asn Val Gln Thr Val Thr Cys Ser Pro Arg Val
 35 40 45
 Lys Ala Leu Pro Leu Ser Pro Arg Lys Arg Leu Gly Asp Asp Asn Leu
 50 55 60
 Cys Asn Thr Pro His Leu Pro Pro Cys Ser Pro Pro Lys Gln Gly Lys
 65 70 75 80
 Lys Glu Asn Gly Pro Pro His Ser His Thr Leu Lys Gly Arg Arg Leu
 85 90 95
 Val Phe Asp Asn Gln Leu Thr Ile Lys Ser Pro Ser Lys Arg Glu Leu
 100 105 110
 Ala Lys Val His Gln Asn Lys Ile Leu Ser Ser Val Arg Lys Ser Gln
 115 120 125
 Glu Ile Thr Thr Asn Ser Glu Gln Arg Cys Pro Leu Lys Lys Glu Ser
 130 135 140
 Ala Cys Val Arg Leu Phe Lys Gln Glu Gly Thr Cys Tyr Gln Gln Ala
 145 150 155 160
 Lys Leu Val Leu Asn Thr Ala Val Pro Asp Arg Leu Pro Ala Arg Glu
 165 170 175
 Arg Glu Met Asp Val Ile Arg Asn Phe Leu Arg Glu His Ile Cys Gly
 180 185 190
 Lys Lys Ala Gly Ser Leu Tyr Leu Ser Gly Ala Pro Gly Thr Gly Lys
 195 200 205
 Thr Ala Cys Leu Ser Arg Ile Leu Gln Asp Leu Lys Lys Glu Leu Lys
 210 215 220
 Gly Phe Lys Thr Ile Met Leu Asn Cys Met Ser Leu Arg Thr Ala Gln
 225 230 235 240

Ala Val Phe Pro Ala Ile Ala Gln Glu Ile Cys Gln Glu Glu Val Ser
 245 255
 Arg Pro Ala Gly Lys Asp Met Met Arg Lys Leu Glu Lys His Met Thr
 260 265 270
 Ala Glu Lys Gly Pro Met Ile Val Leu Val Leu Asp Glu Met Asp Gln
 275 280 285
 Leu Asp Ser Ser Lys Gly Gln Asp Val Leu Tyr Thr Leu Phe Glu Trp Pro
 290 295 300
 Trp Leu Ser Asn Ser His Leu Val Leu Ile Gly Ile Ala Asn Thr Leu
 305 310 315 320
 Asp Leu Thr Asp Arg Ile Leu Pro Arg Leu Gln Ala Arg Glu Lys Cys
 325 330 335
 Lys Pro Gln Leu Leu Asn Phe Pro Pro Tyr Thr Arg Asn Gln Ile Val
 340 345 350
 Thr Ile Leu Gln Asp Arg Leu Asn Gln Val Ser Arg Asp Gln Val Leu
 355 360 365
 Asp Asn Ala Ala Val Gln Phe Cys Ala Arg Lys Val Ser Ala Val Ser
 370 375 380
 Gly Asp Val Arg Lys Ala Leu Asp Val Cys Arg Arg Ala Ile Glu Ile
 385 390 395 400
 Val Glu Ser Asp Val Lys Ser Gln Thr Ile Leu Lys Pro Leu Ser Glu
 405 410 415
 Cys Lys Ser Pro Ser Glu Pro Leu Ile Pro Lys Arg Val Gly Leu Ile
 420 425 430
 His Ile Ser Gln Val Ile Ser Glu Val Asp Gly Asn Arg Met Thr Leu
 435 440 445
 Ser Gln Glu Gly Ala Gln Asp Ser Phe Pro Leu Gln Gln Lys Ile Leu
 450 455 460
 Val Cys Ser Leu Met Leu Leu Ile Arg Gln Leu Lys Ile Lys Glu Val
 465 470 475 480
 Thr Leu Gly Lys Leu Tyr Glu Ala Tyr Ser Lys Val Cys Arg Lys Gln
 485 490 495
 Gln Val Ala Ala Val Asp Gln Ser Glu Cys Leu Ser Leu Ser Gly Leu
 500 505 510
 Leu Glu Ala Arg Gly Ile Leu Gly Leu Lys Arg Asn Lys Glu Thr Arg
 515 520 525
 Leu Thr Lys Val Phe Phe Lys Ile Glu Glu Lys Glu Ile Glu His Ala
 530 535 540
 Leu Lys Asp Lys Ala Leu Ile Gly Asn Ile Leu Ala Thr Gly Leu Pro
 545 550 555 560

 <210> 45

 <211> 462

 <212> PRT

 <213> Homo sapiens

 <400> 45
 Met Ala Ser Asn Ser Ser Ser Cys Pro Thr Pro Gly Gly Gly His Leu
 1 5 10 15
 Asn Gly Tyr Pro Val Pro Pro Tyr Ala Phe Phe Phe Pro Pro Met Leu
 20 25 30
 Gly Gly Leu Ser Pro Pro Gly Ala Leu Thr Thr Leu Gln His Gln Leu
 35 40 45

Pro Val Ser Gly Tyr Ser Thr Pro Ser Pro Ala Thr Ile Glu Thr Gln
 50 55 60
 Ser Ser Ser Ser Glu Glu Ile Val Pro Ser Pro Pro Ser Pro Pro Pro
 65 70 75 80
 Leu Pro Arg Ile Tyr Lys Pro Cys Phe Val Cys Gln Asp Lys Ser Ser
 85 90 95
 Gly Tyr His Tyr Gly Val Ser Ala Cys Glu Gly Cys Lys Gly Phe Phe
 100 105 110
 Arg Arg Ser Ile Gln Lys Asn Met Val Tyr Thr Cys His Arg Asp Lys
 115 120 125
 Asn Cys Ile Ile Asn Lys Val Thr Arg Asn Arg Cys Gln Tyr Cys Arg
 130 135 140
 Leu Gln Lys Cys Phe Glu Val Gly Met Ser Lys Glu Ser Val Arg Asn
 145 150 155 160
 Asp Arg Asn Lys Lys Lys Lys Glu Val Pro Lys Pro Glu Cys Ser Glu
 165 170 175
 Ser Tyr Thr Leu Thr Pro Glu Val Gly Glu Leu Ile Glu Lys Val Arg
 180 185 190
 Lys Ala His Gln Glu Thr Phe Pro Ala Leu Cys Gln Leu Gly Lys Tyr
 195 200 205
 Thr Thr Asn Asn Ser Ser Glu Gln Arg Val Ser Leu Asp Ile Asp Leu
 210 215 220
 Trp Asp Lys Phe Ser Glu Leu Ser Thr Lys Cys Ile Ile Lys Thr Val
 225 230 235 240
 Glu Phe Ala Lys Gln Leu Pro Gly Phe Thr Thr Leu Thr Ile Ala Asp
 245 250 255
 Gln Ile Thr Leu Leu Lys Ala Ala Cys Leu Asp Ile Leu Ile Leu Arg
 260 265 270
 Ile Cys Thr Arg Tyr Thr Pro Glu Gln Asp Thr Met Thr Phe Ser Asp
 275 280 285
 Gly Leu Thr Leu Asn Arg Thr Gln Met His Asn Ala Gly Phe Gly Pro
 290 295 300
 Leu Thr Asp Leu Val Phe Ala Phe Ala Asn Gln Leu Leu Pro Leu Glu
 305 310 315 320
 Met Asp Asp Ala Glu Thr Gly Leu Leu Ser Ala Ile Cys Leu Ile Cys
 325 330 335
 Gly Asp Arg Gln Asp Leu Glu Gln Pro Asp Arg Val Asp Met Leu Gln
 340 345 350
 Glu Pro Leu Leu Glu Ala Leu Lys Val Tyr Val Arg Lys Arg Arg Pro
 355 360 365
 Ser Arg Pro His Met Phe Pro Lys Met Leu Met Lys Ile Thr Asp Leu
 370 375 380
 Arg Ser Ile Ser Ala Lys Gly Ala Glu Arg Val Ile Thr Leu Lys Met
 385 390 395 400
 Glu Ile Pro Gly Ser Met Pro Pro Leu Ile Gln Glu Met Leu Glu Asn
 405 410 415
 Ser Glu Gly Leu Asp Thr Leu Ser Gly Gln Pro Gly Gly Gly Arg
 420 425 430
 Asp Gly Gly Gly Leu Ala Pro Pro Gly Ser Cys Ser Pro Ser Leu
 435 440 445
 Ser Pro Ser Ser Asn Arg Ser Ser Pro Ala Thr His Ser Pro
 450 455 460

<210> 46

<211> 1531

<212> PRT

<213> Homo sapiens

<400> 46

```

Met Glu Val Ser Pro Leu Gln Pro Val Asn Glu Asn Met Gln Val Asn
1      5      10      15
Lys Ile Lys Lys Asn Glu Asp Ala Lys Lys Arg Leu Ser Val Glu Arg
15     20     25     30
Ile Tyr Gln Lys Lys Thr Gln Leu Glu His Ile Leu Leu Arg Pro Asp
35     40     45
Thr Tyr Ile Gly Ser Val Glu Leu Val Thr Gln Gln Met Trp Val Tyr
50     55     60
Asp Glu Asp Val Gly Ile Asn Tyr Arg Glu Val Thr Phe Val Pro Gly
65     70     75     80
Leu Tyr Lys Ile Phe Asp Glu Ile Leu Val Asn Ala Ala Asp Asn Lys
85     90     95
Gln Arg Asp Pro Lys Met Ser Cys Ile Arg Val Thr Ile Asp Pro Glu
100    105    110
Asn Asn Leu Ile Ser Ile Trp Asn Asn Gly Lys Gly Ile Pro Val Val
115    120    125
Glu His Lys Val Glu Lys Met Tyr Val Pro Ala Leu Ile Phe Gly Gln
130    135    140
Leu Leu Thr Ser Ser Asn Tyr Asp Asp Asp Glu Lys Lys Val Thr Gly
145    150    155    160
Gly Arg Asn Gly Tyr Gly Ala Lys Leu Cys Asn Ile Phe Ser Thr Lys
165    170    175
Phe Thr Val Glu Thr Ala Ser Arg Glu Tyr Lys Lys Met Phe Lys Gln
180    185    190
Thr Trp Met Asp Asn Met Gly Arg Ala Gly Glu Met Glu Leu Lys Pro
195    200    205
Phe Asn Gly Glu Asp Tyr Thr Cys Ile Thr Phe Gln Pro Asp Leu Ser
210    215    220
Lys Phe Lys Met Gln Ser Leu Asp Lys Asp Ile Val Ala Leu Met Val
225    230    235
Arg Arg Ala Tyr Asp Ile Ala Gly Ser Thr Lys Asp Val Lys Val Phe
240    245    250    255
Leu Asn Gly Asn Lys Leu Pro Val Lys Gly Phe Arg Ser Tyr Val Asp
260    265    270
Met Tyr Leu Lys Asp Lys Leu Asp Glu Thr Gly Asn Ser Leu Lys Val
275    280    285
Ile His Glu Gln Val Asn His Arg Trp Glu Val Cys Leu Thr Met Ser
290    295    300
Glu Lys Gly Phe Gln Gln Ile Ser Phe Val Asn Ser Ile Ala Thr Ser
305    310    315    320
Lys Gly Gly Arg His Val Asp Tyr Val Ala Asp Gln Ile Val Thr Lys
325    330    335
Leu Val Asp Val Val Lys Lys Lys Asn Lys Gly Gly Val Ala Val Lys
340    345    350
Ala His Gln Val Lys Asn His Met Trp Ile Phe Val Asn Ala Leu Ile
355    360    365
Glu Asn Pro Thr Phe Asp Ser Gln Thr Lys Glu Asn Met Thr Leu Gln
370    375    380

```

Pro Lys Ser Phe Gly Ser Thr Cys Gln Leu Ser Glu Lys Phe Ile Lys
 385 390 395 400
 Ala Ala Ile Gly Cys Gly Ile Val Glu Ser Ile Leu Asn Trp Val Lys
 405 410 415
 Phe Lys Ala Gln Val Gln Leu Asn Lys Lys Cys Ser Ala Val Lys His
 420 425 430
 Asn Arg Ile Lys Gly Ile Pro Lys Leu Asp Asp Ala Asn Asp Ala Gly
 435 440 445
 Gly Arg Asn Ser Thr Glu Cys Thr Leu Ile Leu Thr Glu Gly Asp Ser
 450 455 460
 Ala Lys Thr Leu Ala Val Ser Gly Leu Gly Val Val Gly Arg Asp Lys
 465 470 475
 Tyr Gly Val Phe Pro Leu Arg Gly Lys Ile Leu Asn Val Arg Glu Ala
 485 490 495
 Ser His Lys Gln Ile Met Glu Asn Ala Glu Ile Asn Asn Ile Ile Lys
 500 505 510
 Ile Val Gly Leu Gln Tyr Lys Lys Asn Tyr Glu Asp Glu Asp Ser Leu
 515 520 525
 Lys Thr Leu Arg Tyr Gly Lys Ile Met Ile Met Thr Asp Gln Asp Gln
 530 535 540
 Asp Gly Ser His Ile Lys Gly Leu Leu Ile Asn Phe Ile His His Asn
 545 550 555 560
 Trp Pro Ser Leu Leu Arg His Arg Phe Leu Glu Glu Phe Ile Thr Pro
 565 570 575
 Ile Val Lys Val Ser Lys Asn Lys Gln Glu Met Ala Phe Tyr Ser Leu
 580 585 590
 Pro Glu Phe Glu Glu Trp Lys Ser Ser Thr Pro Asn His Lys Lys Trp
 595 600 605
 Lys Val Lys Tyr Tyr Lys Gly Leu Gly Thr Ser Thr Ser Lys Glu Ala
 610 615 620
 Lys Gln Tyr Phe Ala Asp Met Lys Arg His Arg Ile Gln Phe Lys Tyr
 625 630 635 640
 Ser Gly Pro Glu Asp Asp Ala Ala Ile Ser Leu Ala Phe Ser Lys Lys
 645 650 655
 Gln Ile Asp Asp Arg Lys Glu Trp Leu Thr Asn Phe Met Glu Asp Arg
 660 665 670
 Arg Gln Arg Lys Leu Leu Gly Leu Pro Glu Asp Tyr Leu Tyr Gly Gln
 675 680 685
 Thr Thr Thr Tyr Leu Thr Tyr Asn Asp Phe Ile Asn Lys Glu Leu Ile
 690 695 700
 Leu Phe Ser Asn Ser Asp Asn Glu Arg Ser Ile Pro Ser Met Val Asp
 705 710 715 720
 Gly Leu Lys Pro Gly Gln Arg Lys Val Leu Phe Thr Cys Phe Lys Arg
 725 730 735
 Asn Asp Lys Arg Glu Val Lys Val Ala Gln Leu Ala Gly Ser Val Ala
 740 745 750
 Glu Met Ser Ser Tyr His His Gly Glu Met Ser Leu Met Met Thr Ile
 755 760 765
 Ile Asn Leu Ala Gln Asn Phe Val Gly Ser Asn Asn Leu Asn Leu Leu
 770 775 780
 Gln Pro Ile Gly Gln Phe Gly Thr Arg Leu His Gly Gly Lys Ser Ser
 785 790 795 800
 Ala Ser Pro Arg Tyr Ile Phe Thr Met Leu Ser Ser Leu Ala Arg Leu
 805 810 815
 Leu Phe Pro Pro Lys Asp Asp His Thr Leu Lys Phe Leu Tyr Asp Asp
 820 825 830
 Asn Gln Arg Val Glu Pro Glu Trp Tyr Ile Pro Ile Ile Pro Met Val
 835 840 845

Leu Ile Asn Gly Ala Glu Gly Ile Gly Thr Gly Trp Ser Cys Lys Ile
 850 855 860
 Pro Asn Phe Asp Val Arg Glu Ile Val Asn Asn Ile Arg Arg Leu Met
 865 870 880
 Asp Gly Glu Glu Pro Leu Pro Met Leu Pro Ser Tyr Lys Asn Phe Lys
 885 890 895
 Gly Thr Ile Glu Glu Leu Ala Pro Asn Gln Tyr Val Ile Ser Gly Glu
 900 905 910
 Val Ala Ile Leu Asn Ser Thr Thr Ile Glu Ile Ser Glu Leu Pro Val
 915 920 925
 Arg Thr Trp Thr Gln Thr Tyr Lys Glu Gln Val Leu Glu Pro Met Leu
 930 935 940
 Asn Gly Thr Glu Lys Thr Pro Pro Leu Ile Thr Asp Tyr Arg Glu Tyr
 945 950 955 960
 His Thr Asp Thr Thr Val Lys Phe Val Val Lys Met Thr Glu Glu Lys
 965 970 975
 Leu Ala Glu Ala Glu Arg Val Gly Leu His Lys Val Phe Lys Leu Gln
 980 985 990
 Thr Ser Leu Thr Cys Asn Ser Met Val Leu Phe Asp His Val Gly Cys
 995 1000 1005
 Leu Lys Lys Tyr Asp Thr Val Leu Asp Ile Leu Arg Asp Phe Phe
 1010 1015 1020
 Glu Leu Arg Leu Lys Tyr Tyr Gly Leu Arg Lys Glu Trp Leu Leu
 1025 1030 1035
 Gly Met Leu Gly Ala Glu Ser Ala Lys Leu Asn Asn Gln Ala Arg
 1040 1045 1050
 Phe Ile Leu Glu Lys Ile Asp Gly Lys Ile Ile Ile Glu Asn Lys
 1055 1060 1065
 Pro Lys Lys Glu Leu Ile Lys Val Leu Ile Gln Arg Gly Tyr Asp
 1070 1075 1080
 Ser Asp Pro Val Lys Ala Trp Lys Glu Ala Gln Lys Val Pro
 1085 1090 1095
 Asp Glu Glu Glu Asn Glu Glu Ser Asp Asn Glu Lys Glu Thr Glu
 1100 1105 1110
 Lys Ser Asp Ser Val Thr Asp Ser Gly Pro Thr Phe Asn Tyr Leu
 1115 1120 1125
 Leu Asp Met Pro Leu Trp Tyr Leu Thr Lys Glu Lys Lys Asp Glu
 1130 1135 1140
 Leu Cys Arg Leu Arg Asn Glu Lys Glu Gln Glu Leu Asp Thr Leu
 1145 1150 1155
 Lys Arg Lys Ser Pro Ser Asp Leu Trp Lys Glu Asp Leu Ala Thr
 1160 1165 1170
 Phe Ile Glu Glu Leu Glu Ala Val Glu Ala Lys Glu Lys Gln Asp
 1175 1180 1185
 Glu Gln Val Gly Leu Pro Gly Lys Gly Gly Lys Ala Lys Gly Lys
 1190 1195 1200
 Lys Thr Gln Met Ala Glu Val Leu Pro Ser Pro Arg Gly Gln Arg
 1205 1210 1215
 Val Ile Pro Arg Ile Thr Ile Glu Met Lys Ala Glu Ala Glu Lys
 1220 1225 1230
 Lys Asn Lys Lys Lys Ile Lys Asn Glu Asn Thr Glu Gly Ser Pro
 1235 1240 1245
 Gln Glu Asp Gly Val Glu Leu Glu Gly Leu Lys Gln Arg Leu Glu
 1250 1255 1260
 Lys Lys Gln Lys Arg Glu Pro Gly Thr Lys Thr Lys Lys Gln Thr
 1265 1270 1275
 Thr Leu Ala Phe Lys Pro Ile Lys Lys Gly Lys Lys Arg Asn Pro
 1280 1285 1290

Trp Ser Asp Ser Glu Ser Asp Arg Ser Ser Asp Glu Ser Asn Phe
 1295 1300 1305
 Asp Val Pro Pro Arg Glu Thr Glu Pro Arg Arg Ala Ala Thr Lys
 1310 1315 1320
 Thr Lys Phe Thr Met Asp Leu Asp Ser Asp Glu Asp Phe Ser Asp
 1325 1330 1335
 Phe Asp Glu Lys Thr Asp Asp Glu Asp Phe Val Pro Ser Asp Ala
 1340 1345 1350
 Ser Pro Pro Lys Thr Lys Thr Ser Pro Lys Leu Ser Asn Lys Glu
 1355 1360 1365
 Leu Lys Pro Gln Lys Ser Val Val Ser Asp Leu Glu Ala Asp Asp
 1370 1375 1380
 Val Lys Gly Ser Val Pro Leu Ser Ser Ser Pro Pro Ala Thr His
 1385 1390 1395
 Phe Pro Asp Glu Thr Glu Ile Thr Asn Pro Val Pro Lys Lys Asn
 1400 1405 1410
 Val Thr Val Lys Lys Thr Ala Ala Lys Ser Gln Ser Ser Thr Ser
 1415 1420 1425
 Thr Thr Gly Ala Lys Lys Arg Ala Ala Pro Lys Gly Thr Lys Arg
 1430 1435 1440
 Asp Pro Ala Leu Asn Ser Gly Val Ser Gln Lys Pro Asp Pro Ala
 1445 1450 1455
 Lys Thr Lys Asn Arg Arg Lys Arg Lys Pro Ser Thr Ser Asp Asp
 1460 1465 1470
 Ser Asp Ser Asn Phe Glu Lys Ile Val Ser Lys Ala Val Thr Ser
 1475 1480 1485
 Lys Lys Ser Lys Gly Glu Ser Asp Asp Phe His Met Asp Phe Asp
 1490 1495 1500
 Ser Ala Val Ala Pro Arg Ala Lys Ser Val Arg Ala Lys Lys Pro
 1505 1510 1515
 Ile Lys Tyr Leu Glu Glu Ser Asp Glu Asp Asp Leu Phe
 1520 1525 1530

 <210> 47

 <211> 258

 <212> PRT

 <213> Homo sapiens

 <400> 47
 Met Leu Pro Leu Cys Leu Val Ala Ala Leu Leu Leu Ala Ala Gly Pro
 1 5 10 15
 Gly Pro Ser Leu Gly Asp Glu Ala Ile His Cys Pro Pro Cys Ser Glu
 20 25 30
 Glu Lys Leu Ala Arg Cys Arg Pro Pro Val Gly Cys Glu Glu Leu Val
 35 40 45
 Arg Glu Pro Gly Cys Gly Cys Cys Ala Thr Cys Ala Leu Gly Leu Gly
 50 55 60
 Met Pro Cys Gly Val Tyr Thr Pro Arg Cys Gly Ser Gly Leu Arg Cys
 65 70 75 80
 Tyr Pro Pro Arg Gly Val Glu Lys Pro Leu His Thr Leu Met His Gly
 85 90 95
 Gln Gly Val Cys Met Glu Leu Ala Glu Ile Glu Ala Ile Gln Glu Ser
 100 105 110

Leu Gln Pro Ser Asp Lys Asp Glu Gly Asp His Pro Asn Asn Ser Phe
 115 120 125
 Ser Pro Cys Ser Ala His Asp Arg Arg Cys Leu Gln Lys His Phe Ala
 130 135 140
 Lys Ile Arg Asp Arg Ser Thr Ser Gly Gly Lys Met Lys Val Asn Gly
 145 150 155 160
 Ala Pro Arg Glu Asp Ala Arg Pro Val Pro Gln Gly Ser Cys Gln Ser
 165 175
 Glu Leu His Arg Ala Leu Glu Arg Leu Ala Ala Ser Gln Ser Arg Thr
 180 185 190
 His Glu Asp Leu Tyr Ile Ile Pro Ile Pro Asn Cys Asp Arg Asn Gly
 195 200 205
 Asn Phe His Pro Lys Gln Cys His Pro Ala Leu Asp Gly Gln Arg Gly
 210 215 220
 Lys Cys Trp Cys Val Asp Arg Lys Thr Gly Val Lys Leu Pro Gly Gly
 225 230 235 240
 Leu Glu Pro Lys Gly Glu Leu Asp Cys His Gln Leu Ala Asp Ser Phe
 245 250 255
 Arg Glu

<210> 48

<211> 378

<212> PRT

<213> Homo sapiens

<400> 48

Met Asp Leu Gly Lys Pro Met Lys Ser Val Leu Val Val Ala Leu Leu
 1 5 10 15
 Val Ile Phe Gln Val Cys Leu Cys Gln Asp Glu Val Thr Asp Tyr
 20 25 30
 Ile Gly Asp Asn Thr Thr Val Asp Tyr Thr Leu Phe Glu Ser Leu Cys
 35 40 45
 Ser Lys Lys Asp Val Arg Asn Phe Lys Ala Trp Phe Leu Pro Ile Met
 50 55 60
 Tyr Ser Ile Ile Cys Phe Val Gly Leu Leu Gly Asn Gly Leu Val Val
 65 70 75 80
 Leu Thr Tyr Ile Tyr Phe Lys Arg Leu Lys Thr Met Thr Asp Thr Tyr
 85 90 95
 Leu Leu Asn Leu Ala Val Ala Asp Ile Leu Phe Leu Leu Thr Leu Pro
 100 105 110
 Phe Trp Ala Tyr Ser Ala Ala Lys Ser Trp Val Phe Gly Val His Phe
 115 120 125
 Cys Lys Leu Ile Phe Ala Ile Tyr Lys Met Ser Phe Phe Ser Gly Met
 130 135 140
 Leu Leu Leu Leu Cys Ile Ser Ile Asp Arg Tyr Val Ala Ile Val Gln
 145 150 155 160
 Ala Val Ser Ala His Arg His Arg Ala Arg Val Leu Leu Ile Ser Lys
 165 170 175
 Leu Ser Cys Val Gly Ile Trp Ile Leu Ala Thr Val Leu Ser Ile Pro
 180 185 190
 Glu Leu Leu Tyr Ser Asp Leu Gln Arg Ser Ser Ser Glu Gln Ala Met
 195 200 205

Arg Cys Ser Leu Ile Thr Glu His Val Glu Ala Phe Ile Thr Ile Gln
 210 215 220
 Val Ala Gln Met Val Ile Gly Phe Leu Val Pro Leu Leu Ala Met Ser
 225 230 235 240
 Phe Cys Tyr Leu Val Ile Ile Arg Thr Leu Leu Gln Ala Arg Asn Phe
 245 250 255
 Glu Arg Asn Lys Ala Ile Lys Val Ile Ala Val Val Val Val Phe
 260 265 270
 Ile Val Phe Gln Leu Pro Tyr Asn Gly Val Val Leu Ala Gln Thr Val
 275 280 285
 Ala Asn Phe Asn Ile Thr Ser Ser Thr Cys Glu Leu Ser Lys Gln Leu
 290 295 300
 Asn Ile Ala Tyr Asp Val Thr Tyr Ser Ser Leu Ala Cys Val Arg Cys Cys
 305 310 315 320
 Val Asn Pro Phe Leu Tyr Ala Phe Ile Gly Val Lys Phe Arg Asn Asp
 325 330 335
 Leu Phe Lys Leu Phe Lys Asp Leu Gly Cys Leu Ser Gln Glu Gln Leu
 340 345 350
 Arg Gln Trp Ser Ser Cys Arg His Ile Arg Arg Ser Ser Met Ser Val
 355 360 365
 Glu Ala Glu Thr Thr Thr Thr Phe Ser Pro
 370 375

<210> 49

<211> 411

<212> PRT

<213> Homo sapiens

<400> 49

Met Ser Lys Arg Pro Ser Tyr Ala Pro Pro Thr Pro Ala Pro Ala
 1 5 10 15
 Thr Gln Met Pro Ser Thr Pro Gly Phe Val Gly Tyr Asn Pro Tyr Ser
 20 25 30
 His Leu Ala Tyr Asn Asn Tyr Arg Leu Gly Gly Asn Pro Ser Thr Asn
 35 40 45
 Ser Arg Val Thr Ala Ser Ser Gly Ile Thr Ile Pro Lys Pro Pro Lys
 50 55 60
 Pro Pro Asp Lys Pro Leu Met Pro Tyr Met Arg Tyr Ser Arg Lys Val
 65 70 75 80
 Trp Asp Gln Val Lys Ala Ser Asn Pro Asp Leu Lys Leu Trp Glu Ile
 85 90 95
 Gly Lys Ile Ile Gly Gly Met Trp Arg Asp Leu Thr Asp Glu Glu Lys
 100 105 110
 Gln Glu Tyr Leu Asn Glu Tyr Glu Ala Glu Lys Ile Glu Tyr Asn Glu
 115 120 125
 Ser Met Lys Ala Tyr His Asn Ser Pro Ala Tyr Leu Ala Tyr Ile Asn
 130 135 140
 Ala Lys Ser Arg Ala Glu Ala Ala Leu Glu Glu Glu Ser Arg Gln Arg
 145 150 155 160
 Gln Ser Arg Met Glu Lys Gly Glu Pro Tyr Met Ser Ile Gln Pro Ala
 165 170 175
 Glu Asp Pro Asp Tyr Asp Asp Gly Phe Ser Met Lys His Thr Ala
 180 185 190

Thr Ala Arg Phe Gln Arg Asn His Arg Leu Ile Ser Glu Ile Leu Ser
 195 200 205
 Glu Ser Val Val Pro Asp Val Arg Ser Val Val Thr Thr Ala Arg Met
 210 215 220
 Gln Val Leu Lys Arg Gln Val Gln Ser Leu Met Val His Gln Arg Lys
 225 230 235 240
 Leu Glu Ala Glu Leu Leu Gln Ile Glu Glu Arg His Gln Glu Lys Lys
 245 250 255
 Arg Lys Phe Leu Glu Ser Thr Asp Ser Phe Asn Asn Glu Leu Lys Arg
 260 265 270
 Leu Cys Gly Leu Lys Val Glu Val Asp Met Glu Lys Ile Ala Ala Glu
 275 280 285
 Ile Ala Gln Ala Glu Glu Gln Ala Arg Lys Arg Gln Glu Glu Arg Glu
 290 295 300
 Lys Glu Ala Ala Glu Gln Ala Glu Arg Ser Gln Ser Ser Ile Val Pro
 305 310 315 320
 Glu Glu Glu Gln Ala Ala Asn Lys Gly Glu Glu Lys Lys Asp Asp Glu
 325 330 335
 Asn Ile Pro Met Glu Thr Glu Glu Thr His Leu Glu Glu Thr Thr Glu
 340 345 350
 Ser Gln Gln Asn Gly Glu Glu Gly Thr Ser Thr Pro Glu Asp Lys Glu
 355 360 365
 Ser Gly Gln Glu Gly Val Asp Ser Met Ala Glu Glu Gly Thr Ser Asp
 370 375 380
 Ser Asn Thr Gly Ser Glu Ser Asn Ser Ala Thr Val Glu Glu Pro Pro
 385 390 395 400
 Thr Asp Pro Ile Pro Glu Asp Glu Lys Lys Glu
 405 410

<210> 50

<211> 593

<212> PRT

<213> Homo sapiens

<400> 50

Met Ser Val Arg Tyr Ser Ser Ser Lys His Tyr Ser Ser Ser Arg Ser
 1 5 10 15
 Gly Gly Gly Gly Gly Gly Gly Cys Gly Gly Gly Gly Gly Val Ser
 20 25 30
 Ser Leu Arg Ile Ser Ser Ser Lys Gly Ser Leu Gly Gly Gly Phe Ser
 35 40 45
 Ser Gly Gly Phe Ser Gly Gly Ser Phe Ser Arg Gly Ser Ser Gly Gly
 50 55 60
 Gly Cys Phe Gly Gly Ser Ser Gly Gly Tyr Gly Gly Leu Gly Gly Phe
 65 70 75 80
 Gly Gly Gly Ser Phe His Gly Ser Tyr Gly Ser Ser Ser Phe Gly Gly
 85 90 95
 Ser Tyr Gly Gly Ser Phe Gly Gly Gly Asn Phe Gly Gly Ser Phe
 100 105 110
 Gly Gly Gly Ser Phe Gly Gly Gly Gly Phe Gly Gly Gly Phe Gly
 115 120 125
 Gly Gly Phe Gly Gly Gly Phe Gly Gly Asp Gly Gly Leu Leu Ser Gly
 130 135 140

Asn Glu Lys Val Thr Met Gln Asn Leu Asn Asp Arg Leu Ala Ser Tyr
 145 150 155 160
 Leu Asp Lys Val Arg Ala Leu Glu Glu Ser Asn Tyr Glu Leu Glu Gly
 165 175
 Lys Ile Lys Glu Trp Tyr Glu Lys His Gly Asn Ser His Gln Gly Glu
 180 185 190
 Pro Arg Asp Tyr Ser Lys Tyr Tyr Lys Thr Ile Asp Asp Leu Lys Asn
 195 200 205
 Gln Ile Leu Asn Leu Thr Thr Asp Asn Ala Asn Ile Leu Leu Gln Ile
 210 215 220
 Asp Asn Ala Arg Leu Ala Ala Asp Asp Phe Arg Leu Lys Tyr Glu Asn
 225 230 235 240
 Glu Val Ala Leu Arg Gln Ser Val Glu Ala Asp Ile Asn Gly Leu Arg
 245 255
 Arg Val Leu Asp Glu Leu Thr Leu Thr Lys Ala Asp Leu Glu Met Gln
 260 265 270
 Ile Glu Ser Leu Thr Glu Glu Leu Ala Tyr Leu Lys Lys Asn His Glu
 275 280 285
 Glu Glu Met Lys Asp Leu Arg Asn Val Ser Thr Gly Asp Val Asn Val
 290 295 300
 Glu Met Asn Ala Ala Pro Gly Val Asp Leu Thr Gln Leu Leu Asn Asn
 305 310 315 320
 Met Arg Ser Gln Tyr Glu Gln Leu Ala Glu Gln Asn Arg Lys Asp Ala
 325 330 335
 Glu Ala Trp Phe Asn Glu Lys Ser Lys Glu Leu Thr Thr Glu Ile Asp
 340 345 350
 Asn Asn Ile Glu Gln Ile Ser Ser Tyr Lys Ser Glu Ile Thr Glu Leu
 355 360 365
 Arg Arg Asn Val Gln Ala Leu Glu Ile Glu Leu Gln Ser Gln Leu Ala
 370 375 380
 Leu Lys Gln Ser Leu Glu Ala Ser Leu Ala Glu Thr Glu Gly Arg Tyr
 385 390 395 400
 Cys Val Gln Leu Ser Gln Ile His Ala Gln Ile Ser Ala Leu Glu Glu
 405 410 415
 Gln Leu Gln Gln Ile Arg Ala Glu Thr Glu Cys Gln Asn Thr Glu Tyr
 420 425 430
 Gln Gln Leu Leu Asp Ile Lys Ile Arg Leu Glu Asn Glu Ile Gln Thr
 435 440 445
 Tyr Arg Ser Leu Leu Glu Gly Glu Gly Ser Ser Gly Gly Gly Gly Arg
 450 455 460
 Gly Gly Gly Ser Phe Gly Gly Gly Tyr Gly Gly Ser Ser Gly Gly
 465 470 475 480
 Gly Ser Ser Gly Gly Gly Tyr Gly Gly Gly His Gly Gly Ser Ser Gly
 485 490 495
 Gly Gly Tyr Gly Gly Gly Ser Ser Gly Gly Gly Ser Ser Gly Gly
 500 505 510
 Tyr Gly Gly Gly Ser Ser Ser Gly Gly His Gly Gly Gly Ser Ser Ser
 515 520 525
 Gly Gly His Gly Gly Ser Ser Ser Gly Gly Tyr Gly Gly Gly Ser Ser
 530 535 540
 Gly Gly Gly Gly Gly Gly Tyr Gly Gly Gly Ser Ser Gly Gly Ser
 545 550 555 560
 Ser Ser Gly Gly Gly Tyr Gly Gly Gly Ser Ser Ser Gly Gly His Lys
 565 570 575
 Ser Ser Ser Ser Gly Ser Val Gly Glu Ser Ser Ser Lys Gly Pro Arg
 580 585 590
 Tyr

<210> 51

<211> 494

<212> FRT

<213> Homo sapiens

<400> 51

```

Met Asp Leu Ser Asn Asn Thr Met Ser Leu Ser Val Arg Thr Pro Gly
1      5      10      15
Leu Ser Arg Arg Leu Ser Ser Gln Ser Val Ile Gly Arg Pro Arg Gly
20
Met Ser Ala Ser Ser Val Gly Ser Gly Tyr Gly Gly Ser Ala Phe Gly
35
Phe Gly Ala Ser Cys Gly Gly Gly Phe Ser Ala Ala Ser Met Phe Gly
50
Ser Ser Ser Gly Phe Gly Gly Gly Ser Gly Ser Ser Met Ala Gly Gly
65
Leu Gly Ala Gly Tyr Gly Arg Ala Leu Gly Gly Gly Ser Phe Gly Gly
85
Leu Gly Met Gly Phe Gly Gly Ser Pro Gly Gly Gly Ser Leu Gly Ile
100
Leu Ser Gly Asn Asp Gly Gly Leu Leu Ser Gly Ser Glu Lys Glu Thr
115
Met Gln Asn Leu Asn Asp Arg Leu Ala Ser Tyr Leu Asp Lys Val Arg
130
Ala Leu Glu Glu Ala Asn Thr Glu Leu Glu Asn Lys Ile Arg Glu Trp
145
Tyr Glu Thr Arg Gly Thr Gly Thr Ala Asp Ala Ser Gln Ser Asp Tyr
165
Ser Lys Tyr Tyr Pro Leu Ile Glu Asp Leu Arg Asn Lys Ile Ile Ser
180
Ala Ser Ile Gly Asn Ala Gln Leu Leu Gln Ile Asp Asn Ala Arg
195
Leu Ala Ala Glu Asp Phe Arg Met Lys Tyr Glu Asn Glu Leu Ala Leu
210
Arg Gln Gly Val Glu Ala Asp Ile Asn Gly Leu Arg Arg Val Leu Asp
225
Glu Leu Thr Leu Thr Arg Thr Asp Leu Glu Met Gln Ile Glu Ser Leu
245
Asn Glu Glu Leu Ala Tyr Met Lys Lys Asn His Glu Asp Glu Leu Gln
260
Ser Phe Arg Val Gly Gly Pro Gly Glu Val Ser Val Glu Met Asp Ala
275
Ala Pro Gly Val Asp Leu Thr Arg Leu Leu Asn Asp Met Arg Ala Gln
290
Tyr Glu Thr Ile Ala Glu Gln Asn Arg Lys Asp Ala Glu Ala Trp Phe
305
Ile Glu Lys Ser Gly Glu Leu Arg Lys Glu Ile Ser Thr Asn Thr Glu
325
Gln Leu Gln Ser Ser Lys Ser Glu Val Thr Asp Leu Arg Arg Ala Phe
340
Gln Asn Leu Glu Ile Glu Leu Gln Ser Gln Leu Ala Met Lys Lys Ser
355
Leu Glu Asp Ser Leu Ala Glu Ala Glu Gly Asp Tyr Cys Ala Gln Leu
370
375
380

```

Ser Gln Val Gln Gln Leu Ile Ser Asn Leu Glu Ala Gln Leu Leu Gln
 385 390 395 400
 Val Arg Ala Asp Ala Glu Arg Gln Asn Val Asp His Gln Arg Leu Leu
 405 410 415
 Asn Val Lys Ala Arg Leu Glu Leu Glu Ile Glu Thr Tyr Arg Arg Leu
 420 425 430
 Leu Asp Gly Glu Ala Gln Gly Asp Gly Leu Glu Glu Ser Leu Phe Val
 435 440 445
 Thr Asp Ser Lys Ser Gln Ala Gln Ser Thr Asp Ser Ser Lys Asp Pro
 450 455 460
 Thr Lys Thr Arg Lys Ile Lys Thr Val Val Gln Glu Met Val Asn Gly
 465 470 475 480
 Glu Val Val Ser Ser Gln Val Gln Glu Ile Glu Glu Leu Met
 485 490

<210> 52
 <211> 361
 <212> PRT
 <213> Homo sapiens

<400> 52
 Cys Asn Trp Phe Cys Glu Gly Ser Phe Asn Gly Ser Glu Lys Glu Thr
 1 5 10 15
 Met Gln Phe Leu Asn Asp Arg Leu Ala Ser Tyr Leu Glu Lys Val Arg
 20 25 30
 His Val Glu Arg Asp Asn Ala Glu Leu Glu Asn Leu Ile Arg Glu Arg
 35 40 45
 Ser Gln Gln Gln Glu Pro Leu Leu Cys Pro Ser Tyr Gln Ser Tyr Phe
 50 55 60
 Lys Thr Ile Glu Glu Leu Gln Gln Lys Ile Leu Cys Ser Lys Ser Glu
 65 70 75 80
 Asn Ala Arg Leu Val Val Gln Ile Asp Asn Ala Lys Leu Ala Ala Asp
 85 90 95
 Asp Phe Arg Thr Lys Tyr Gln Thr Glu Gln Ser Leu Arg Gln Leu Val
 100 105 110
 Glu Ser Asp Ile Asn Ser Leu Arg Arg Ile Leu Asp Glu Leu Thr Leu
 115 120 125
 Cys Arg Ser Asp Leu Glu Ala Gln Met Glu Ser Leu Lys Glu Glu Leu
 130 135 140
 Leu Ser Leu Lys Gln Asn His Glu Gln Glu Val Asn Thr Leu Arg Cys
 145 150 155 160
 Gln Leu Gly Asp Arg Leu Asn Val Glu Val Asp Ala Ala Pro Ala Val
 165 170 175
 Asp Leu Asn Gln Val Leu Asn Glu Thr Arg Asn Gln Tyr Glu Ala Leu
 180 185 190
 Val Glu Thr Asn Arg Arg Glu Val Glu Gln Trp Phe Ala Thr Gln Thr
 195 200 205
 Glu Glu Leu Asn Lys Gln Val Val Ser Ser Ser Glu Gln Leu Gln Ser
 210 215 220
 Tyr Gln Ala Glu Ile Ile Glu Leu Arg Arg Thr Val Asn Ala Leu Glu
 225 230 235 240
 Ile Glu Leu Gln Ala Gln His Asn Leu Arg Tyr Ser Leu Glu Asn Thr
 245 250 255

	Leu	Thr	Glu	Ser	Glu	Ala	Arg	Tyr	Ser	Ser	Gln	Leu	Ser	Gln	Val	Gln
			260						265						270	
	Ser	Leu	Ile	Thr	Asn	Val	Glu	Ser	Gln	Leu	Ala	Glu	Ile	Arg	Ser	Asp
5			275					280					285			
	Leu	Glu	Arg	Gln	Asn	Gln	Glu	Tyr	Gln	Val	Leu	Leu	Asp	Val	Arg	Ala
			290					295				300				
	Arg	Leu	Glu	Cys	Glu	Ile	Asn	Thr	Tyr	Arg	Ser	Leu	Leu	Glu	Ser	Glu
	305					310					315				320	
10	Asp	Cys	Lys	Leu	Pro	Ser	Asn	Pro	Cys	Ala	Thr	Thr	Asn	Ala	Cys	Glu
					325				330						335	
	Lys	Pro	Ile	Gly	Ser	Cys	Val	Thr	Asn	Pro	Cys	Gly	Pro	Arg	Ser	Arg
			340					345						350		
	Cys	Gly	Pro	Cys	Asn	Thr	Phe	Gly	Tyr							
			355					360								
15	<210> 53															
	<211> 3282															
20	<212> DNA															
	<213> Homo sapiens															
25	<400> 53															
	atgaaggaga	tggtaggagg	ctgctgcgta	tgttoggagc	agagggggctg	ggcggagAAC										60
	cgcgtgtgtc	actgcgagtg	gcacgcgtgc	agcgtggcgc	tcacccaagc	ttgtctatggc										120
	atcgcttcagg	tgccaaacggg	acccgtggtc	tgccggaaat	gtgaattcca	ggagcgagca										180
	gcacgggtga	gggtgtgagct	gtgcaccacac	aaagacgggg	cattgaagag	gactgataat										240
	ggaggctggg	gcacagtggt	gtgtgcacct	tacatccccc	aggtgcaatt	tgccaaagtg										300
30	ctcaccatgg	agcccatcgt	gctgcagtag	gtgcctcatg	atcgcttcaa	caagacctgt										360
	tacatctcgg	aggagaaggg	cggggagagc	aaggcgccct	cgggagcctg	catgacctgt										420
	aaocggccatg	gatgtcgaca	agctttccac	gtcacctgtg	cccaaatggc	aggcttctgt										480
	tgtgaggag	aagtgtcgga	ggtaggaacac	gtcaagtact	gogggtactg	cgctggagga										540
	ttcagcaaga	tgaagacatc	coggoacacg	agcgggggag	cogggaggag	aagccgggtca										600
35	ggagggtggca	gcattgggggg	aggtaggcagt	ggtttcatct	atcgaggagg	ccagaagaag										660
	gcctccaccat	ccacgcagca	ggagaagcac	cccacccacc	acgagagggg	ccagaagaag										720
	agtcgaagag	acaaagaacg	ccttaagcag	aagcacaaga	agcggcctga	gtgcaccccc										780
	agcatctcca	ccccggccgt	ggccccact	gtgacaagg	tctctctctc	ggcttctctc										840
	tctctccacc	acgagggcag	cacgcaggag	acctctgaga	gcagcagggga	gtcaaggggg										900
40	aaaaagtctt	ccagccatag	cctgagtcac	aaagggaaga	aactgagcag	tgggcaagggt										960
	gtgagcagtt	ttacctctcc	ctctctctct	tctctctctc	ctctctctct	ctctgggggg										1020
	ccctctcaagc	ctgcagctctc	gtccctgcag	agctccctgc	ttgagcagaa	gggtgctctc										1080
	ctggagcagc	cagaggagga	caagtactcc	aagcccccag	ccccccccc	ttcagccccc										1140
	ccctctccct	cagctcccat	gcgccccaag	gtgacacctt	ttgagcagaa	gggtgctctc										1200
	tatggctttg	ggcccatcag	gcgcttctcc	accacccact	ccagctcagg	ccgggcccgg										1260
45	ggccctctcc	ctggggagata	taagtctccc	caogtcaogg	ggctctgggc	ctcggaaggg										1320
	accacacaaac	ggatgccccg	actgagtgcc	agccctgtgc	ctgcgtgaga	gaccctgtag										1380
	acaggccctga	aggagaagaa	gcacaaagcc	agcaagagga	gcgcgccatg	ggcggccagt										1440
	cccaaggggca	cccggaacaa	ggagggcact	gggggccacc	ctgcgcccat	cttgcccgagt										1500
	gcccagctggc	ctggctttac	cgccactgct	gcctcacctc	tctctggagg	ttccctggct										1560
50	agctcggggc	tgggaggtct	gtctctccga	acctttgggc	ctcttgggag	ctgcggccag										1620
	ttgagcctgg	agtcocccct	actaggggca	ggcatctaca	ccagttaata	ggaccaccat										1680
	ttccacagtg	ctgggagtgt	gcgggctgtc	tgacagcacc	ctctctctct	cgacctcctg										1740
	ggggccccag	ggacccctgg	ctgccccccg	ctcagccgct	ccccgttcaa	cagcacccct										1800
	ccctctccct	ctgctctctat	ctccacccact	cagggttttt	ctctggcgtg	ctctaccttt										1860
55	agcctccctt	ctacccacat	ctttggaaac	ccatgggggt	ccgttaatcc	ccctctctcc										1920
	caagctgaga	gcagccacac	agagccagac	ctggaggact	gcagcttccg	gtgtgtgggg										1980

	accctcccctc	aggagagctct	gtcttccatg	tcccccata	gcagcctccc	gcgactcttc	2040
	gaccagacag	ccctctgacc	ctgtgggggc	ggccagttag	accggcgggc	cccagggaag	2100
	actaacatgg	acgagcttct	ggagaagcag	ggcgacgggg	aggccggcgt	caacatcggt	2160
5	gagatgtctga	cgcgctgca	cgcgctgcag	aaggagaaac	agcggtctga	agagcagatc	2220
	ctgagcctga	cgcccaaaaa	ggagggcgctg	cagattctca	acgtgcagct	ctctgtgccc	2280
	ttccctgcc	tgccctgtgc	ccgtccctgcc	gcacaaggcc	ctgtccctgg	gcccctatggc	2340
	ctgcctccccc	aaaggcgagg	cagcgactcc	ttgagcaacca	gcaagagccc	ctcggggaag	2400
	agcagctcag	gcccgggcaa	ctcgctgtcc	actctctctg	aggacccaca	ctcaggctgc	2460
	ccgagccgca	gcagctctgt	ctgtctcttc	cacagcaacc	ccccaacgct	gcccctccctc	2520
10	cagcagccca	ctgcgaactct	gcccctggcc	ctgcctgggg	ccctccgccc	actcccgccc	2580
	cagcccgca	acgggttggt	ccgggaaccc	ggggcagcgg	ggctgggggg	catgcccatt	2640
	ctgtgaggggc	tgtttggggg	gctgggaagg	agtggggggc	tgcccctcaa	tgccctcatt	2700
	gggggggtga	atggggccgc	tgcccccaac	cccgcaagct	tgagccaggg	tgggggggcc	2760
	cccagcgtgc	agctgcgaag	ctgtctcaac	agccttaacg	agcagcagag	acatctcctt	2820
15	cagcagcaag	agcagcagct	ccagcaactc	cagcagctcc	tgccctcccc	gcagctgacc	2880
	caggaaacac	agactgttgt	ctaccagatg	atccagcaag	tcacagcaaa	acggggagctg	2940
	cagcgtctgc	aga ttgctgg	gggtccccc	ctgcctatgg	ccagcctctg	ggcagggaag	3000
	ttccaccctgc	tgctgtctgc	gggtacccct	ggcctgtctg	ccacacgctc	ggcaccacc	3060
	ctgtgtcccg	ctggagccct	agtggctccc	tcgtctggca	acaaacaaag	ttctatggcc	3120
	cgacgagctg	cagctgcagc	agttagcagca	gcagggggac	ctccagttct	ctactgccc	3180
20	acccaacctc	ttcccaagct	ctggggagca	gagggggaag	gggttgggcc	caaaaggagg	3240
	acgcgtgaca	aaggagcttc	agccaaaccg	gaaaaaggct	aa		3282
	<210> 54						
25	<211> 2227						
	<212> DNA						
	<213> Homo sapiens						
30							
	<400> 54						
	gagagccoga	acaggaagag	ggtacagctt	tgtgcaggtc	acatgcccac	tgacgcccct	60
	cagcctctgg	ttcccagagc	ggacttttga	agctgaactg	cttttgttgc	tggaagacct	120
35	atgtttataat	ttacccctggg	tggaaccagg	tgtaacaaaa	gggcaacgct	cccagctccc	180
	cccactcccg	accocggaa	catgcactgg	actacaagg	taaaaaac	agagctgaac	240
	cccaccccca	tgtgtgccc	ctgcgggggg	tacttcatcg	acgccaccac	tatgtggag	300
	tgctgcattt	ccctctgcaa	aaactgcact	gtgcgtaccc	tggaagccaa	caaaactctg	360
	cccatgtgtg	acgtgcaggt	ccataaaaa	cggccgctgc	tgagcatcag	gtctgacaaa	420
40	acacttcaag	acaattgtct	caaaattgtc	ccctggcttt	ttaaaagatg	gatgaacagg	480
	cgccgggagt	ttctatgcag	gtaccccctg	acggaggctc	ccaaaggctc	caatgaggag	540
	cgccggcagg	ttctggagca	ggagaagggg	gctctgagtg	atgatgagat	tgctagccct	600
	ttccatgaat	ttatcgaagg	tgccaggggc	cgggatgaga	agaaggggcc	ccctggagaa	660
	ggggatgggg	acaaaagaga	aacaggggtg	cgtctcctgc	gatgccacgc	agccatgaac	720
45	gtcatgtcat	ttccgaagtt	ttccgcgaac	aaagtggatg	tgcccagcaa	gtacaaaggt	780
	gaggtttctg	acggagggga	gccaactgaag	gaatactaca	ccctcatgga	ctcgcctcac	840
	aatccacctc	cgaggcgga	cgggccctct	ccctcaagtg	acggtgtcca	gcacgctccc	900
	aagcggtcca	ccctagccac	gggtgccaac	ccctccggag	gcaccaaac	cagcgggggc	960
	tcagagtggt	agtaagtgac	cgacaaggct	cccagccctg	ccacccctgc	agccacctcc	1020
	ttccctctgc	ccagcccgcc	cccccatacc	catggtctgc	ccagttccca	tgggccctcc	1080
50	gcccaccacc	ctacccctcc	ccctccctcc	tgacagacga	ctggggccac	cacagctgac	1140
	aaagggggta	gcttgaactg	ccctgcagca	ccatccctca	ccagcagggg	ggcgcaagat	1200
	actgtccaa	ggcgctccgt	gcccocctta	ctgtgaggcc	gcagcaccct	ttccctcttc	1260
	cagccaagcc	ttcccaactc	ttcccaactt	ttctggggcc	tttttccact	ttctcttact	1320
	ttcccaagct	ttcccaactc	gggggtgggg	ttttgggttt	ataataaat	ataataat	1380
55	atgtacatag	gaaaacaaca	ataataatc	ttattttcta	tggaaccaac	agattaaatt	1440
	aaatgccaca	ggaacaacaac	tttatgtgtg	tgtgtatgtg	tggaaaatgg	tggtcatttt	1500

	ttttgggggg	ggtcttgtgt	aaatttgotgt	ttttgggggt	gootggagat	gaactggatg	1560
	ggccactgga	gtctcaataa	agctctgcac	catctotgct	gtttcccaag	gcaagtggtg	1620
	gttggggggg	cccttcagac	ccaaagcttt	aggcatgatt	ccaactggct	gcataagga	1680
5	tcaggttaga	attgttttct	tctctcccoo	ttctctctcc	catcttggct	gctgtctctg	1740
	ctctgacacg	tggccgcccc	cgcggttggt	gaatgtccag	aaattgtctaa	gaacagtgcg	1800
	ttttacaatat	gcagtttata	ccgtgttctg	aggagcaagt	gcaaggttga	ggtggcaccct	1860
	gcataccctc	ctctctcttg	agtggaacac	tggtgcaaat	aatagatagt	tctgtctctt	1920
	tttttttttt	tctctgtgtg	tggtggcctt	gcatacttta	tctgttggaa	aagaagatct	1980
	aggccctcag	aggtctcagc	tcttggagga	gggttaaggg	tttagcattg	tgaagcgtgt	2040
10	caacccacac	aaccttcaacc	tcacccggga	accctcaact	gcaggactgt	ggtgtggagt	2100
	tcacctgggg	ccctgagctg	aagtgggggt	gggttaacct	cacacaagca	cagatccacc	2160
	actttgcacg	aggcaaacag	ggaattccgc	cgatactgac	gggtccacag	agtcgtgcgc	2220
	acactcg						2277
15	<210>	55					
	<211>	4283					
	<212>	DNA					
20	<213>	Homo sapiens					
	<400>	55					
25	ttgcgggaaa	gagccaaacc	ctggcgttgg	ggggcccggg	cggggagccc	ctcccgoggt	60
	ccacagcgac	gctgtccacg	ccctctctcc	cttccggctc	cgccacgggg	ccccgagggcg	120
	ttcggaggccc	aggcgggttt	ctgtcaggcc	cggggaggag	ggggcgggcg	ggcgcccgct	180
	gcctcccgcg	gacggggcgt	accacgggga	cggggaggac	ggggccaggg	gaactgcaggg	240
	gggctgcacc	gcccgggggg	ggggttgagg	gcccggccgc	gggctccccc	ggggcgggggg	300
	cgagtcgggg	gcgtggggcg	gaoggaacca	ccggggcggg	gtgggaggtg	acgggagcggg	360
30	cgcgacccatg	gcgcggtgag	ggagccgggg	tggggatcgg	tcggggggag	gcctgaggcc	420
	gctggcttgt	gcgctgtctc	gcgcgcccc	ctctttcgcc	gcgcgcccgc	ccgcgccggg	480
	catgtcgttc	aaactgcacca	gcacccaggg	ggtggcggtg	gcgcgctcca	gcgcgcagcaa	540
	gaaccaagac	aagaagaagc	attctgtgtg	ccagaaagt	aagctattcc	gggcccagoga	600
	gocgatccct	agcgctctga	tgctgggggt	gaacacacag	atcaatgagc	tgagccaatgt	660
35	toctgttctc	gtcatgtctaa	tgccagatga	cttcaaaagc	taacagcaaga	tcaaggttggg	720
	caatcatctc	ttcaataaag	agaaacctgc	cagccgcttt	aagttttaag	agttattgcoc	780
	catggtgttc	ogaaacacct	gggagaggtt	tggaattgat	gatccaggatt	accagaatttc	840
	agtgaacggc	gaacggcgct	tcacacgtga	cagccagggt	cggtgtggca	ccgctttctc	900
	caccacacct	gaacggcgct	ttgtcatcaa	gactgtgtgc	agccgaggag	tgggcgagat	960
40	gcacaacatc	ttaaagaagt	accaccagtt	tatagtggag	tgctcatggca	acaacgctttt	1020
	gcacacagtt	ctgggcatgt	acgcgctgac	ctgttgatgt	gttgaaacct	acatgtgtgtg	1080
	taccaggaac	ctcaatgaag	atoggctcac	tgctcatcgc	aagtatgaac	tcaaggggttc	1140
	taocgtgtcc	agagaagcga	gcgacaaagg	gaagggccag	gacttgcccaa	cattcaaaaga	1200
	caatgacttc	ctcaatgaag	ggcagaagct	gcatgtggga	gaggagagta	aaaagaaactt	1260
	ctcggagaaa	ctgaagcggg	acgttgagtt	cttgccacag	ctgaagatca	tggaatcacag	1320
45	ccgtcgtgtg	ggcatccacg	acgttggaac	gcagagagcg	gaggagatga	aggttggaagg	1380
	gcgggcagag	gcagaggagt	gtgagaaatg	tggggtgggt	ggcaaacctac	tctgtcccta	1440
	tggaacacct	ccggacacgc	ctggcaacct	ccctcagctt	ctcgtgttgt	gatgtctctg	1500
	ggaatttogac	ccctctgttg	acgtctatgc	catgaaaagc	catgaaagtt	cccccagaaga	1560
	ggagggtgat	ttcatggcca	tcatttgatat	ccctacgcga	taogatacaca	agaagaagaag	1620
50	tgccactgct	gcacaaaagc	tgaaacacgg	ggcaggggcc	gagatctcga	gcttgaaccc	1680
	tgagcagta	tcocaaagct	tcacagagtt	tatgtccacc	atcctgaagc	agttctctct	1740
	taacctcagc	cagagccaga	gagctgggata	tggggtcggg	gatcggggag	taggttgagag	1800
	gggtattttg	ggctagatgg	gagggtggga	gcagagtggg	gttttggagg	gcttttagcaa	1860
	tgagactgca	gcctgtgaca	cogaaagaga	cccttagctga	agagggaggg	gagtggtctgt	1920
	gtgtgcacct	gctccacagg	tgtaacccca	ccctctgctt	acccctgatt	ttttctccccc	1980
55	atttgacacc	cagggttaaaa	aggggttccc	tttttggtac	cttgttaacct	cttgagataac	2040

	cttggggcta	gagatgaact	ogtgggttta	tttggggttt	gtttotgaaa	tttcatgtgt	2100
	caagggttgc	tatttataat	caatatttoat	caagcctaccc	aacctcccca	tottttgtga	2160
	gcctocagtt	cccttcaatt	aaagagatac	ccagtagaac	caagcacaagg	gtccttccag	2220
5	aaaccaagtc	tatggaTGcc	agattggaga	cgctcagacac	ctcgccctgc	tgcatcttgt	2280
	cttgtctgga	tttaactttgt	aatttatgga	gtattgtgca	caacttctcc	caactttccc	2340
	ttggattcaa	gtgaaaaatg	ttgcattatt	gtcccaatcc	ctctcgaata	caacaggtca	2400
	acaocagaga	totoagatca	gaatcaagaga	tctcagaggg	gaataagttc	atcctcatgg	2460
	gatgggtgag	ggcagggaag	oggtgtgggt	ctggttctcag	agaaacctgt	atcctcatgg	2520
10	gatgataccc	caagcccccac	cgctgtcttag	cccttggagt	tcaggaagtc	tctctgtaaa	2580
	gocctgcctc	caataggtca	agaggaaacta	gagtaacctt	ggatttatca	gactccatcc	2640
	gtttaaatgg	ttatttccct	ttgggaaaaac	ttcagaaaact	gatgtatcaa	atagggcctc	2700
	gtgcctctga	tctatttccct	tcttccctctc	gaactccctcc	caggcaactct	tacttctagc	2760
	cgaaacttta	gcctctggga	gatctccaaag	cgactggagt	gcttttttagc	agagacacct	2820
15	ogttaagctc	ogggatgacc	ttgtaggaga	tctgtctccc	tgctcctgga	gagttacagc	2880
	cagcaaggtg	cccccatctt	agagtgttgt	gtccaaaactg	gaggtggctt	ccagtattca	2940
	tgaggtgtgt	atccaggaaa	tcagtttggg	aggcttgatg	tggtgtttga	cctgtgcctca	3000
	gccttggggc	tggttttccct	tggtgcccgg	ctctagactt	ttagccagatc	tgccagccca	3060
	aggctttttt	ggaaggagtg	gcttccctgca	gggtgtccac	ctgcctctgg	agcctggccac	3120
20	ccaggccctc	agaactgagc	caacaggctgc	tctgtgcaatc	agagaaaacg	ctctgtttgt	3180
	ctgcaatggg	ggaggtacat	tctgtgcaatc	tctcccccct	tcaaccaggga	atctgtggatt	3240
	tggtatgaga	tatgtctcaga	cttctagata	accccaaaaga	tggtgaagatc	gcttgtgaaa	3300
	ccattttgaa	tgaattagatt	gggttccctgt	ggctccctcc	aaacccctggc	aaagcccaagt	3360
	tcogaagcag	gaaccagcac	tgctctctgt	ccctgactcc	agcataatgg	tcagggaaga	3420
	atggagacgg	cattcttggga	cttcaactggg	gcctgtggat	tggaaggga	acactcttga	3480
25	agaggcagat	gggggtcaaa	ccactgacct	ggcccccagg	agggggcaata	ggtaggtctg	3540
	aaacaactgc	ccagacacac	tacatgaact	agggaaactg	aaacccaactg	gctcaatggc	3600
	aaacaacaa	tgacttggga	aagggaattat	gtaggaaata	tggtttggaat	tgatttcccc	3660
	acgtcataat	gaagaaatga	agtttggatc	ctgcctccgt	caggccagcc	actctctgaag	3720
	cttggaaagc	tgctctccag	ctcccaaac	tgcccaagcc	cagcttccga	agcaggaacc	3780
30	agcaactgtc	ctgtgctctc	ctccacagcat	ataggtccag	aaagaaatga	cagccgcatc	3840
	ctggacttca	ctgggtccgc	tggtattggat	gggaaacact	ctgggaagagg	cagatggggg	3900
	tcaaaacact	gccttggccc	cagggaagggg	ccataggtag	gtctgaaaca	ctgcgcgaag	3960
	accaactcat	gacttaggga	acttgaaacc	aaactggctca	tggaagaaac	aaatttgaat	4020
	tggaagagg	attatgtagg	aaatgtgtt	ggacttgatt	tcccccacgt	ataatgaaga	4080
	atggaagttt	ggatctgtct	ctogtcaagg	gcagcatctc	tgaagcttgg	aaagctgtct	4140
35	tccagcagcc	tcogtggcct	cgggttccca	ooggtctctc	tgaatttggg	ctgctgatca	4200
	tggtgcata	atgtgtatgg	aaagtgtaac	acattctctc	tggttaagaa	cgactaccag	4260
	gtatcatact	tgcttaacat	tga				4283
	<210>	56					
40	<211>	6140					
	<212>	DNA					
45	<213>	Homo sapiens					
	<400>	56					
	ggggccgacg	ccctgagccag	ggcccccctcc	ctogtcaagg	cogggggcagc	aagcaggccg	60
50	ggggcaggtc	ogggcaccca	ccatgcgagg	ccagctctgc	ctcctgggtgc	tggtgtctag	120
	ggaggctgac	ogggcgctga	gccccagacc	cggagcaggt	caagatgagg	gccccaggtc	180
	tggtatgggt	gccaaaggga	cogtgcgggg	ctggaaacgg	agagcccgag	agagccctgg	240
	ccatgtgtca	gagccggaca	ggacccagct	gagccaggac	ctgggtgggg	gccccctggc	300
	catggacaag	ctgcacagta	acagggaacc	gggtgtggag	gacacaccca	gctattatgt	360
55	gtcccgctct	tatggcccca	gogagcccca	cagccgggaa	ctgtgggtag	atgtggccga	420
	ggccaaacgg	agccaaagtga	agatccacac	aatactctcc	aaacccaccc	ggcaggcttc	480

	gagagtgggt	ttgttcotttg	atttccottt	ctacgggcat	cotctggcgc	agatcaccat	540
	agcaactggg	ggcttcactt	tcattggggg	cgtgatccat	oggaagtctc	caagctactc	600
	gtatgttgag	ccocctgatg	ccaacttcaa	ccctggctac	tcogacaaat	ccagcttgtt	660
5	ttactttggc	aatggggacg	tcctttgtgt	tcagtgggac	caogtttatc	tcocaggctg	720
	ggagagcaag	ggcagtttca	ccctccaggc	agctctgcac	catgacggcc	gcatttgttt	780
	tgccatacaa	gagatcccta	tgctgttccc	ggaaatccgc	tcctcccaag	atccctgtca	840
	aaocggccta	tcoggatgctt	tcattgatctt	caatccatcc	coggatgtgc	cagaatctcg	900
	cagagaggag	atctttgaa	acaccccgat	agagctggac	cccagcaagg	tcaccagcat	960
	ctggcgctgt	gagtttcaacc	cattgcgcag	ctgcctgcag	cataggagct	gtgacgcctg	1020
10	catgtctcga	gcagctgaact	tcacactgcag	ctgtgtccat	gtcctccaga	gatgtctccg	1080
	tggtcttgac	cgctatcgcc	aggagtgat	ggactatggc	tgctgcacag	aggcagaggg	1140
	caggatgtgc	gagggacttcc	aggataggga	cgcgaactca	tcctccctgt	acactctgtt	1200
	cagccoccat	gatggagacc	tcacccatcc	ctcctctccc	ctcttccatg	acagcctcac	1260
	cacagaaagt	gacacccaag	tgaaatccct	tcgaggagga	gaocgcoctc	agaaacacct	1320
15	gtcccccagg	acaaaggcca	ctcctgttga	cctgggcaac	atcgtgggca	tcgtgtctgc	1380
	agtcctctcc	gtggcgccca	tcactcttgc	tggaatttac	atcaatttac	tggaatttac	1440
	caatgtctgc	ctcttctcca	tccttctcca	gggtggagcc	tgggccagca	tgaaagtttg	1500
	cagccacccc	gacacttcca	cccttgagca	cccttgagcc	tgggccagca	tgaaagtttg	1560
	cttcctggag	gctgagcagt	gctgagaaac	acagttctcc	cctttgaaga	ctttgagcgc	1620
	acagaaaaag	cagtttaaaag	aaagaagaga	agtgactctt	cctggcctcc	ccacagctgc	1680
20	cctggcgctga	gatgagatgg	tggttttatgg	ctccagagct	tcgtttctgt	tcgttcagcc	1740
	accccgaaat	ttgaaagggg	ggccaaaaaa	caaccacatg	gattttttat	agggacaaac	1800
	acctaattct	atactgtttt	gatgaaaggg	ttctctctgt	tgctttgtta	catgtgaaac	1860
	cgagaaagac	taacataact	aactccattt	ttgttttaag	ggcctttaoc	tattctgcga	1920
	octagtcctag	gataaacttt	gagcacagga	ataaaacgca	aaaacaggaa	tcactcoggt	1980
25	tgcaaaaact	actctgggat	taaaagggaa	gcattgaaac	agctaaactg	ttttgttaaa	2040
	gattttatag	aatgaggggg	tttggctatt	gtccacatgac	agactgttag	ccagggacaa	2100
	agaagttctg	caaacccccc	ctgggacccc	ctgggtgtcc	agatgtctgc	ggttgttaag	2160
	ccctctcttt	cccccgaact	aaacataaaa	gacaaaggca	agccgcatac	attttaagac	2220
	ggtttcttgc	gacacttatg	ccacatcttc	ttggtttgtc	ggctctccga	aataaagtcc	2280
30	ctttctcttg	tcctaacctc	tgctctccaa	ctgtattgct	atgacgcagc	agacagagtt	2340
	aatttggact	cagttacagg	ctgtccaatg	tcctgtctgt	agcagttcca	gagcctcccc	2400
	gaccccatcc	ctggagagtag	ccagatagcc	agatgocctg	ctcctggcca	cctttaaagc	2460
	ccctgcatac	gacacaggtt	aactaaagtc	aagattgggg	ctcctgcaat	ccaggttccc	2520
	tgactccaca	agctgtgtcc	tgccacaggt	cagtggtcca	ccacgcaact	ccagcgaact	2580
35	tgggaggctg	aggaaggcgg	atccactgaa	gtccagaagt	tgagacacag	ctggccaaac	2640
	taattaaaat	gtctctacta	aaaatacaaa	aaattagctg	ggtgtggtga	cgcttgctgt	2700
	tatccagctc	actcaaggag	ctgagacacg	agaaatccat	gaaocctggg	ggcagaggtt	2760
	gcagtgagct	cagatagttg	cactgcactc	cagcctgggt	gacagagoga	gactccgctt	2820
	caaaaaaaaa	aaaagaaga	agaaotccat	gcctatagag	ttggcatttt	agcccccagc	2880
40	ctctgtagct	tgaaagccca	aagaaggtat	tcctctccac	tggttaaacac	agttatagttg	2940
	ctctcagccc	ctggggcag	ttatccatgg	agggaaagca	gagaaagaga	gagaaagaga	3000
	ctccaaaggg	aaactgcatt	tttaggcttt	gctctcttcc	ctctgcoctt	ctactcagaa	3060
	ccaatcaact	ctgcactaaa	acatgtttac	cctgcactca	agggccttcc	cccaacotgc	3120
	agccacgctc	tcctctgggt	agcttgcctt	cgcagagocac	atttcaactg	tggggctccc	3180
	tattctgatg	gcctgtctgg	tgccgggttt	actcactgcc	ctgtttctgt	ctcagttgct	3240
45	gtacataact	ccaaaggcag	gacttgcctg	ataaatattt	ttcctctctc	gaacttggtt	3300
	ttatagtcac	taaaagccaa	tcgggtggct	agagggctcc	ttgagacata	ctcagcaggg	3360
	aaotcgaggt	ggattctgtt	gagagggcaa	gcacgtgagt	ggttgggaca	cagggcagct	3420
	gcattggagg	gacttttttt	gagacaggg	ctcactgtgt	cccccagggc	taaggtgcoc	3480
	aaagacacca	ggttgggag	gcaactggc	actaactgct	ttccctggag	ctcgtatgtg	3540
	ccbtgggggt	ggggagcgtt	aggggtctac	ggctgocgtg	gatgggtgtg	ccagcttgtt	3600
50	gaagtaccta	actcctgtcc	ttgtctgact	gacacagact	gcagggcgcg	gcccacagac	3660
	ccatgtctcc	atctggtcat	actccatagc	tacaaagtta	acotctctca	aactctggag	3720
	aactggatct	gtcccaataa	cgcttatttg	gcacagcctg	atggctctct	ctcgttactcc	3780
	cagcaacttg	ggagcctgag	gtggggaggt	tgcttgagcc	caggggtttg	agacagcgtt	3840
	gggcaacaac	acacaaaaat	ccaggtgtgg	tggtgtgcac	ctgtagtcoc	agctactagg	3900
	gaggtcgagc	caggagagac	acttgagccc	gggaggttga	gggtgcagtg	gggggtcata	3960
55	atcctgccc	tgtaactcag	cctgggtgac	agagtgagac	ctcgtctccg	aaaaaataaa	4020

	aaaaaaaaga	acggaaaaag	aatgtcttac	attgtcaggg	atcctgtaga	caatcattaa	4080
	ctctatgaga	tgcttggttc	tatttttttg	ggagactctg	tcocagtggt	ttgggttaag	4140
	aatccatag	gocctctctg	gtgacacatc	tctagtaact	ttgttcataa	acaaacaggc	4200
5	catctgcgcg	caaatacatc	cactccocat	gcacgtgaca	tcctatgggt	agccagggtc	4260
	tgctttgaat	gaggcgagg	catctggaac	ttctcttgcc	tgacagggtc	agcagcagag	4320
	gcttcacgcg	atocacaccc	cttctccac	tcttgacatt	ctttcccttc	agggatccaa	4380
	aatggttgcc	cgagatccca	gtgggaaaac	gtgtgctaga	gttggggagt	gagatgagtg	4440
	gtctgtccca	tggaatccag	ccacagcagg	aactgcceca	ctggccattt	gagacacaca	4500
	cagggtgtga	atgctctgct	gggtggctgt	gcttccctca	ttcagagagc	tctgtttacg	4560
10	cccactgtgt	ctcttagaag	cttgaaggga	acccaactct	ttgtgtgact	gtcctttttc	4620
	ttctccaaat	tcagacccctc	cttccacccg	caccccctca	ctccacccctc	agctcttctc	4680
	tgccgtgttt	atcaagcaga	gctgaggccc	caogtttcca	actctgattg	tcacttgcat	4740
	cttccaaag	gataaaccca	ggagcaactg	gaaaaccatc	agccaagcgt	tcggatgagt	4800
	ctggttaatg	gtccacccccc	gaocagattc	ccttacactt	aactcacttc	ttcttttgge	4860
15	aatgacccct	atgacatgta	taaatgggta	tgactaagaa	gaggctgtga	tctaacattt	4920
	atttgtctgc	attttttaact	ctggggagaa	gcagccccaa	ctcatcactg	ggaaagaact	4980
	ccocctgcga	accagcttaaa	tttgataa	taaaacccctc	gcccctaaaa	cttctcaacg	5040
	agctggggag	ttgtgtggcaa	ctttccaa	caaggctctg	ttggggagct	gaggcaggag	5100
	atggccagggt	gcagtggtct	acogcgttag	tcocaggtag	gatca tccca	ctgcatctgt	5160
20	gtatcgttga	gctcaggagt	tcaaggctgc	agagagctat	ctccctgtac	atgttatctt	5220
	ttaaaaataa	atttttaaaa	tttgtgtgtt	ttatcagggg	tttctgtgag	atgaagatatg	5280
	tgtagtgttg	ttgtgtgtgt	tgtagatcag	ctgtttta	gtttgtgagc	atgaagatatg	5340
	cacacacagg	tattttgttg	ctaaagagat	tggaacaagt	tgtagctgtg	ctcagggtctc	5400
	agctgtgttt	gttaaatgta	gagataaaca	atgacaagag	ctgccaagca	acccaactat	5460
	tcacaaagca	aagtgtttcac	caactaaagt	acaacttc	ctggtgtcag	gcaagggttaa	5520
25	ggctctctct	cccttagttc	ctggaacaga	ctccagagat	ggcagtgaag	actgatcaga	5580
	ggctgcacct	agactccctg	cccaagcaaa	octacacacg	aagatgcagt	gtccacagta	5640
	tggtgogccg	aatctctgtc	tcocaaaaac	taoctgaact	taattgttag	attcaagaatg	5700
	ctggggactg	agggcaacca	gocctctaaa	acaacatgta	tcca tgtgtt	tagtgttaaac	5760
	tctctgcact	gattctcagt	gttaataata	aaagggaagc	ttcttttaca	actcctgctg	5820
30	gtgtcacaag	aagtgtcaac	ggatttggag	tgccattccg	aagatcaaca	caactcactt	5880
	ggttctgatg	gctgtcgaa	tcogacttct	tcgtgtgagc	atgactgtgt	gaaacagctc	5940
	cagctatctg	ctcactccag	gtgctttctc	caactcctg	caacacctcc	aagagaaaac	6000
	gocataaaag	aaaccccgac	tgctttacta	tattgtgtct	taaatccctg	gaagttaaac	6060
	ccatgcattt	ttactctact	tcgtgaggca	tacaataaat	ctgagaaagt	cta tgcctg	6120
	aaaaaaaaaa	aaaaaaaaaa					6180
35	<210> 57						
	<211> 2098						
40	<212> DNA						
	<213> Homo sapiens						
45	<400> 57						
	gcaggagcac	gtggagaggg	cggtgtgaca	cagoggcagc	tcacggccgg	ccggcgagcg	60
	acatgtgaaga	tatacaaaac	aatgoggaa	tgaaaagcac	tcaggagcag	ctgtgtcccg	120
	cagaaagtgc	agcgggtttg	aatgactaca	gtttaaccaa	atctcatgaa	atggaaaagt	180
	tggaacagtgc	agaagggcca	gccaatgaag	atgaagacat	aggagatgat	tcaattgaag	240
50	tgaaagtga	atacagtga	agagatgaga	atgttttaaa	gtcagaaccc	atgggaaatg	300
	cagaagagcc	tgaattccct	tacagctatt	caagagaata	taattgaata	gaaaacatta	360
	agtgtagagag	acattgtgtc	tcattcgata	gtacagggc	aaccagtgga	gtatgtgaact	420
	cgatgtgttg	tggttatctc	tgcatcagct	tcaatgtctt	aatgtttcat	aagcgaagcc	480
	atactgtgtg	acgcccaatt	cagtgttaac	atgtgtgggc	atcttttact	cagaaaaggta	540
	acctctcccg	ccacattaaa	ctgcacacag	gggaaaaacc	tttttaagtgt	caactctgca	600
55	actatgcactg	ccaagaaga	gatgogctca	cggggcactc	taggacacat	ctgtgtggaga	660

	aaccotacaa	atgtgagttt	tgtggaagga	gttacaagca	gagaagttoc	cttgaggagc	720
	acaaggagcg	ctgcogtaca	tttcttcaga	gcactgaccc	aggggacact	gcaagtggcg	780
	aggcaawgca	catcaaaagca	gagatgggaa	gtgaaagagc	totogtaactg	gacagattag	840
5	caagcaattgt	ggcaaaacga	aaaagctcaa	tgcttcagaa	attcattggg	gagaagggcc	900
	actgotttga	tgtcaactat	aattoaaagt	acatgtatga	gaaagagagt	gagctcactac	960
	agacocgcag	gatggaccaa	gcacatcaata	acgcatcag	ctatcttggc	gcogaagccc	1020
	tgtgccccct	gtgtccagaca	ocgootgtct	ccacctogga	gatgtgttca	gttatccagca	1080
	cgatgtatcc	catgacccct	acccgggctg	agatgtcaaa	cggtgcccc	caagagctgg	1140
10	aaaggaagaag	catctcctct	ccagagaaga	gcgtgccttc	tgagagagcg	ctctctccca	1200
	acaatagtgtg	ccacgaactcc	acgggacactg	acagacaacca	tgaagaacgc	cagaatccaca	1260
	tctatccagca	aaatccactg	gtcctgtctct	gggcccgcga	tgggatgcga	cttcttgaagg	1320
	aggttccocg	ctcttaacgaa	ctcctcaagc	cccccgcact	ctgcccaga	gactctgtca	1380
	aagtgatoga	caaggaaaggg	gaggtgatgg	atgtgtatcg	gtgtgacac	tgccgcgtcc	1440
	tcttccctgga	ctatgtgtatg	ttoacgatto	acaaggcctg	ccacggcttc	cgtagacott	1500
15	tcgagtgtaa	catgtgtgga	gatogaagcc	atgatcggt	tgaattctog	tctccactag	1560
	ccagagagga	acacagaagc	ctgctgaagt	gaatctctgg	ctccagggat	tgctctctatg	1620
	tattccagcat	ogtctctaaa	aacaggtgac	ctcgcctaac	agattgtctct	caaaacatac	1680
	tcagtcccaa	actctctttc	atacactttt	tagctgtgtt	cacaggggta	gcccagagaa	1740
	cactgtctct	cttcagaagt	tattgcagg	tattgacttt	gtgaaacott	gtgaaacott	1800
20	tgcttttccca	tcagggaact	gaattttatg	gaatttaaaa	gccccaaaag	tatttggtca	1860
	ttatctctca	cagcagtgga	atgagtggtg	ccggagagtg	gctatatgaa	acattctcttc	1920
	tgagatatac	caaccacacg	tggaaaagcc	tttcagctat	acatgcaaat	ccacaagagc	1980
	gaagagotga	ccagctgacc	ttgtctggaa	ccctgacctc	tctgcccctc	acaggtctgaa	2040
	gggttaagat	ctaattctcc	taattctaaa	gacagctcaa	gagtaagtaa	aagaacag	2098
25	<210> 58						
	<211> 2947						
	<212> DNA						
30	<213> Homo sapiens						
	<400> 58						
35	atgcacaatc	ctctccccc	gcccacccca	ccgtgtctcc	ctccacccct	ccactttctat	60
	caggccaaca	cagagcagcc	caagctgagt	agagatgagc	agcggggtcg	agggccctcc	120
	ttacaggagca	tttgcaaaag	gaccaaagctg	aagaagggtga	ccaacattaa	tgatcggagt	180
	gctcccatcc	togagaagcc	gaagggaagc	agtgtgtgtg	atggctctcg	aggagctgcc	240
	ctgcagccca	agggaggtct	cttccaaagga	ggagtgtctga	agcttcgacc	tggtggagagc	300
40	aaaggaaggt	caggaacact	agctggtaag	ccagccctgc	aaatcccccag	ttctcagagct	360
	cgctgcgccaa	ggcctccagt	atctgcgcgc	agcgggctgc	ctcaggtatga	tacagacagc	420
	agcggccgct	actgcacaga	actgcocccg	atgcagagac	ctcttttacc	ggactctctct	480
	gggccttaata	ccacccagcag	taaggccatg	aagcagcagt	ctctctcccc	tcccccacaa	540
	ccccccgggc	ggcgtgcaca	cgccacccccc	acacctctgc	ctatgcacag	cagcaaaagcc	600
45	cccgcttaca	acagagagaa	accccttgcca	ccgacgcctg	gacaaaaggt	tcacccctgg	660
	cgagagggag	ctcctgtctcc	acccccagtc	aaaccaactc	cttccctgtg	gaatatccga	720
	acaggacaaa	gtggccagtc	ctctgtctct	actctccgcg	cttacccgca	ccctcctggg	780
	gtccccaatg	gacccctcag	ccccactaat	gagtcagccc	ctgagctgac	gacagagagc	840
	aattctttgc	ataggagaac	acccagggct	gtcagagggc	tgagcacctcc	tcacccaccc	900
	tcggcctccc	catctttact	gagttaatag	ccacccctcc	cagcccgaga	ccctccagct	960
50	cggggagcag	ctctccaccc	ccccacaccc	gtgatccgag	atgggtgccc	ggatgtctcc	1020
	ctcccccacc	caacataccg	aatgcatggg	tcagaaacccc	cgagcccgag	aaagccccc	1080
	ccctccaccc	caaggagccc	agctggggcc	cccccctctc	ctccacccgc	ctcagggaat	1140
	ggccacagag	attctatacc	actgtctcgg	ctttctctgg	atgattttga	gtccaaagtat	1200
	ctctctccat	cagtagtaaga	cttctctgct	ccagagaagt	ataaacactt	tcaggagata	1260
55	tatccacaga	aaacaaaccc	agctgcocct	ggagccccc	ctctgcccac	catctccagc	1320
	tgaagcctgg	ctgtgtcccg	ttctccagga	aaaggaatga	cttctctctc	ttccagatg	1380

	gtoccttoca	ttoccttcaa	acctgcattga	gagctccttaa	catgtttcttc	caatgcacac	1440
	aagcctcaga	ctccaaatgt	cctccagct	caactccatc	tatgcacctc	atctctggat	1500
	ttggtgata	gactctatat	tgacagtagg	atctcaaac	ctgcacacat	ccttctctca	1560
5	gcaagccttg	ctagccacat	gaggaacaag	ttctcgtgtc	ttctcgtctc	ctcttgggga	1620
	aaggtgcctt	gttggtgata	attaaactac	tgtagggcca	gggtggagaa	tggtactcct	1680
	tcctctcct	gtccactgtg	ggggagcctt	ggcaggata	ttatatctca	tcatttagga	1740
	ggctgcatg	accagagcctt	atgggtggga	ggggagcatt	tttagtgaag	caagaaagga	1800
	gtttgccag	aagtgtactg	ttttaaaggt	catatttggga	gaaagggcaa	ggaattgggt	1860
	ctgtcttatt	tttgggggta	ttttgttttt	gttctcaact	gctgcccccc	caccccacca	1920
10	cccaagggat	aaattgggata	taaacactaa	atactaatca	gttgaaactta	acattttaata	1980
	aaaagaaagg	gtgaaataaa	ctgaagacaa	ttttagaact	agtcaagtct	ctgcagcaaa	2040
	gggaacagg	gccatttgaa	ccctctggga	ccctcaaccc	cactgcttca	gggtgctagg	2100
	ctgagggatg	tttttctctc	cccttaacgc	caatgccttt	gaaagaaaag	tcactttttg	2160
	tgaggggcat	cattcattcc	tgattcccaa	accccaaaaa	cctctggtgg	gagataggaa	2220
15	gatagggcgt	gggcctgggg	cttaaacctca	atcttgtgtc	tgccctcagtc	ttttctgact	2280
	ggccttgagg	ttgtcagttg	ctcttttgtt	ctctcagccc	ctggaaggtg	ctccaggata	2340
	acaaagagg	gcaggttgaa	gcccctcagt	gaaggagctg	gcttctgggg	gctgcagaag	2400
	acttttaagt	cctgcctgta	ctgaagttca	cagccacact	ctctgagcag	actcttctctg	2460
	ttcctttctc	tacccacctt	gcttccacag	gactgcacgg	tttaacacag	cagagtaacag	2520
20	aaggggtgaag	aagtgaacag	aggcttatga	agatatccag	atactctctt	atgccaggaa	2580
	gcacaaagac	tttgttgaga	tttgctcag	ttcagtagat	gttctcttgg	agccagccat	2640
	agggtgttct	tttgtctctc	gggtctcaaa	gagcacagag	aaaaaggagg	tcctccagctc	2700
	aggtaggaag	ctgatctgat	gaggactctt	tttttctoga	cagcaggatg	gggtctcttg	2760
	gctocacaca	ccagatgctt	tggtttctta	caactgttgc	tatgtgtaga	gggtgctcag	2820
	agctgtgcat	gagagcaagg	agacacatgg	tactctttga	aatggatggg	gaaatttagc	2880
25	ttaaaaattt	aatcacgaga	ttgcgcact	gcactccagc	ctggggcgaca	gagccagact	2940
	cogtctc						2947
	<210>	59					
	<211>	784					
30	<212>	DNA					
	<213>	Homo sapiens					
35							
	<400>	59					
	gagcgtgtgc	gcagtggaag	ctagacccgg	tttactggaa	ttgctctgac	gatcgagggg	60
	tcctagtaca	cgcgaactcat	gtctattatg	tcctataacg	gaggggcoct	catggccatg	120
40	aaggggaaaga	actgtgtgctc	catcgctgca	gcagcggcgt	tcgggatacca	ggccagagatg	180
	gtgacccacg	acttcacgaa	gatctttccc	atgggtgacc	ggctgtacat	cggtctggcc	240
	gggctgcgca	ctgaagttcca	gacagttgcc	cagcgcctca	agttcggctc	gaactgtgat	300
	tgattgaagg	aaggtcggca	gatcaaacct	tataccctca	tgagcatggt	ggccaaacctc	360
	ttgtatgaga	aacgggtttgg	cccttactac	actgagccag	tcattgcccg	gttggaacctg	420
	aagaccttta	agcctctcat	ttgctctcta	gaactcactg	gctgccccat	ggtgactgat	480
45	gactttgttg	tcagtggcac	ctgcgcgcga	caaatgtacg	gaattgttga	gtccctcttg	540
	gaccccaaca	tggaatccgga	tcacgtgttt	gaaaccaatc	ccccagccat	gctgaatgct	600
	gtggacccgg	atgcagttgc	aggcatggga	gtcattgtct	caatcatoga	gaaggacaaa	660
	atcaccccca	ggacactgaa	ggccccgaatg	gactaacctc	gttccacag	cccccttttt	720
	ttcttttttt	tgaataaaaa	tagcctgtct	ttcaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	780
50	aaaa						784
55							

	accoccttgctc	cotcttttgag	ggagggggagc	tatgcttagga	ctccaaacctc	agggactcggg	2940
	gtggcctcgog	ctagcttctt	ttgatactga	aaacttttaa	ggtggggagg	tggcaggga	3000
	tgtgcttaat	aaatcaattc	caagcctcac	ctg			3033
5	<210>	61					
	<211>	1174					
10	<212>	DNA					
	<213>	Homo sapiens					
15	<400>	61					
	aagctctctac	cccgccggcg	agccaggagg	aaaggatggc	cgccctggcg	gcccgggttg	60
	tactcttagc	tggggcagcg	gogctggcga	goggtcccca	ggcgagccgt	gagccgggtg	120
	accggaactg	cgtactgcag	tgcgaagagc	agaactgctc	tggggggcg	ctgaatcaact	180
	tcgcgtcccg	ccagcccaatc	tacatgagtc	tagcagcgctg	gacctgtcgg	gacgaactgta	240
20	agtatgagtg	tatgtgggtc	acogtggggc	tctaacctcca	gggaaggctca	aaagtgcctc	300
	agttccatgg	caagtgggcc	ttctcccggt	tctgttcttt	tcaagagccg	gcactggccg	360
	tggcctcggt	tctcaatggc	ctggccagcc	tggtagctct	ctgcgcctac	cgacactctg	420
	tgcacgcctc	ctcccccattg	taccacacct	gtgtggcctt	cgccctgggtg	tcctccaatg	480
	catgtgtctg	gtccacagtc	ttccacacca	gggacactga	cctcacagag	aaaatggact	540
	actctgtgtg	ctccactgtc	atctctacact	caatctacact	gtgctgcgtc	aggacogtgg	600
25	ggctgcagca	cccagctgtg	gtcagtgctc	tcggggctct	cctgctgctc	atgctgacog	660
	tgcacgtctc	ctaacctgagc	ctcactccgct	togactatgg	ctacacacct	gtggccaacg	720
	tggcttatgg	cctggtcaac	gtggtgtgtg	ggctggcctg	gtgcctgtgg	aacacagcgc	780
	ggctgcctca	ogtgcgcaag	tgcgtgtgtg	tgtctctgct	gctgcagggg	ctgtccctgc	840
	tgcagctgct	tgactctcca	cogctctctc	gggtctctga	tgcctctgct	atctggcaca	900
30	tcagcaccat	ccctgtccac	gtcctctttt	tcagctttct	ggaagatgac	agcctgtacc	960
	tgtgaaagga	atcagaggac	aagtccaagc	tgtttgaagc	agactggtat	tttctctctg	1020
	ccctgacccc	ttgtccctct	ttgaggaggg	ggagctatgc	taggaactca	acctcaggga	1080
	ctcgggtggc	ctgcgctagc	ttcttttgat	actgaaaact	tttaaggtgg	gaggggtggca	1140
	agggatgtgc	ttataaatac	aattccaagc	ctca			1174
35	<210>	62					
	<211>	3167					
	<212>	DNA					
40	<213>	Homo sapiens					
45	<400>	62					
	aagctctctac	cccgccggcg	agccaggagg	aaaggatggc	cgccctggcg	gcccgggttg	60
	tactcttagc	tggggcagcg	gogctggcga	goggtcccca	ggcgagccgt	gagccgggtg	120
	accggaactg	cgtactgcag	tgcgaagagc	agaactgctc	tggggggcg	ctgaatcaact	180
	tcgcgtcccg	ccagcccaatc	tacatgagtc	tagcagcgctg	gacctgtcgg	gacgaactgta	240
50	agtatgagtg	tatgtgggtc	acogtggggc	tctaacctcca	gggaaggctca	aaagtgcctc	300
	agttccatgg	caagtggggc	ttctcccggt	tctgttcttt	tcaagagccg	gcactggccg	360
	tggcctcggt	tctcaatggc	ctggccagcc	tggtagctct	ctgcgcctac	cgacactctg	420
	tgcacgcctc	ctcccccattg	taccacacct	gtgtggcctt	cgccctgggtg	agaaaactga	480
	gtccacagcaa	cttgcccaag	gacacacagg	aaatgcagag	ccagggaactg		540
55	aaocctggca	gtctggctgt	agggcttgca	ttcttaatga	taccactaac	tcocaaactct	600
	gaggaagggg	tgctccctcaa	tgcactgtgtc	tgggtccacag	tcttcacac	cagggaactct	660

	gaootcacag	agaaaatgga	ctacttctgt	gootccaoty	toactactaca	otcaatotac	720
	ctgtgtctg	tcaggtgag	ctgootgggt	ggctgcaagg	gcaaaatoga	acootggggg	780
	cagaaaaggg	tcacccagcc	ttccoottgg	ggoottcttc	actagtctcc	caacacccac	840
5	gcccccaac	cccccacaca	tcagctgtcc	tgggtgagga	ctctggggta	ggactggggg	900
	ccotggctcc	tgacaaggag	ctgtagcaot	tgttgocag	ctgtggocgt	tttgggtggg	960
	agaggtctag	tgacttcag	ggccatgoc	caatgttggg	gggaggagat	gtccaggga	1020
	atgtgtctgt	ggggatgggc	oaootgoot	ctgagcaacc	ctggagcgtg	gggcaggacc	1080
	tgtgggtctg	agcacccag	tgtgtcagt	goottoogg	ctctctgtgt	gtccatgtgt	1140
	acoottgca	ttctctcaot	gagootcaot	cgottogact	atgtgtacaa	ccotgtgggc	1200
10	aacgttgcta	ctgtgtcgt	oaootgggt	ctgtgtcgt	ctgtggaacc	gtggaaccag	1260
	cggtgtgtgt	ctcaogtgc	caagtgtgt	gtgtgtgtgt	tgtgtgtgt	gggggtgtgt	1320
	ctgtctogag	tgtctgaatt	ccacogctgt	ttctgggtcc	tgtgtccca	tgcacatgtg	1380
	cacatcagca	ccatccotgt	ccacgtctct	ttttcagct	ttctgggaag	tgacagcctg	1440
	taactgtgga	aggaatcaga	ggacaggttc	aaactggact	gaagacottg	gagcagctgt	1500
15	gccccagtg	ggatccotgt	ccogootctg	tggootccot	ttccocctca	acoottgaga	1560
	tgtattttct	ttttccactt	cttgaaottg	gaactgaagg	atgtggggcc	agaaatcagt	1620
	ggocagccca	ccocctgtgt	gootccacca	goottggaat	ctgttctagg	gaaggcctcc	1680
	cagcatctgt	gactogagag	tgggcagccc	ctctacotcc	tggagctgaa	ctgggggtga	1740
	actgagtgtg	ttcttagctc	taocggggag	acagctgoot	gtttctctcc	ccacagcctc	1800
20	ctcccccact	ccocagctgc	ctggctgggt	ccggaagccc	tctgtctacc	tgggagacca	1860
	gggacacag	gcttaggga	tacaggggg	ccocctctgt	taocaccccc	ccacotctct	1920
	caggacacaa	ctaggtgtgt	ctggatgctt	gttctttggc	cagccaaagt	tcacggcgat	1980
	ttctcccatg	ggatctttg	ggacccagct	gtctgggatt	ggagggagtt	tcacccctgac	2040
	cggtgcccct	ggcaggttcc	caggagcctt	caoccatctc	cccttccagg	ccagggctcc	2100
	agccaagccc	ggccaaggat	ccgtgtgtgt	tgtctgggtg	agagcctgct	acogtgtgtc	2160
25	gggagtggtg	ggcaggtctg	gtgcataggt	gacagggccg	tgagcatggg	ccctgggtgtg	2220
	tgttaggtctg	ggcaggtctg	ggcaggtctg	agacgggtgt	tgtoggggaa	gaggtgtgtg	2280
	ttcacaagtgt	gtgtgtgtgt	gggtgtgtgt	gtttagcgtg	gggttagggga	acgtgtgtgt	2340
	cggtgtgtgt	gggcatgtga	gatgagtgac	tgocgggtgaa	tgtgtccaca	gttgagaggt	2400
	tggagcagga	tgaggggaat	ctgtccacat	caataatcac	ttgtggagcg	ccagctctgt	2460
30	ccaagagccc	acctgggggg	ccagccagga	ccctccactg	gocaggctgc	ctgtgtgtgt	2520
	gttccctgtc	tgtgtccctc	ttgcccgcct	ccctccaaac	taacagggtc	cccccacacac	2580
	agtgccctcc	agaagcagcc	ccctggaggg	agaggaagga	aaatggggtg	ggctgggggt	2640
	ctctccatcc	ctctttttct	cttgcccttg	catggctggc	cttccocctc	aaaaocctcc	2700
	ttccocctgt	ggcagccctc	ttgccaatag	ctgatttttg	ggagggagaa	ggggcgattt	2760
35	gagggagaa	gggagaaagc	ttatggctgt	ctctgggttc	ttccctccoc	agaggttctt	2820
	actgttccag	gggtggccca	gggcaggcag	gggcacacac	atgccttgcc	ccctggtaag	2880
	gtgacccctg	ccattttaaca	gcagccctgg	catgttctct	cccccacagg	atagaaatga	2940
	gggagctcca	gaacttttcc	atcccaaaag	cagtctccct	gggtgaagca	gactggattt	3000
	tgtctctctg	ccctgacccct	gtctccctct	tgagggaggg	gactatgct	aggaatccaa	3060
	ccctagggga	ctgggtggcc	tgcgtcagct	ctttttgata	ctgaaaaact	ttaaggtggg	3120
40	aggggtgcaa	gggatgtgtct	taataaatca	attccaaagc	toactgtg		3167

<210> 63

<211> 2733

45 <212> DNA

<213> Homo sapiens

50

55

<220>
 <221> misc_feature
 <222> (2694)..(2694)
 <223> n=a, c, g or t
 <220>
 <221> misc_feature
 <222> (2724)..(2724)
 <223> n=a, c, g or t
 <400> 63
 agggagaaag gatggcgggc ctggcgggoc ggttggttct gctagctggg gcagggggoc 60
 tggcgagcgg ctcccaggcg gaccgtgagc oggtgtacog ctgactgogta ctgcagtgcg 120
 aagagcagaa ctgctctggg ggogctctga atcacttoog ctcccggcag ccaatctaca 180
 tgagtctagc aggcctgagc ttctgggagc actgtaahta tgagtgtatg tgggtcaocg 240
 ttgggctctca cctccaggaa ggtcacaaag tgcctcaagt ccatggcaag tggccctctt 300
 cccggttctg gttctttcaa gagcggogaa oggocgtggc ctgctttctc aatggcctgg 360
 ccagcttggt gatgctctgc cgtacogcga cctctgtgoc agcctctccc ccaatgtacc 420
 acaactgtgt ggctctggcc ttgggtgtccc tcaatgtcat gtctgtgtcc acagtcttcc 480
 acaccaggga cactgaacct cagagaaat ggaactatct tgtgctctcc gtatcttaca 540
 ctcaatctac ctgtgctgcg tcaggacogt ggggctgcag caccagctgt tggctcaagt 600
 ccttcogggc tctctctgct ctcatgctga cctgtgacgt ctctacctg agcctcatcc 660
 gcttgcacta tggctacaa ctgtgtggca acgtggctat tggcctggtc aacgtggtgt 720
 ggtggctggc ctggtgctgc tggaaocagc ggcggtgccc tcaagtgcgc aagtgcgtgg 780
 tgggtgctct gctgctgcag gggctgtccc tgcctgagct gcttgacttc ccaocgctct 840
 tctgggtctt ggaagccat gccatctggc acatcagcac catctctgtc caagtctctc 900
 ttttcagctt tctgggaagt gacagcgtg accgtctgaa ggaatcagag gacaagttaa 960
 agctggactg agaccttgga cccagctggg gatctgtgg gactctggcc ccgcctgct 1020
 ggctcctct ctcccctcaa cctctgagat gattttctct tttcaacttc ttgaacttgg 1080
 acatgaagga ttgtggccca gaatcatgtg gccagccac cccctgttgg cctcaocag 1140
 ccttgaggtc tgtttatagg aaggctctcc agctcttggg actcgagagt gggcagccoc 1200
 tctactctct ggaactgaat ggggtggaa ctagtgtgtt cttagctcta ccgggaggac 1260
 agctgctgtt ttctctccca ccagctctct ccccaatcc cagctgctct ggtctgggtcc 1320
 tgaagcogtc tctctactcg ggaagaccag gtaccacag ccttagggat acagggggtc 1380
 cctctctgtt accacccccc accctctctc aggaacacac taggtgtgctc tggatgtctg 1440
 ttctttggcc agccaaggtt caocggcatt ctcccaatg ctaggttgag gaccagctgt 1500
 ctgggactgg gaaggagtct caccctgacc gttgcctag ccaaggttccc agggagcctc 1560
 accatactcc ctttcaaggc cagggtctca gcaagccac ggaaggaatc ctgtgctgct 1620
 gtctggttga gagcctgcca ccgtgtgtgc ggaagtgtgg ccaagctgag tgcattagggt 1680
 acagggccgt gagcatgggc ctgggtgtgt gtgagctcag ccttaggtgc cagtggtgga 1740
 gacgggtgtt gtoggggaag aggtgtggct toaaagtgtg tgtgtcagg ggggtgggtgt 1800
 gttagctgtg ctgaggggaa cgtgtgtgoc cgtgtgtgtt ggaatgtgag atgagtgaat 1860
 gcoagtgaaat gtgtccacag ttgagaggtt ggaagcagg cctggggcga cagcagcagg 1920
 aataatcaat ttgtggagoc cagctctgoc ttggtgtcat gctggcctcc cagcagcagg 1980
 cctctcaagg ccaggctgoc ccaacacaa ctgctgggctc ttccctgtct ccaagcctcc 2040
 ctgcaaacct ccagggttcc cctggggctc ttccctctcc gaagcagccc ctggagggga 2100
 gaggagaagaa aatggggatg gctggggctc cctttctctc cctttctctc ttgctctgac 2160
 atggctggcc ttcccttcca aaacctccat tccctgtgct ccaagcctctc tgccatagoc 2220
 tgattttagg tggcaggaag gggcgatttg agggagaag taaggctggg 2280
 tctggtttct tcccttccca gagggtctta ctgttccag gtggccacag gcagcagggc 2340
 cacactatgc ctgcgcctgt gtaaaaggtga cccctgcct tatccagcag ccttggaatg 2400

	ttcctgcctcc	acagggaatag	aatggaggga	gctccagaaa	ctttccatcc	caaaggcagt	2460
	ctcgttggtt	gaagcagact	ggatttttgc	ctgcctcctg	accccttgct	ccctcttgag	2520
	ggaggggagc	tatgtctagga	ctccaaacct	agggactcgg	gtggcctcgg	ctagctcttt	2580
5	tgatacttga	aaacttttaa	ggtggggagg	tggaaggga	tgtgttaag	ggcgccgaa	2640
	ttcaaaaagc	ttctcagag	tactcttaga	goggccggcg	gccatctgat	tttccacccc	2700
	gggtggggta	ccaggttaag	tgttccccat	atc			2733
	<210>	64					
10	<211>	2546					
	<212>	DNA					
	<213>	Homo sapiens					
15							
	<400>	64					
	aagctcctcc	cccgcgggcg	agccaggggg	aaaggatggc	cggcctggcg	gcgcgggttg	60
	ctcgtctagc	tggggcagcg	gcgctggcga	ggcgctccca	ggggagccgt	gagccgggtg	120
20	acgcggactc	cgtactgcag	tgccaaggagc	agaaactgtc	tggggcgogt	ctgaaactact	180
	tcgcgtcccg	ccagcacaac	tacatagagc	tagcaggctc	gaactgtcgg	gcagactgta	240
	agtatagagt	tatgtgggtc	acgcgtgggc	ctacactcca	ggaaggtcac	aaagtgcctc	300
	agttccatgg	caagtggccc	ttctcccggt	tcctgttctt	tcaagagcgc	gcactggcgc	360
	tgccctggtt	ctccaaatgg	ctggccagcc	tgtgtatgct	ctgcgcgtac	cgccactctg	420
25	tgccagcctc	ctcccccctg	taccacacct	gtgtggcctt	cgccgtgggtg	tcctccaatg	480
	catgtgtctc	gtccacagtc	ttccacacca	gggacactga	ctccacagag	aaaaatggact	540
	actctgtctc	ctccactgtc	atctctacct	caactctact	gtgctgctgc	aggccttggtc	600
	aaactgtgtg	ggtggctggc	ctgggtgctg	tggaaaccgc	ggcggtctgc	tcactgtgcg	660
	aaagtgcgtg	tgtgtgtctt	gctgtgcag	ggcctgtccc	tgtctgagct	gcttgacttc	720
	ccacgcgtct	ctctggctct	ggatgcccac	gccatctggc	acatcaagac	catctctgtc	780
30	caactctctc	ttttccagct	ctctggagat	gacagcctgt	acctgtctga	ggaatcaag	840
	gacaaagttc	agctggagct	aagacacttg	acgcagctct	ccccagtggt	gatctgtccc	900
	ccgcctcgtc	ggctccctct	ctcccccctc	cccttgagat	gatctctctc	tttccaaactc	960
	ttgaacttgg	acatgaaggc	tgtgggcccc	gaatcatgtg	gccaagccccc	ccctgtgttg	1020
	ccctccccc	cccttgagct	tgttccaggg	aaggccctcc	agcatctggg	actctgagag	1080
35	gggcagcccc	ctctacccct	ggagctgaac	tgggttgga	ctgagtggtt	ctctagctct	1140
	acgcggggga	cagctgcctg	ttctccccc	accagcctcc	tccccccctc	ccagctgtcc	1200
	tggctgggtc	ctgaagccct	ctgtctacct	gggagacagc	ggacccaggg	cccttagggt	1260
	acaggggggc	ccctctctgt	accaccccc	acccctctcc	aggacccac	taggtgtgtc	1320
	tggtatgctt	ttctttggcc	agccaaaggt	ccagggagat	ctcccccctg	gatcttgagg	1380
40	gacaaagctg	ctgggattgg	gaaggagttt	ccccctgacc	gttgccctag	ccaggttccc	1440
	aggagggcct	accatactcc	ctttccaggg	cagggtctca	gcaagccccc	ggcagaggtc	1500
	ctgtctgctc	ctctgttgga	gagcctggcc	cgtgtgtgtg	ggagtgtgtg	ccaggtctgg	1560
	tgcataggtg	acagggcgct	gagcctgggc	ctgggtgtgt	gtgagctcag	gcctaggtgc	1620
	gcagtggtga	gaggggtgtt	ctgggggaag	aggtgtgtgt	tcaagtgctg	tgtgtgcagg	1680
	gggtgggtgt	cttagctggg	gttaggggaa	cgtgtgtgtg	cgtgtgtgtg	ggcctgtgag	1740
45	atgagtgact	gcgcgtgaa	gtgtccacag	ttgagaggtt	gagcagagat	gaggtgaatc	1800
	tgttccatcc	aaataactact	tgtggagcgc	cagctctgct	caagagccca	ccctggggcg	1860
	cagccagggg	ctctccatgg	ccaggtctgc	tgtgtgcatg	ttccctgtct	gggtccctgt	1920
	tgcccccctc	ctccaaacct	ccaggggtcc	ccacacacca	gtgcctccca	gaagcagccc	1980
	ctcggaggga	gagggaaggc	aatggggatg	gctggggctc	ttccatctct	ccctttccct	2040
50	ttgctctgca	atggctggcc	ttcccccctc	aaactctcat	ctccctgtgc	ccctccctct	2100
	tgccatagcc	tgatttttgg	gaggaggaag	ggggatctgt	agggagaggg	ggagaaagct	2160
	tatgtgttga	cttggtttct	ttccctccca	ggggctctta	ctgtccagag	gtggccccc	2220
	ggcagggcag	ggccacaccta	tgcctgccc	ctggttaagg	tgacccctcc	caattaccag	2280
	cagccctggc	atgttctcgc	ccccacaggc	tagaatggag	ggagctccag	aaacttccca	2340
	ttccaaaagg	agttctcgtg	gttgaagcag	actggaatct	tgtctgtccc	ctgacccctt	2400
55	gtccctcttt	gagggagggg	agctatgcta	ggactccaac	ctcagggact	cgggtgggct	2460

	gogctagcctt	otttttagatac	tgaaaaacttt	taagggtggga	gggtggcgaag	ggatgtgctt	2520
	aataaatcaaa	ttccaagcct	cacctg				2546
5	<210>	65					
	<211>	2683					
	<212>	DNA					
10	<213>	Homo sapiens					
	<400>	65					
15	aagctcctcc	cccgggcgog	agccaggggag	aaaggatggc	oggcctggcg	gcgcgggttg	60
	tcctgtctag	tggggcgagc	gogctggoga	goggcctccca	ggggcgacgt	gagccgggtgt	120
	acogogaact	ogtaactgcag	tgcgaagagc	agaactgtctc	tggggcgogct	ctgaatcaact	180
	tcogctccog	ccagcccaatc	tacatgagtc	tagcaggctg	gacctgtcg	gaogactgta	240
	agtataggtg	tatgtgggtc	acogttgggc	tcctacctcca	gggaagtgac	aaagtgtctc	300
	agttccatgg	caagtggccc	ttctcccggt	tcctgtttct	tcgaagagcg	gcacogggcg	360
20	tggtccgtgt	ttcoaatggc	ctggccagcc	tgggtatgct	ctgocgtatc	cgccactctg	420
	tgcgaacctc	ctcccccatt	taccacacct	gtgtggcctt	ogcctgggtg	tcctccaagt	480
	catgggtctg	tgcacagtc	ttccacacca	gggaacctga	ctccacagag	aaaaatggact	540
	actttctgtc	ctccactgtc	atctacacct	caatctacct	gtgtgtgtgc	aggacogtgg	600
	ggctgcagca	ccagctgtgt	gtcagtgtct	tcogggctct	ctgtctgtct	atgtctgacg	660
25	tgcactcttc	ctacactgagc	ctcatccgct	tgcactatgg	ctacacacct	gtggccaaag	720
	tggtctatgg	ccctggccaac	gtgtgtgtgt	ggctggcctg	gtgctgtgtg	aaocagggcg	780
	ggctgcctca	ogtgcgcaag	tgcgtgtgtg	tggtctgtct	gtgtgcaggg	ctgtccctgc	840
	tgcagctgct	tgcactccca	ccgtctctct	gggtccctga	tgcccatgac	atctggcaca	900
	tcagcaacct	ccctgtccac	gtctctcttt	tcagctctct	ggagatgac	agcctgtacc	960
	tgtctgaagg	atcacagagc	aagtccaagc	tggactgaag	acottggagc	gagctgtccc	1020
30	caagtgggat	ccctgcccgc	ccctgtctgc	ctccctctct	ccctcaaac	ttagatgat	1080
	ttctctcttt	caactctctg	aacttggaca	tgaaggatgt	gggcccagaa	tcattgtggc	1140
	agcccccacc	ctgttggccc	tcaccagct	tggagtctgt	tcctagggaag	gcctcccgac	1200
	atctgggact	cgagagtggt	cagccctct	acotccctga	gttgaaactg	ggttgaactg	1260
	agtggtctct	tagctctacc	gggaggacag	ctgctgtgtt	ccctccccc	agccctccct	1320
35	ccacatcccc	agctgtccgt	ctgggtccct	aagccctctg	tcctccctgg	agaccaggga	1380
	ccacaggcct	tagggatata	gggggtccct	ttctgttacc	ccctccccc	ccctccagg	1440
	acacacactg	gtgtgtctgt	atgtctgttc	tttgccacg	caaggttcc	ggcagttctc	1500
	cccatgggat	cttgagggac	caagctgtct	ggattgggaa	ggagtttcc	ccctgaactg	1560
	gcctatgcca	gggtccacag	aggcctccac	atactccctt	tcaggggcag	ggctccagca	1620
40	agccccaggg	aaggatctct	tgtctgtgtc	tggttgagag	octcccaac	tggttcggga	1680
	gtgtggccca	ggctgtgtgc	ataggtgaca	gggcctgtg	catgggocgt	ggtgtgtgtg	1740
	agctcagccc	taggtgcgca	gtgtggagac	gggtgtgttc	ggggaagagg	tgtgtgttca	1800
	aagtgtgtgt	gtgcaggggg	tgggtgtgtt	agcgtgggtt	aggggaaact	gtgtgcogct	1860
	gctgtgtggc	atgttagatg	agtgactgac	gttgaaatgt	tcacacagtg	agaggttgga	1920
	cgaggatgag	ggaatccctg	cccatccaat	aatcaactgt	ggagcgccag	ctctgcccca	1980
45	gaocccacct	gggcggagag	ccaggagctc	tcactggcca	ggctgcctgt	gtgcagtgtc	2040
	ccgtgtctgt	gcocctttgc	ccgctccctg	caaacctcc	agggctccca	caacacagtg	2100
	ccctccagaa	gcagcccttc	ggaggcagag	gaaggaaact	ggggatggct	ggggctctct	2160
	ccatccctct	ttctccctgt	ccctccagtg	gctgcctctc	ccctccaaaa	ccctccattc	2220
	ccctgtccca	gcocctttgc	catagctgta	ttttggggag	gagggaaggg	cgatttgagg	2280
50	gagaaggagg	gaaagcttat	ggctgggtct	ggtttctctc	ctccacagag	gtgtcttaact	2340
	ttccagggtg	gcgccagggc	aggcaggggc	ccacacttat	ctgcgocctg	gttaaaggta	2400
	ccctcccaat	ttaccagcag	ccctggccat	ttctggcccc	gaagcagact	aatggaggga	2460
	ctcccgaaaa	ctttccatct	caaaagccag	ctcgtgtgtt	tatgtctagg	ggattttctc	2520
	tctgcocctg	acccctgttc	ccctctttag	gtgggggagc	aaactcttaa	ctccaaactc	2580
55	agggactggg	gtggcctggg	ctagctctct	ttgatactga	ggtggggagg		2640
	tggcaaggga	tgtgtctaat	aaatcaatto	caagctcac	ctg		2683

<210> 66

<211> 2341

<212> DNA

<213> Homo sapiens

<400> 66

aaagcttccctc cccggcggcg agccaggggag aaaggaaggc oggcctggcg ggcgggttgg 60
 tootgctagc tggggcagcg gcgctggcga gcgctccca gggcgacgt gagcgggtgt 120
 acogcgactg cgtactgcag tgcgaagagc agaactgcctc tggggcgct ctgaatcact 180
 tccgtctccg ccagcccaatc tatgtgggtc aocgttgggc tctacctcca ggaaggtoac aaagtgcctc 240
 agtatgagtg taatgtgggtc tctcccggtt tootgttctt tcaagagcgc gcattggccg 360
 tggcctcgtt tctccatggc cttggccagcc tgggtgatct ctgcgcctac cgcactcttg 420
 tgcacacctc cttcccccag taaccacacct gttgtggcctt cgcctgggtg tccctcaaatg 480
 catgtgtctc tgcacagtc ttcacaccca gggcacactg cctcacagag aaaaaggact 540
 actctgtgtc cttccactgc atctacacct caatctacct gtctgtcgtc agcttctctg 600
 aagatgacag cctgtacctg ctgaagggaat cagaggacaa gtccaagctg gactgaagac 660

cttggagaga gtctgccccg gtggggatcc tgcctccccc ctgctggcct cctctctccc 720
 ctcaaccctt gagatgattt tctcttttca actctctgaa cttggacatg aaggatgttg 780
 gcccagaatc atgtggccag cccacccctt gttggccctc accagccttg gactctgttc 840
 tagggaaggc cttccagcat ctgggactcg agagtgggca gccctctac cttctgggac 900
 tgaactgggg tggaaactgag tgtgttctta gctctacogg gaggacagct gactgtttcc 960
 tcccccagag cctcctccccc acatcccccag ctgctggcct gggctctgaa gccctctgtc 1020
 taacttggag accaggggacc acaggccctta gggatacagg gggctccctt ctgttaacca 1080
 ccccccacct cctccagagc accaactaggt ggtgctggat gctgttctt tggccagcca 1140
 aggttcaagg ogattctccc catgggatct tgaagggaaca agotgtgtgg attgggaagg 1200
 agtttcaacc tgaccgttgc cctagccagg ttcaccagg gctcaacct actcctttc 1260
 agggccaggc cttccagcag cccagggcga ggaatctgt ctgtgtctg gttgagagcc 1320
 tgcacccgtg tgtoggggat gtggggcagg ctgagtgcat aggttgacag cctgtgagca 1380
 tgggcctggg tgtgtgtgag cttcaggccta ggtggcaggt ggtgtgttag cgtgggttag 1440
 gggaaggggt tggcttcaaa tgggtggcat gtgagatgag tgactgcagg tgaatgtgtc 1500
 cacagttgag aggttggagc aggatgaggg aatctgttca ccatcaataa tcaactgttg 1560
 agogccagct ctgccccaga cgcacacctg ggggacagcc agggagctct catggccagg 1620
 ctgctgtgtg gcatgttccc tgtctgttgc cctcttgcac gccctcaaca aacctcaacg 1680
 gtgtcccca caacagtgcc cttccagaagc agccctctgg aggcagagga aggaataatg 1740
 ggatgtctgg ggcctctctc atctcctttt tctccttggc tctccttggc tggccttccc 1800
 ctccaaaaac tccattccccc tgcctgcagc cctcttgcca tagcctgatt ttggggagga 1860
 ggaaggggcg atttgaggga gaaggggaga aagcttatgg ctgggtctgg ttcttccctt 1920
 tcccagaggg tcttactgct ccaagggtgac cccagggcag cagggggcca caactgccc 1980
 aggaatagaa aaaggtgacat cctgcacatt accagcagcc ctggcatgtt cctgcccacc 2040
 agcagactgt tggaggggag tccagaaact tccatcccca aaggcagctt cctgtgttga 2100
 gctaggact atttttgttc tgcctctgac cctctgtccc tctttgaggg aggggagcta 2160
 acttttaagg gggggagggt ggaactgggt cgtctgtcgt agcttctttt gataatgaaa 2220
 g 2280
 2340
 2341

<210> 67

<211> 2109

<212> DNA

<213> Homo sapiens

<400> 67

gattggcgcg	gagctgccag	cggggaggct	gcagcgcggg	gttgttacag	ctgtctggagc	60
agcagcggcc	cccgctcccg	ggaaocgttc	ccgggcogtt	gatctctggc	ccacacagaa	120
cagcagagag	gggcagcagg	atgaatgtgg	gcacagcgca	cagcagagtg	aaocccaaca	180
cgccgggtgat	gaacagccgt	ggcaatctggc	ttctctactgt	ctgtggccatc	ggctctctccc	240
acatctgtct	gctgagcatc	ccgtttgtga	gtgtctctgt	ctgtctggacc	ctcacaaccc	300
taattccaca	catgggcatg	tatatctctc	tgcaacgggt	gaagggggaca	ccctttgaga	360
cccgccagca	gggcaaggcg	aggctgtctaa	cccactggga	gcagaaggat	tatggggtcc	420
agttcaacggc	ctctcgggaag	ttcttgacca	tcacacccat	cgctctgtac	ttctctccaca	480
gctctacacac	taagtacgac	cagatccatt	ttgtgtctcaa	caocgtgtcc	ctgatgagcg	540
tgcttatccc	caagctgcgc	cagctccacg	gagtcoggat	ttttggaaac	aataagtaact	600
gagagtgcag	ccctctccccc	tgcccagggt	ggcaggggag	gggtagggtta	aaaggcatgt	660
gctgcaacac	tgaagacaga	aagaagaagc	ctctgggacac	tgccagagat	gggggttgag	720
ctctctggcct	aatttccccc	ctcgtctccc	ccagtagcca	acttgagta	gctgttagtg	780
gggtctgggt	aggcccccctg	ggctctgacc	ttttctgaat	tttttgatct	cttctcttttg	840
ctttttgaaat	agagactcca	tggaattgtgt	catggaattgg	gctgggctcc	tggttgcaac	900
atggcaacag	cagttgcgac	aggaggccag	gggaaaaacc	ccgtctcaact	tggttgccct	960
caggcaagcca	aagcaacttta	acccctgcac	agggagcaga	gggcgggtacg	gctctctggat	1020
tgtttcaactg	tgattctctag	gtttttttoga	tgccatgcag	tggtgtgcttt	tgctgatgga	1080
agcaagtgtg	ggatgggtct	ttgctctctc	gggtaggggag	ctgtcttaact	caagtcocag	1140
gcttttggca	gctctctctgc	aacccacccg	gggtccctgg	tgggagtgagg	gagggtccagg	1200
ttggggaaag	atggggtaga	gtgtagatgg	cttggttcca	gaggtagagg	ggccagggctt	1260
gctgcacatc	tgccctgtgtg	gaggttgggg	agctgtaggga	gagctagtga	gtcgagactt	1320
agaagaatgg	ggccacacag	cagcagaggga	ctggtgttaag	ggagggagggg	gtagggacag	1380
aagctagacc	caatctctctt	tgggatgtgtg	gcaggggagg	aagcaggctt	ggagggttaa	1440
tttaccocaca	gaatgtgata	gtaatagggg	agggaggctg	ctgtgggttt	aaactcctggg	1500
ttggctgttg	ggtagacagg	tggggaaaag	gcccgtgagt	caatgttaagc	acaggtccaca	1560
cttggccctg	actcctcggg	gggtatgggg	aagctgtgac	tgccactggg	cagggcctct	1620
tcctctgag	gcccctacccc	cttaactctc	tcactgcagac	tggggtctgt	ggtgagagtc	1680
catgtggcag	ccacatgtgg	cgttgtgagg	ccaccccatg	gaactgtgct	gtagggaactg	1740
ctgtaggatc	ccctgtccaag	cagcaccagag	gaaggggcaa	ccctggcctg	agccttctcc	1800
ctctccactg	cagagaagaa	agtgaggccg	gagccctgag	gtgtgggtcc	ctgtgtctct	1860
ctcccccaggt	tggaactagg	gcagtgttaa	ttttgaaaag	ttttctccct	actgaccctg	1920
ttccagctgt	ccaaagggaac	aggagaggct	actgggctgt	aatctttcta	agcaaatgtgt	1980
catctccacac	ccgtgtgtatc	atagggaact	ttccacttaa	gcgaataaaa	agggcaacat	2040
gaataggatt	tttactccct	ttgtacagta	ttctgaggaa			2100
gtttctgttt						2109

<210> 68

<211> 2423

<212> DNA

<213> Homo sapiens

<400> 68
gagagccogag ctagogacga gcaagtgttg ogggcgcggg ogcgcgggga ggttggtgag 60
gcgtagcoggg agccagagag tctcttgttc cegtcccaag gtccggcggt caccctcccg 120
gcgcgcagctg ccggtcccggt aactcccggg cctgtccctg gcccccgggt tctgcactcc 180
5 gctgcgcogga gcgcgcggcc ogggcggcac cgcgcggccc catgaggagg gacgtgaacg 240
gagtgacocaa gagcaggttt ctaggtgtct caaatagtga tgaagctgta atcaataaaa 300
aacttccocaa agaactcttg ttacggatat tctcttttct tctgttgttt agatgtgtt accctgtgoc 360
gctgtgtacaa ggttccocagg gcoctggaaat ttctggctct ggaatggcagt aactggcagc 420
10 gaattgtcct gaattgtact atttgtattc ctaggggata ttctggcgcg agtagtgagg aatatttcaa 480
aacgatgtgt gggctctttta cgaaaagtta gctcttgttg atgtcttgga gtgggagaca 540
atgcaataag aacacttgca gaaaactgaa gtcaacattg aaactaactg ctaaattgggt 600
gtacaaagac aacagacaga acagtgtact cgccttagca gttctgttcc aaactaaggc 660
aactgtacct gactctctgt acatcaataa caaacattgt tctaaagct ctagtgagg 720
ga tgtcactt gttggagcag ttgaacattt cctggttgta ccaagttaacc aaggaatggca 780
15 tccaagcaat agtgaggggg tgtgggggtg taaggacttc aagtacatag gtgcacactg ccttgaaatg 840
agctagaaga ttgtctgcga atcacagatg gcoctgtgct tgtccaaact cactataatg agaggtgoc 900
aactgtcgaac ttgtctgcga atcacagatg gcoctgtgct tgtccaaact cactataatg agaggtgoc 960
ataagtatac atcoctttgt cggcttagaa gctccaaact taatggaaat ggcaagatgt tccaattaa 1020
ctctagtgtca gaactgcocaa ctagocaggga attgccaatg acttgaaaag atggaactgt 1080
20 cagatgtggg cgtttaccact ctagocaggga attgccaatg acttgaaaag atggaactgt 1140
aagagttgtt tcagataaaca gatgcacat taatccaaact tctatacaac tbtctctcag 1200
ttoaagtatt gagtctgtct ctagctgagc ctagctgagc ctagctgagc gctggacaac ttaccactaa 1260
ggagtgtggg ctagccoccat ctagccoccat ctagccoccat ctagccoccat ttaccactaa 1320
25 tcaagatga atccctggag ctagctgaaga ctagccoccat ctagccoccat ttaccactaa 1380
ttgaactgca gcaaaataca ogggtgtgga atccoccat ctagccoccat ttaccactaa 1440
ttaaagtcca cgcacttct ctagccoccat atccacttca gcaacatccc atccacttca 1500
agcgctctgt cagatgtctc agagatccag tggagtctgt cgaatgtctc aga tcccag agccacacat 1560
gagtatttaa tgacacttct aggtaggagg atccactgct tgaatgtctc ggaactgaat 1620
30 gttctaggaa aggggttaaa agccoccatc taataattta ctaattgtct tgaagtgtcc 1680
acatacacat acacacocct accoccatc taataattta ctaattgtct tgaagtgtcc 1740
ttgtgatggc catagggcaa gataggtgg gcocttagtg ggaactccag caaaocccgt gttaatgtct 1800
ataagaaaaa taatattttt ctaggggggt catcttagtg ggaactccag caaaocccgt gttaatgtct 1860
tbtgtttttt taaatttttt catcttagtg ggaactccag caaaocccgt gttaatgtct 1920
tctctccggg tcaagaaaaa catcttagtg ggaactccag caaaocccgt gttaatgtct 1980
35 aaaaatttct tgaattttt aatgtgtaga tgcctagcat gactcaaatg aaggtctgtc cttctatgtc 2040
tctctgtttt catggoccaa tgcctagcat gactcaaatg aaggtctgtc cttctatgtc 2100
gtgtttttat atttggocaa gagggttctt aaatgttttg cttctatgtc cttctatgtc 2160
aatgttaatt ttacacactt aaaaatttct aaatatttcc cttctatgtc cttctatgtc 2220
cattctattt tgcctccccc cacaacttct tgcctatgtc cttctatgtc cttctatgtc 2280
tcaactcaat gtacacactt tgcctatgtc cttctatgtc cttctatgtc cttctatgtc 2340
40 tggggaattt caaaactttt ttaaanaaat attatacta atattttttt taattctaaa 2400
ttttaaanaa aaaaaaaa aaa 2423

<210> 69
<211> 1841
46 <212> DNA
<213> Homo sapiens

<400> 69
agctggggacc ggaggggtgag ccgggcagag gcagagacac acggggagag gaggagaggc 60
tgagggggagg agctgggagaa ggagggggaga actcagagag cggagagacac cgagagccag 120
agacacccagg gcggggggag gcggggggagc gaggcagaga cctggctcag cgagcgcggg 180
5 ggagagaccc cgaactccga gacgctcggg gcggcggcag ccggggcgag cgaactccca 240
agctggggacc ggaggggtgag ccgggcagag gcagagacac acggggagag gaggagaggc 60
tgagggggagg agctgggagaa ggagggggaga actcagagag cggagagacac cgagagccag 120
agacacccagg gcggggggag gcggggggagc gaggcagaga cctggctcag cgagcgcggg 180
5 ggagagaccc cgaactccga gacgctcggg gcggcggcag ccggggcgag cgaactccca 240
agctggggacc ggaggggtgag ccgggcagag gcagagacac acggggagag gaggagaggc 60
tgagggggagg agctgggagaa ggagggggaga actcagagag cggagagacac cgagagccag 120
agacacccagg gcggggggag gcggggggagc gaggcagaga cctggctcag cgagcgcggg 180
5 ggagagaccc cgaactccga gacgctcggg gcggcggcag ccggggcgag cgaactccca 240
agctggggacc ggaggggtgag ccgggcagag gcagagacac acggggagag gaggagaggc 60
tgagggggagg agctgggagaa ggagggggaga actcagagag cggagagacac cgagagccag 120
agacacccagg gcggggggag gcggggggagc gaggcagaga cctggctcag cgagcgcggg 180
5 ggagagaccc cgaactccga gacgctcggg gcggcggcag ccggggcgag cgaactccca 240
agctggggacc ggaggggtgag ccgggcagag gcagagacac acggggagag gaggagaggc 60
tgagggggagg agctgggagaa ggagggggaga actcagagag cggagagacac cgagagccag 120
agacacccagg gcggggggag gcggggggagc gaggcagaga cctggctcag cgagcgcggg 180
5 ggagagaccc cgaactccga gacgctcggg gcggcggcag ccggggcgag cgaactccca 240
agctggggacc ggaggggtgag ccgggcagag gcagagacac acggggagag gaggagaggc 60
tgagggggagg agctgggagaa ggagggggaga actcagagag cggagagacac cgagagccag 120
agacacccagg gcggggggag gcggggggagc gaggcagaga cctggctcag cgagcgcggg 180
5 ggagagaccc cgaactccga gacgctcggg gcggcggcag ccggggcgag cgaactccca 240
agctggggacc ggaggggtgag ccgggcagag gcagagacac acggggagag gaggagaggc 60
tgagggggagg agctgggagaa ggagggggaga actcagagag cggagagacac cgagagccag 120
agacacccagg gcggggggag gcggggggagc gaggcagaga cctggctcag cgagcgcggg 180
5 ggagagaccc cgaactccga gacgctcggg gcggcggcag ccggggcgag cgaactccca 240
agctggggacc ggaggggtgag ccgggcagag gcagagacac acggggagag gaggagaggc 60
tgagggggagg agctgggagaa ggagggggaga actcagagag cggagagacac cgagagccag 120
agacacccagg gcggggggag gcggggggagc gaggcagaga cctggctcag cgagcgcggg 180
5 ggagagaccc cgaactccga gacgctcggg gcggcggcag ccggggcgag cgaactccca 240
agctggggacc ggaggggtgag ccgggcagag gcagagacac acggggagag gaggagaggc 60
tgagggggagg agctgggagaa ggagggggaga actcagagag cggagagacac cgagagccag 120
agacacccagg gcggggggag gcggggggagc gaggcagaga cctggctcag cgagcgcggg 180
5 ggagagaccc cgaactccga gacgctcggg gcggcggcag ccggggcgag cgaactccca 240
agctggggacc ggaggggtgag ccgggcagag gcagagacac acggggagag gaggagaggc 60
tgagggggagg agctgggagaa ggagggggaga actcagagag cggagagacac cgagagccag 120
agacacccagg gcggggggag gcggggggagc gaggcagaga cctggctcag cgagcgcggg 180
5 ggagagaccc cgaactccga gacgctcggg gcggcggcag ccggggcgag cgaactccca 240
agctggggacc ggaggggtgag ccgggcagag gcagagacac acggggagag gaggagaggc 60
tgagggggagg agctgggagaa ggagggggaga actcagagag cggagagacac cgagagccag 120
agacacccagg gcggggggag gcggggggagc gaggcagaga cctggctcag cgagcgcggg 180
5 ggagagaccc cgaactccga gacgctcggg gcggcggcag ccggggcgag cgaactccca 240
agctggggacc ggaggggtgag ccgggcagag gcagagacac acggggagag gaggagaggc 60
tgagggggagg agctgggagaa ggagggggaga actcagagag cggagagacac cgagagccag 120
agacacccagg gcggggggag gcggggggagc gaggcagaga cctggctcag cgagcgcggg 180
5 ggagagaccc cgaactccga gacgctcggg gcggcggcag ccggggcgag cgaactccca 240
agctggggacc ggaggggtgag ccgggcagag gcagagacac acggggagag gaggagaggc 60
tgagggggagg agctgggagaa ggagggggaga actcagagag cggagagacac cgagagccag 120
agacacccagg gcggggggag gcggggggagc gaggcagaga cctggctcag cgagcgcggg 180
5 ggagagaccc cgaactccga gacgctcggg gcggcggcag ccggggcgag cgaactccca 240
agctggggacc ggaggggtgag ccgggcagag gcagagacac acggggagag gaggagaggc 60
tgagggggagg agctgggagaa ggagggggaga actcagagag cggagagacac cgagagccag 120
agacacccagg gcggggggag gcggggggagc gaggcagaga cctggctcag cgagcgcggg 180
5 ggagagaccc cgaactccga gacgctcggg gcggcggcag ccggggcgag cgaactccca 240
agctggggacc ggaggggtgag ccgggcagag gcagagacac acggggagag gaggagaggc 60
tgagggggagg agctgggagaa ggagggggaga actcagagag cggagagacac cgagagccag 120
agacacccagg gcggggggag gcggggggagc gaggcagaga cctggctcag cgagcgcggg 180
5 ggagagaccc cgaactccga gacgctcggg gcggcggcag ccggggcgag cgaactccca 240
agctggggacc ggaggggtgag ccgggcagag gcagagac

	ccgcctccgc	gcoctccct	aggggggc	ggtattttta	tcogtgogog	aacagccctc	360
	ctcctcctct	ogcgcacag	ccgcgcgcct	gcgcggggga	gcccagca	gacgcgcgc	420
	gggacccgc	gtgcgcgc	ccgcgcgc	ogccaccc	gcgcgcgc	gacccacag	480
5	acgcacagaa	gacacagtt	tcggtgccgc	ogccccctag	ccagctcgac	ccgcgcgcag	540
	tgagagtgat	ccgcgcgcag	agacccacgc	ctgcctatgt	gttcgcgc	tcagagcact	600
	ctcaccagca	ggaggaagcc	tccccccac	agagagcc	agggagggg	ccacctc	660
	agtcgaagag	acccaccc	tgctcctaca	ccccccttc	gctgaaagct	gtgcgcgc	720
	ttgtgtagtc	tcacctgcag	tcctacagca	atttgaaaga	gaacccagcc	tcagagggag	780
10	aggaagagtc	gggggagctt	ogggagctgg	gttatccaa	agaggaagat	gagggagga	840
	aggaagagtc	tgaggaagag	gaaggaagag	aggacagca	ggctgaagtc	ctgaaggtc	900
	tcaggcagtc	tgctgggcga	aagacacct	gtggccaggg	tgaggaggg	ccctgggagc	960
	gcccacccc	tcggatgcag	tcggagagag	atggaggtc	tgaggacaa	gtggaagacc	1020
	cagcactaag	tgagcctggg	gaggaaacct	agcccccct	ccctctgag	ccctggcact	1080
	agggccacag	ctgcctcct	ccaggagga	gtggagggga	catcctgtt	ccccagaaac	1140
15	ccactctatc	ctcccccct	tttgtctct	tcctcctgc	tgctagggct	goggtctct	1200
	actctataga	gactaaggt	ggtctgtgt	tgctgtgtt	cccccctgc	agtcctct	1260
	agagacccct	ggcactgtct	gctcgtatgc	cccccctgc	agtcctct	ccctcacc	1320
	agggggaggt	gggagtgtg	ccgcgcgc	ctggaaaac	cagaaaaag	ggaaacggga	1380
	tttgccttc	acactctac	tcccacagtc	tcctccctg	gaacacggag	cccccacggg	1440
20	caggacccct	agactctggg	aaaggaggtc	ctgagaaact	tgaggtaacc	ttagatctt	1500
	ttctacccac	ttctctatgg	aggattccaa	ctcaccacgt	ctcaccacgt	ctctacacag	1560
	ggtccaggac	taaggcgttt	ttctccatg	ccccaacatt	ttgggaatct	tcctctaatc	1620
	acccctgtct	ccctgtgtg	ccctggagat	ggactggcag	agacccctt	gtgtgtgtt	1680
	gtgctttgat	gcccaggaatg	ccgcctagtt	tatgtccccc	gtgggggaca	cagggggggg	1740
	cgccaggttt	tccttgtccc	ccagctgtct	tgcccccct	ccctctctcc	ctgactccag	1800
25	gctggaaccc	ctccctgtct	gtaataaact	ttgttaaata	a		1841
	<210>	70					
	<211>	748					
30	<212>	DNA					
	<213>	Homo sapiens					
35	<400>	70					
	ggccgcgcag	aggggggagc	oggggcagac	gtccgtagcg	cccccctccg	aggaggtoga	60
	gcccggcagc	ggggctccgc	tcgtgtgtga	gtactgtgaa	ccctgcgcct	tcagggcgac	120
	ctacctggag	ctggccagctg	ctgtgaagga	gcagtatccg	ggcctcgaga	tcgagctogog	180
40	ccctgggggc	acaggtgcct	ttgagataga	gataaatgga	cagctgggtg	tcctccagct	240
	ggagaaatgg	ggcttccct	atgagaaaga	tcctattgag	gcccctcga	gagccagtaa	300
	tgagaaaccc	ctagaaaaga	tcacccacag	ccgtccctcc	tgctcctc	tgtagctgca	360
	caggaactct	ggctcctgct	ctgttctggg	gtccaaacct	tggtctccct	ttggtcctgc	420
	tgaggactcc	ccctgcctct	tcccctcact	tagctcctta	gcacagagac	ccctggccctc	480
45	actttgcctc	ttgggtacaa	agaaggaata	gaagattccg	tgcccttggg	ggcagggagag	540
	agacactctc	catgaaacct	ctccacagca	ccctacaccc	ccctccacag	gtaagtgccc	600
	acgaaagccc	atgcacactc	tcgcctcgtg	aatccctgtc	tgatgcacac	gattttatct	660
	attctccctc	aaacccagggc	aatgtcagct	attggcagta	aagtgccgct	acaaacacta	720
	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa				748

<210> 71

<211> 795

<212> DNA

<213> Homo sapiens

<400> 71

taoggtgtgog	agaagacgac	agaagctaga	cccaatctcc	tttgggatgt	gggcaggag	60
ggaagcaggc	ttggagggtt	aatttaccca	cagaatgtga	tagtaaatgg	ggaggggagc	120
tgatgtgggt	tttaactctg	ggttggctgt	tggttagaca	ggtgggga	aggccogtga	180
gtcatgttaa	gcacaggctc	aacttggccc	tgactcctgc	gggggtatgg	gggaagtgtg	240
acagaaacga	tgggtgtctgt	ggtcattctg	aggccctcac	cccttaactt	cctcatcacg	300
actggcactg	ggcaggggcct	ctcatgttgc	agccacatgt	ggcgttttga	ggccacccca	360
tgtgggggtc	gtggtgagag	tcctgttaga	tcctgtctca	agcagcacag	aggaagggcg	420
aagactgtgc	ctgttaggcac	tgtctcagcc	tgccagagaag	aaagttaggc	cgggagcctg	480
agcctggggt	ggagcctctc	ccctccccc	gttggactag	ggcgagtgtt	aattttgaaa	540
aggttggggt	ccctgtgtcc	tcttccaggg	gtccaggga	acaggagagg	tcactggggc	600
tgttttctcc	ccctgagccc	tgcattctcc	accccggtga	tcattagggaa	ctttcactct	660
aaaaattctt	taagcaaaat	gtgaatagga	tttttaactc	ctttgtacag	tattctgaga	720
aacgcacata	aaagggcacc	atgtttctgt	taaaaaaaaa	aaaaagtacg	caaaaaaaaa	780
aaaaaaaaaa	aaaaaa					795

<210> 72

<211> 2356

<212> DNA

<213> Homo sapiens

<400> 72

ggcacgaggg	oggaagtgc	ctctagaggg	gtgggtgaaa	tggcagttga	oggtcctgtg	60
gaatagatcc	cgogaggtag	cccccgaact	atttctctac	gtttctctct	gatcctcccg	120
aaatcttcca	gatoocgtga	gtgaggaatc	gtctccaccc	tcattggggg	ogggagacctg	180
aatctgaaga	agagctggca	cccgacgaac	ctcagggaatg	tggagaaagt	gtggagagcc	240
gagcagaagc	atgagctgga	cggaagaag	attgaggagc	ttcagcggga	gctcgagagaa	300
gagcagagcc	gggaagagat	gcagggctat	cgggaggatg	tggggccgct	caagaaaaaa	360
gaagaaaagt	tggactggat	gtaccagggt	ctgggtggga	tgggtgaacg	tgaccagctac	420
ctgctggggc	gcccatttga	caaatatgtt	tttggaaga	tggaggagaa	ggaggcaggc	480
tgtcttctgt	aaaacaggact	tctccacggc	tctattcttt	cccatcagg	tgccaattcc	540
ctctctgaca	tggccagcaa	gatoocggag	gaoccatctt	tcattctcag	gaagaaaggag	600
ggggagaaaa	aacgagaggt	atataataat	ccagtgaaaa	tgaaagaaat	caagaatttg	660
ttgcgaatga	tgtctggaaaa	aaaggagag	aagaaaaaga	aggagagaa	aaagaagcac	720
aagaaacata	agcacagaag	ctcgagtagt	gatcgttcca	gcagcgagga	tgagccaggt	780
gcagaggagat	caacagaagaa	gatggcaaat	tcctccctgt	ttttgtccaa	agtcctctgga	840
tatggtctac	aggtccggaa	ctctgacogt	aaocagggtc	ttcagggctc	cttgacagca	900
gagcaaaaga	gagggcaatg	gatgaagaac	cattccagat	ccagaagctc	ctcccactca	960
cccccaagac	atgccagcaa	gaagagcacc	aggggaagcag	ggtccoggga	caggaggtct	1020
cgatccctgg	cgagaaggtc	acogtcccca	agaccagca	aattgcacaa	ctctaaagtg	1080
aaacagagag	agacagggcca	aactaggagc	ccatccacta	aaaaagaggt	ctaccaaaag	1140
cgacatgctc	ccggatcacac	cagaaaaact	ctcgacaggg	aattagagcg	aaaaacggcaa	1200
gagatgatgg	aaaaocgcaa	atggaggggg	gaggagagac	tgaacatctc	caagaggcat	1260
gctaaggatg	aggaacoggga	gcagaggcta	gagaagctgg	actccoggga	tgggaagtct	1320

	atocacogca	tgaagctgga	gagtgcaatc	acttctccoc	tggaggatog	ggtgaagogg	1380
	aaatatctact	ctttacacag	aactctoggt	gctctggaga	agaaactttat	gaaagatga	1440
5	aaactgtoc	ctctcttatt	gggtttctogt	cattttccag	ggaagctgct	gaacctttaa	1500
	ttctctttat	aagagctcaa	tgaactctct	tcacagatgt	caaaacacaa	gtgttcaaa	1560
	tgacctgtct	tcattgagtc	ctgaaacacg	tcactctctt	tgagagctag	tgtgactgtc	1620
	tttgggtggac	actcagtaac	tttgggtttt	gaactcttaa	oggggtggga	tggtgaccta	1680
	toggtggggag	ctgtgtgtgc	actctgggaag	gctgttctct	gggtgtgtga	tgtttatcat	1740
	gcoactctct	tcttcaactgt	gcoaacacag	ctatttcaat	gctctagagt	acacacagac	1800
10	cttcagaaac	ctctctgtgt	tcacacagat	agattgtgtg	tactgagaca	aatgaacgtt	1860
	tacttgattt	agaaagataat	gtgacagaa	gatgtcaggt	taggtcaaa	ccaaagggat	1920
	gaacaaatct	ggaanaatcaa	acaaacacaa	aagcctcaaa	tgaaactgtta	actattttgat	1980
	ctttggatgt	aaaattgttaa	tgcgtatatg	tacaaatgta	caattttttac	atgttttttaa	2040
	aaaaggtttag	ctttgttgaaa	atacctgtgt	tggtcaatga	ctttactggg	taatgagaacc	2100
	acattgaacc	ttgatggcaa	gtaatacaat	aaggcaggcc	agctgtgttt	tctctctgaa	2160
15	tctgctgtgt	ttaggaggag	ctgtgggttta	tcgacagagt	ctggagatc	tattcttttc	2220
	cactgctgtc	agtctccaat	gtaggcagtg	taaaaggtata	gtaaaatgat	tttaggagtc	2280
	agaaccaa	tgccaatatg	ctccatggct	ccaaaggaa	aataaatg	aagtttttaa	2340
	aaaaaaaa	aaaaaa					2356
20	<210>	73					
	<211>	1646					
	<212>	DNA					
25	<213>	Homo sapiens					
	<400>	73					
30	gtggaatgtc	atcagttaa	gctattttca	ttttttttgt	ggatcttcag	tggtctcagg	60
	ccatctggat	gtatacatgc	aggtcacagg	gaatatgat	gcttagcttg	gggtcacagg	120
	ctcgacacct	caggctgcaa	aatgtggaag	atttaaatca	ttgaacaaat	acactctccc	180
	caaaaactga	aattggcttc	gtttctctgag	ttgtctcagg	ogcaatgttc	agcgattttg	240
	aggaaatcac	aagaattgtta	gttaaaggaga	tgtagctctg	aggggataat	attgcoogtta	300
	gaagccttgt	tgatgctgat	agattccgct	gcttccatct	gggtggggag	aagagaactt	360
35	tottttgtag	coggcactac	acacacaggc	tcacactgat	ggacattctg	gacacacatg	420
	gggacaagtg	gttagatgaa	ctggattctg	ggctccaaag	tcaaaaggct	gagtttcaaa	480
	ttctggataa	tgtagactca	aoggagagtg	tgatagttag	attacocaaa	gaaataacaa	540
	tttcaggcag	tttccagggc	ttccacacac	agaaaatcaa	gatctoggag	aacoggatat	600
	ccacagcagta	tctggctaac	cttgaaacaa	ggaagctgaa	gagggaacta	cocttttcat	660
40	tccatcaaat	taatacagaga	gaaaacctgt	atctgtgtgac	agaaactctg	gagacgttaa	720
	aggaggaac	cctgaaacag	gaccggcaat	ataaattttg	gagccagatc	tctcagggcc	780
	atctcagcta	taaacacaa	ggccaaagg	aagtgaacat	cccccacaa	cggtctctga	840
	gctctcagtg	aaagcagctt	gtttccocca	acaaggagac	gatgagaaag	tctttgggtt	900
	ogggagattc	cagaacaatg	aggagagaat	tgaggacat	ggagagtgtc	ctcagggaac	960
	tgacagagga	gaagagaaaa	gatgtgctaa	actccctogo	taagtgtctc	ggcnaaggagg	1020
45	atatctcgga	ggatctagag	caaaagatat	ctgaggtctc	gtgtctcggg	gagctacaca	1080
	tgaggagacc	agacaagcct	ctctcaagca	goccttttaa	tgctgtctgg	gtctgtgtag	1140
	aagcgtgtgc	aaaagccatt	ctggactctc	tggtgctcct	ctgagagctc	ctggaagagc	1200
	agcagtttgt	ggctgaggcc	ctggagaagg	ggaacccctc	tctgtttga	gaacagggtg	1260
	aactctgtcat	ggagcagaac	tggtgatgag	tggtcagcag	ctctcctgat	atggactatg	1320
50	acocctgagg	acgaattctc	tgtgtgtgtg	atgtgtgtgt	ctctatctct	ctggagctgtg	1380
	ctgagggggc	tacactgtct	ttctcttaac	tacaaaagcc	gtttctccoc	ccaaagcctc	1440
	gggtttctcc	tttaacagtc	tgtctctca	gocactgoc	ctacactctc	gtcaccagtg	1500
	ggacactctt	aaaaacagca	gocaaacatt	ctttgtatgt	tcocatttgc	tcattgttaa	1560
	cattccaaac	cagcctggat	ttcatcatg	gaactctgat	taaaagtggc	aggtgtgtgca	1620
	gttataaaaa	aaaaaaaa	aaaaaa				1646
55							

<210> 74

<211> 3340

<212> DNA

<213> Homo sapiens

<400> 74

oggggogccca	gagacagagc	ogcctcagat	atcctgtctgg	atgacattgt	cottaccocat	60
tctctctctcc	taocagacga	gaattttctg	caggagctac	accagtaoct	tgttogggga	120
ggaggcatgg	agggcctcga	agggctgggg	oggaagcaag	octgtctagc	catgtctctc	180
catttcttgg	acacctacaa	ggggctgctt	caagaggaag	aggggggogg	ccacatcctc	240
aaggatctat	aootgtcaat	taagaaggac	gagtcctctt	accagggoot	cogagaggac	300
actctgaggg	tgccaccagot	ggtggagacg	gtggaactaa	agattccaga	gggagaaccag	360
ccaccocagca	agcagggtgaa	gcoactctct	ogcoactctc	gocggataga	ctootgtctg	420
cagacacoggg	tggocttccog	gggctctgat	gagatctctt	gocgtgtata	catgocctgac	480
caotctatgt	tgacccaatg	cagocgoott	tcagcatctg	tgacaggcat	tctgggctct	540
gtgaogggaga	aaacttcaata	ttcagaggag	oogogggggc	gtgaggattc	octcatcctg	600
gtagctgtgt	ootctctctg	agagaaggctc	ottctocagc	ccactgagga	ctgtgttttc	660
acogcagacg	gcaatcaacag	ccacotgttt	gocgttactc	gggacagcta	tgaggctctg	720
gtgcocctcc	cogagagagat	ccaggtctcc	oocggagaca	cagagatcca	cogagtggag	780
octtgaggacg	ttgcocaaoca	octaaotgoc	ttcactggg	agctgttccg	atgtgtgcac	840
gagctggagt	togtggaccta	ogtgttccac	ggggagocgg	gocgocggga	gaocggocaa	900
ttggagotgc	tgctgcagcg	ctgcagocag	gtcacgcact	gggtggccac	agaagtgtgt	960
ctctgcaggg	cccogggcga	gocogocgag	ctgctcaaga	agttcaatca	gatocggoc	1020
ctctgcaggg	ttgagaacagg	octgtgtct	ttctacogcg	tggtctatgg	ctgtggacaa	1080
gocgtgtgta	gocgootctg	actcaootgg	gagaagctgc	cagggaatct	caagaacttg	1140
ttctgcocaa	ttgagaacact	gacggacccc	tgacgggaac	acaaagcta	cogagaagtg	1200
atctccaaaa	tgaaagccccc	tgtgattccc	ttcgtgctcc	tgactcccaa	agaactgact	1260
ttcctgocog	aaggggatga	gaacootgtg	gatgttttgg	tgacaatoga	gaactgtgac	1320
tcagtgtgocg	aaaaagtggag	gaacatccgc	aaataccgga	gocggccccc	ttgocctggac	1380
atggaggcat	cccccactca	octgcagacc	aaggcctatg	tgocccagtt	taaggtcatc	1440
gacaaocaga	actctctctt	ogagctctcc	tacacagctg	aggcaaaacg	tcagtggag	1500
tgagggtctc	agtcagaccc	gocagatcct	tgggccactc	gcaactcaagc	acttgcacog	1560
atgtctcctac	caacatctga	catcttccgc	gtggagacac	ttcctgtctc	acogggaaaga	1620
ggtgatagga	tttaccocctg	gaoccaaag	tcgttctcat	ctgctgaagt	ccctccocca	1680
ttgtctcctc	aagocaaaaac	tacactttgc	tggtctctgt	ccoccttgag	aaaggggata	1740
gaagotctc	tcctctatgt	octcccatcg	agatctgttc	tggggatgga	gctcccaact	1800
tcctcttctga	gcaggaaaga	atgtctgctc	ccactctgtc	ttgcagagtg	ggatgtgtgg	1860
agggatgtgc	agootctctc	tcacacacct	gtccagcttc	ttcctgttca	ggcctgggac	1920
ccocaggaat	attatgtgc	ogtgtgtgtg	tgtgtgtgtg	tgtgtgtgtg	tgtgtgtgtg	1980
tgtgtctctc	ttataggagc	agggagtcat	ctgttaattg	aggggtgagt	ttgtgtgtgc	2040
tgggtagggg	tcctctgtgt	tggtgtctac	ctgtctactc	ctgcocctgg	atgggtgggg	2100
gtgctttctc	caccccacaa	ctccctgtct	agctcttgt	gtgcocctgc	atgcocaggc	2160
ttgttgagcca	aggtgtcttt	tggggcaggg	agttagcaga	gtggggaggg	gttaccocac	2220
agoccttgca	agtcocccac	tcaggootct	ggaaggtcca	gggatgggct	ctgatgagag	2280
ggtaaaagat	atcaggggaa	acacaggcct	cagctgocct	gaggacocct	ccocctgcct	2340
gcagtgggct	ccggttagagc	agtaacagga	gctagggttg	tcctgtctcc	acactcctgc	2400
tttttgggat	gtcttaactgc	taaggaggga	gttgacatcc	cccttctggc	ctactgtctc	2460
gacacccaag	acatgtgtct	ogtccctctc	tcctagactc	tcocctttgtc	ctccccaatg	2520
agctgggtgg	gggtgtgctcc	ctataagtgg	gcaaggcagcc	ccaaagtggt	gttaccocac	2580
ggcagagact	gtaaaggggc	cactggactc	tggaagggoc	tttattacoc	ttactccctc	2640
ccctctccat	ccacagctcc	aaggcctgag	gggtgcaggg	gctccctggca	ctactgtggg	2700
gaggtttctc	ggccacagact	ccocctctct	ctctggcacc	ctctttccct	tttgaagaga	2760
cagcaacacg	cgtagacaaa	gcagctgtg	ctcctgtcat	gaggggtgat	atatttttta	2820
cccaaaagctc	tggaattgtga	catttatctt	ttaaaaatca	aagagggaaa	gagocctgtg	2880

	tcattatgtga	acattgttatc	ataggtaatg	ttgtacagac	cotttttac	agtgatctgt	2940
	cttgttctctg	cagcaaaaat	octotatgga	cataggagg	gctgtgtccc	atgctctctt	3000
	gcoctgacag	tgtoctatgg	gcoctctctt	gctcctctgc	cctcctctgc	tactgtgat	3060
5	gcactgtctct	ctcctctgag	coctctgctt	cccagctctc	ctcctgaccc	ctctcaacag	3120
	ccttggaact	caaactgcga	caacccctctg	ggtctggacac	tggggaccac	tggcccaagtc	3180
	ttggctgtctg	cttaacccta	gcocttgatg	ctgcccagg	accccagcc	cctccctggt	3240
	gcoctgcagc	tttaacacag	tgaacctatg	gtatgttaca	ggcgoggtg	tcattgcaga	3300
	aaocgtctgg	tggagaagaa	gocgataaag	tctatgaate			3340
10	<210>	75					
	<211>	4005					
	<212>	DNA					
15	<213>	Homo sapiens					
	<400>	75					
20	gggcaacagt	ctgcccacct	gtggacacca	gactctggga	gctcctgggt	agcaagtggag	60
	atctctctgg	tgctagtgag	gctggttgaa	gacacagagt	aaactgcaga	ggtcacccac	120
	ccaacactgt	ccaaggtgat	gtccagccca	ctgctggcag	gaggccatgtg	tgctcagctgtg	180
	gcyctctgtg	atgagccca	gaggacccctg	caccagccac	ccagccccag	ccttgccaccc	240
	catgtgtctt	actacaccac	ggaaggtctg	ggagccagag	cctgtatggo	ccctctgccc	300
25	tgtcaatgggc	ccctctggcc	actccagcaa	gcccacagg	tggaggccaa	agccacctgc	360
	ttcctgcctg	ccctctggga	gaaggccttg	gggaccccag	aggaccttga	ctctacatt	420
	gacttctcac	tggagagcct	caatcagatg	atcctggaa	tggacccac	cttccagctg	480
	cttcccccag	ggactggggg	ctcccagctc	gagctggccc	agagccacat	gtcaatgaga	540
	aagaaggagg	aatctgaagc	cttggacata	aagtacatog	aggtgaacct	cgccagatca	600
	aggtgcacag	attggcccga	gcactgctcc	agccctctg	tcaccccctc	cttgcctccc	660
30	cctcgacgtg	gtggcctcct	ccttccacga	gactctcccc	gagagacacg	aagcagcagt	720
	gagagcctca	tctctctctg	gaacacgggc	agggggccac	agccgcctct	gccccctca	780
	gagggctctc	ccctctgacc	cccaaatctc	cccagcactc	caatcccttg	catggggagc	840
	aaaggcctga	gccccactgg	tttgggctcc	agccggatct	ctgtgctgtc	agccagccca	900
	cggtctggag	gctctggccc	acagcggggc	agccagctcc	ctctgcactc	cagcaactcc	960
35	gtgtctgatg	tcagcttatat	gttctggaa	agtcacagca	actctctctc	cagcctccac	1020
	agccttggtc	cagtgctcct	gtgtacaa	cccagtgact	tcacggctcc	cagaaacccc	1080
	accctaacca	tggggccaccc	cagaaacccc	acactctccc	cactgtgcac	agactctccc	1140
	agcatctctc	ccccaatcat	ccccaactcc	atgtgtggac	tacccaatgt	gctgtatcaac	1200
40	ggctctccag	aaocagggtc	ttctccaccc	caagcggacc	caggacaccc	caactcctgt	1260
	caactctggag	ctgctctctc	cagcaacccc	tgctcagcca	ccaggagcaa	cagccagacc	1320
	ctgtcaagtg	cccccctttac	caatctgcca	gaggtctccg	caaactctac	cagagagcaa	1380
	atgaagttcg	tgtatggacac	atctaaatac	tggtttaagc	tcataaggga	tgctcagact	1440
	gcaatcgagc	cctctgagaa	ggaggagcca	ggggcttttg	ttcccgctgc	tgctcagact	1500
45	tacagaggtc	cctctggcct	ggccctgaag	gtgcaggagg	tcataaggga	tgctcagact	1560
	cgacacagtg	agacagccaa	tgacactctc	cgacactctc	ggagcctctc	tgctcagact	1620
	ggagtgcaat	tcaaaaggagc	agatgaggag	ccctactctg	tcaccaatcc	cagagagcaa	1680
	tgccacagct	ccatcaatggc	cctggccctg	ccctgcaaac	ccagcctctg	cagagagcaa	1740
	ctgggaggtg	cagatggggc	ctcggaactc	acagacagcc	tgagagaccc	cagagagcaa	1800
	ctctggcgtg	cccaacccct	gtacctgagc	tcagtgagcg	acatctctcc	cagagagcaa	1860
	ctggccctgc	agaaagccat	ctccacccac	tttgaggagg	acatctctcc	cagagagcaa	1920
50	gtgtgctcat	tcgaagtcaac	agagcagggc	atcaactctg	ctgatgtcca	gaggaaggtg	1980
	tttttctggc	gcaattaccc	actcaacccc	ctccgctctc	gtgtgtatga	ccttgagcaa	2040
	cgggaagtg	aggaactactg	caaacctccc	tggtctcttg	gggtttgtgg	caagagagcc	2100
	acagagcctc	aggaagaggt	atgcccaccc	tttgaggagg	atgaactggt	ccagcagacc	2160
	tcggcaggtc	tggcctgggt	gaactgctct	ctgcaggagc	cagaaaggat	gtggggaga	2220
55	gactgcctgt	gcaactaccc	aacaccccca	ggggctcctc	aaggagcccc	cctccacccc	2280

5 ctgaatgggt gtggcttgtg gccatattga cagaccatc tatgggacta gggggattgg 240(
 catcaagttg acacccottga aactgctatg gccttcagca gtcaccatca tccagacccc 246(
 cogggocctca gtttctctcaa tcatagaaga agaccaatag acaagatcag ctgttcttag 252(
 atgtctgtgg gatatttgaa atgctctccc atgattctga agcatgcaca cctctgaaga 258(
 ccctctgcat aaaaataact ccaaggaccc ctgaccccca tcgacctggg cctgtccacc 264(
 acacagctct gagcaagaga cctgcagccc ctgtttctgt gcagacagca ggtgocctgg 270(
 gtgacccac ggggctctctg gcttgacgtc ggtgaagtgc aagaactgac tacaaaaag 276(
 gaatggatag actctatttc cttccatctc tgttctctgt ttccttttcc caactttctg 282(
 gtggcttttt gggttccccc agccaggatg ctgcaggcca agctgggtgt ggtatttagg 288(
 cagctcagc aggggggaact tgttcccatg gtccagaggag aocccagctg cctgaccccc 294(
 ctgcagatg agtatcaccc catcttttct ttccacttgg ttttattttt tatttttttt 300(
 gagacagagt ctcactgtca cccaggctga actgcagctg tgtgatctag gtcactgtca 306(
 acctccacct cccaggttca agcaattatc ctgcctcagg ctcccgagta gctgggatta 312(
 caggcatgtg caactcacc agctaatttt gtatttttag tagagacagg gtttaccat 318(
 gttgccagg ctggtcttga actcctgaac gcaggtaatc caactgcttc ggcttcccaa 324(
 agtgtcggga ttacaggcgc aagccaccac gccagctctc tttccattcc ttgataggcg 330(
 agtattccaa agctgggtatc gtagctgccc taagtgtgca tattaggcgg ogggggcaga 336(
 gataaggggc atctctctgt gattctgccc cagctcctgt cttgctgagc cctcccccac 342(
 ccacagctcc aacacacaca cacacacaca cacacacaca cacacacaca cacacacaca 348(
 cacgcccctc tactctctatg tggcttccaa cagcctcaca gccacacagg ggaagcagag 354(
 agtcaagaat gcaaagaggc cgcttcccta agaggcttgg aggagctggg ctctatccca 360(
 caccccaccc caccccaccc caccccaccc tccagaagct ggaacattt ctcccgagg 366(
 cctgagttcc taaggaaacc accctacagg ggtggaaggg agggtcaggg aagaaaccca 372(
 ctctgtctct aogaggagca agtgcctgcc ccctccacgc agccagccct gccaaagtgt 378(
 cattatcttt ggcccaaggct gggctcagag gttatgattt cagccctggg ctgcagagg 384(
 aggctgagat cagcccccac agccagtggt cgagcactgc cccgccgcca aagctgtcag 390(
 aatgtgagat gaggttctca aggctcacagg cccagtcacc agcctggggg ctggcagagg 396(
 ccccatata ctctgtatca gtcctctatc tgaaaaataa aatgt 4005

<210> 76

<211> 1093

<212> PRT

<213> Homo sapiens

<400> 76

Met Lys Glu Met Val Gly Gly Cys Cys Val Cys Ser Asp Glu Arg Gly
 1 5 10 15
 Trp Ala Glu Asn Pro Leu Val Tyr Cys Asp Gly His Ala Cys Ser Val
 20 25 30
 Ala Val His Gln Ala Cys Tyr Gly Ile Val Gln Val Pro Thr Gly Pro
 35 40 45
 Trp Phe Cys Arg Lys Cys Glu Ser Gln Glu Arg Ala Ala Arg Val Arg
 50 55 60
 Cys Glu Leu Cys Pro His Lys Asp Gly Ala Leu Lys Arg Thr Asp Asn
 65 70 75 80
 Gly Gly Trp Ala His Val Val Cys Ala Leu Tyr Ile Pro Glu Val Gln
 85 90 95
 Phe Ala Asn Val Leu Thr Met Glu Pro Ile Val Leu Gln Tyr Val Pro
 100 105 110
 His Asp Arg Phe Asn Lys Thr Cys Tyr Ile Cys Glu Glu Thr Gly Arg
 115 120 125
 Glu Ser Lys Ala Ala Ser Gly Ala Cys Met Thr Cys Asn Arg His Gly
 130 135 140

Cys Arg Gln Ala Phe His Val Thr Cys Ala Gln Met Ala Gly Leu Leu
 145 150 155 160
 Cys Glu Glu Glu Val Leu Glu Val Asp Asn Val Lys Tyr Cys Gly Tyr
 165 170 175
 Cys Lys Tyr His Phe Ser Lys Met Lys Thr Ser Arg His Ser Ser Gly
 180 185 190
 Gly Gly Gly Gly Ala Gly Gly Gly Gly Ser Met Gly Gly Gly
 195 200 205
 Gly Ser Gly Phe Ile Ser Gly Arg Arg Ser Arg Ser Ala Ser Pro Ser
 210 215 220
 Thr Gln Gln Glu Lys His Pro Thr His His Glu Arg Gly Gln Lys Lys
 225 230 235 240
 Ser Arg Lys Asp Lys Glu Arg Leu Lys Gln Lys His Lys Lys Arg Pro
 245 250 255
 Glu Ser Pro Pro Ser Ile Leu Thr Pro Pro Val Val Pro Thr Ala Asp
 260 265 270
 Lys Val Ser Ser Ser Ala Ser Ser Ser Ser His His Glu Ala Ser Thr
 275 280 285
 Gln Glu Thr Ser Glu Ser Ser Arg Glu Ser Lys Gly Lys Lys Ser Ser
 290 295 300
 Ser His Ser Leu Ser His Lys Gly Lys Lys Leu Ser Ser Gly Lys Gly
 305 310 315 320
 Val Ser Ser Phe Thr Ser Ala Ser Ser Ser Ser Ser Ser Ser Ser
 325 330 335
 Ser Ser Gly Gly Pro Phe Gln Pro Ala Val Ser Ser Leu Gln Ser Ser
 340 345 350
 Pro Asp Phe Ser Ala Phe Pro Lys Leu Glu Gln Pro Glu Glu Asp Lys
 355 360 365
 Tyr Ser Lys Pro Thr Ala Pro Ala Pro Ser Ala Pro Pro Ser Pro Ser
 370 375 380
 Ala Pro Glu Pro Pro Lys Ala Asp Leu Phe Glu Gln Lys Val Val Phe
 385 390 395 400
 Ser Gly Phe Gly Pro Ile Met Arg Phe Ser Thr Thr Thr Ser Ser Ser
 405 410 415
 Gly Arg Ala Arg Ala Pro Ser Pro Gly Asp Tyr Lys Ser Pro His Val
 420 425 430
 Thr Gly Ser Gly Ala Ser Ala Gly Thr His Lys Arg Met Pro Ala Leu
 435 440 445
 Ser Ala Thr Pro Val Pro Ala Asp Glu Thr Pro Glu Thr Gly Leu Lys
 450 455 460
 Glu Lys Lys His Lys Ala Ser Lys Arg Ser Arg His Gly Pro Gly Arg
 465 470 475 480
 Pro Lys Gly Ser Arg Asn Lys Glu Gly Thr Gly Gly Pro Ala Ala Pro
 485 490 495
 Ser Leu Pro Ser Ala Gln Leu Ala Gly Phe Thr Ala Thr Ala Ala Ser
 500 505 510
 Pro Phe Ser Gly Gly Ser Leu Val Ser Ser Gly Leu Gly Gly Leu Ser
 515 520 525
 Ser Arg Thr Phe Gly Pro Ser Gly Ser Leu Pro Ser Leu Ser Leu Glu
 530 535 540
 Ser Pro Leu Leu Gly Ala Gly Ile Tyr Thr Ser Asn Lys Asp Pro Ile
 545 550 555 560
 Ser His Ser Gly Gly Met Leu Arg Ala Val Cys Ser Thr Pro Leu Ser
 565 570 575
 Ser Ser Leu Leu Gly Pro Pro Gly Thr Ser Ala Leu Pro Arg Leu Ser
 580 585 590
 Arg Ser Pro Phe Thr Ser Thr Leu Pro Ser Ser Ser Ala Ser Ile Ser
 595 600 605

Thr Thr Gln Val Phe Ser Leu Ala Gly Ser Thr Phe Ser Leu Pro Ser
 610 615 620
 Thr His Ile Phe Gly Thr Pro Met Gly Ala Val Asn Pro Leu Leu Ser
 625 630 635 640
 Gln Ala Glu Ser Ser His Thr Glu Pro Asp Leu Glu Asp Cys Ser Phe
 645 650 655
 Arg Cys Arg Gly Thr Ser Pro Gln Glu Ser Leu Ser Ser Met Ser Pro
 660 665 670
 Ile Ser Ser Leu Pro Ala Leu Phe Asp Gln Thr Ala Ser Ala Pro Cys
 675 680 685
 Gly Gly Gly Gln Leu Asp Pro Ala Ala Pro Gly Thr Thr Asn Met Glu
 690 695 700
 Gln Leu Leu Glu Lys Gln Gln Asp Gly Glu Ala Gly Val Asn Ile Val
 705 710 715 720
 Glu Met Leu Lys Ala Leu His Ala Leu Gln Lys Glu Asn Gln Arg Leu
 725 730 735
 Gln Glu Gln Ile Leu Ser Leu Thr Ala Lys Lys Glu Arg Leu Gln Ile
 740 745 750
 Leu Asn Val Gln Leu Ser Val Pro Phe Pro Ala Leu Pro Ala Ala Leu
 755 760 765
 Pro Ala Ala Asn Gly Pro Val Pro Gly Pro Tyr Gly Leu Pro Pro Gln
 770 775 780
 Ala Gly Ser Ser Asp Ser Leu Ser Thr Ser Lys Ser Pro Pro Gly Lys
 785 790 795 800
 Ser Ser Leu Gly Leu Asp Asn Ser Leu Ser Thr Ser Ser Glu Asp Pro
 805 810 815
 His Ser Gly Cys Pro Ser Arg Ser Ser Ser Ser Leu Ser Phe His Ser
 820 825 830
 Thr Pro Pro Leu Pro Leu Leu Gln Gln Ser Pro Ala Thr Leu Pro
 835 840 845
 Leu Ala Leu Pro Gly Ala Pro Ala Pro Leu Pro Pro Gln Pro Gln Asn
 850 855 860
 Gly Leu Gly Arg Ala Pro Gly Ala Ala Gly Leu Gly Ala Met Pro Met
 865 870 875 880
 Ala Glu Gly Leu Leu Gly Gly Leu Ala Gly Ser Gly Gly Leu Pro Leu
 885 890 895
 Asn Gly Leu Leu Gly Gly Leu Asn Gly Ala Ala Ala Pro Asn Pro Ala
 900 905 910
 Ser Leu Ser Gln Ala Gly Gly Ala Pro Thr Leu Gln Leu Pro Gly Cys
 915 920 925
 Leu Asn Ser Leu Thr Glu Gln Gln Arg His Leu Leu Gln Gln Gln Glu
 930 935 940
 Gln Gln Leu Gln Gln Leu Gln Gln Leu Leu Ala Ser Pro Gln Leu Thr
 945 950 955 960
 Pro Glu His Gln Thr Val Val Tyr Gln Met Ile Gln Gln Ile Gln Gln
 965 970 975
 Lys Arg Glu Leu Gln Arg Leu Gln Met Ala Gly Gly Ser Gln Leu Pro
 980 985 990
 Met Ala Ser Leu Leu Ala Gly Ser Ser Thr Pro Leu Leu Ser Ala Gly
 995 1000 1005
 Thr Pro Gly Leu Leu Pro Thr Ala Ser Ala Pro Pro Leu Leu Pro
 1010 1015 1020
 Ala Gly Ala Leu Val Ala Pro Ser Leu Gly Asn Asn Thr Ser Leu
 1025 1030 1035
 Met Ala Ala Ala Ala Ala Ala Ala Ala Val Ala Ala Ala Gly Gly
 1040 1045 1050
 Pro Pro Val Leu Thr Ala Gln Thr Asn Pro Phe Leu Ser Leu Ser
 1055 1060 1065

Gly Ala Glu Gly Ser Gly Gly Gly Pro Lys Gly Gly Thr Ala Asp
 1070 1075 1080
 Lys Gly Ala Ser Ala Asn Gln Glu Lys Gly
 1085 1090

<210> 77

<211> 344

<212> PRT

<213> Homo sapiens

<400> 77

Met His Arg Thr Thr Arg Ile Lys Ile Thr Glu Leu Asn Pro His Leu
 1 5 10 15
 Met Cys Ala Leu Cys Gly Gly Tyr Phe Ile Asp Ala Thr Thr Ile Val
 20 25 30
 Glu Cys Leu His Ser Phe Cys Lys Thr Cys Ile Val Arg Tyr Leu Glu
 35 40 45
 Thr Asn Lys Tyr Cys Pro Met Cys Asp Val Gln Val His Lys Thr Arg
 50 55 60
 Pro Leu Leu Ser Ile Arg Ser Asp Lys Thr Leu Gln Asp Ile Val Tyr
 65 70 75 80
 Lys Leu Val Pro Gly Leu Phe Lys Asp Glu Met Lys Arg Arg Arg Asp
 85 90 95
 Phe Tyr Ala Ala Tyr Pro Leu Thr Glu Val Pro Asn Gly Ser Asn Glu
 100 105 110
 Asp Arg Gly Glu Val Leu Glu Gln Glu Lys Gly Ala Leu Ser Asp Asp
 115 120 125
 Glu Ile Val Ser Leu Ser Ile Glu Phe Tyr Glu Gly Ala Arg Asp Arg
 130 135 140
 Asp Glu Lys Lys Gly Pro Leu Glu Asn Gly Asp Gly Asp Lys Glu Lys
 145 150 155 160
 Thr Gly Val Arg Phe Leu Arg Cys Pro Ala Ala Met Thr Val Met His
 165 170 175
 Leu Ala Lys Phe Leu Arg Asn Lys Met Asp Val Pro Ser Lys Tyr Lys
 180 185 190
 Val Glu Val Leu Tyr Glu Asp Glu Pro Leu Lys Glu Tyr Thr Leu
 195 200 205
 Met Asp Ile Ala Tyr Ile Tyr Pro Trp Arg Arg Asn Gly Pro Leu Pro
 210 215 220
 Leu Lys Tyr Arg Val Gln Pro Ala Cys Lys Arg Leu Thr Leu Ala Thr
 225 230 235 240
 Val Pro Thr Pro Ser Glu Gly Thr Asn Thr Ser Gly Ala Ser Glu Cys
 245 250 255
 Glu Ser Val Ser Asp Lys Ala Pro Ser Pro Ala Thr Leu Pro Ala Thr
 260 265 270
 Ser Ser Ser Leu Pro Ser Pro Ala Thr Pro Ser His Gly Ser Pro Ser
 275 280 285
 Ser His Gly Pro Pro Ala Thr His Pro Thr Ser Pro Thr Pro Pro Ser
 290 295 300
 Thr Ala Ser Gly Ala Thr Thr Ala Ala Asn Gly Gly Ser Leu Asn Cys
 305 310 315 320
 Leu Gln Thr Pro Ser Ser Thr Ser Arg Gly Arg Lys Met Thr Val Asn
 325 330 335

Gly Ala Pro Val Pro Pro Leu Thr
340

<210> 78

<211> 416

<212> PRT

<213> Homo sapiens

<400> 78

Met Ser Ser Asn Cys Thr Ser Thr Thr Ala Val Ala Val Ala Pro Leu
1 5 10 15
Ser Ala Ser Lys Thr Lys Thr Lys Lys His Phe Val Cys Gln Lys
20 25 30
Val Lys Leu Phe Arg Ala Ser Glu Pro Ile Leu Ser Val Leu Met Trp
35 40 45
Gly Val Asn His Thr Ile Asn Glu Leu Ser Asn Val Pro Val Pro Val
50 55 60
Met Leu Met Pro Asp Asp Phe Lys Ala Tyr Ser Lys Ile Lys Val Asp
65 70 75 80
Asn His Leu Phe Asn Lys Glu Asn Leu Pro Ser Arg Phe Lys Phe Lys
85 90 95
Glu Tyr Cys Pro Met Val Phe Arg Asn Leu Arg Glu Arg Phe Gly Ile
100 105 110
Asp Asp Gln Asp Tyr Gln Asn Ser Val Thr Arg Ser Ala Pro Ile Asn
115 120 125
Ser Asp Ser Gln Gly Arg Cys Gly Thr Arg Phe Leu Thr Thr Tyr Asp
130 135 140
Arg Arg Phe Val Ile Lys Thr Val Ser Ser Glu Asp Val Ala Glu Met
145 150 155 160
His Asn Ile Leu Lys Lys Tyr His Gln Phe Ile Val Glu Cys His Gly
165 170 175
Asn Thr Leu Leu Pro Gln Phe Leu Gly Met Tyr Arg Leu Thr Val Asp
180 185 190
Gly Val Glu Thr Tyr Met Val Val Thr Arg Asn Val Phe Ser His Arg
195 200 205
Leu Thr Val His Arg Lys Tyr Asp Leu Lys Gly Ser Thr Val Ala Arg
210 215 220
Glu Ala Ser Asp Lys Glu Lys Ala Lys Asp Leu Pro Thr Phe Lys Asp
225 230 235 240
Asn Asp Phe Leu Asn Glu Gly Gln Lys Leu His Val Gly Glu Glu Ser
245 250 255
Lys Lys Asn Phe Leu Glu Lys Leu Lys Arg Asp Val Glu Phe Leu Ala
260 265 270
Gln Leu Lys Ile Met Asp Tyr Ser Leu Leu Val Gly Ile His Asp Val
275 280 285
Asp Arg Ala Glu Gln Glu Glu Met Glu Val Glu Glu Arg Ala Glu Asp
290 295 300
Glu Glu Cys Glu Asn Asp Gly Val Gly Gly Asn Leu Leu Cys Ser Tyr
305 310 315 320
Gly Thr Pro Pro Asp Ser Pro Gly Asn Leu Leu Ser Phe Pro Arg Phe
325 330 335
Phe Gly Pro Gly Glu Phe Asp Pro Ser Val Asp Val Tyr Ala Met Lys
340 345 350

Ser His Glu Ser Ser Pro Lys Lys Glu Val Tyr Phe Met Ala Ile Ile
 355 360 365
 Asp Ile Leu Thr Pro Tyr Asp Thr Lys Lys Lys Ala Ala His Ala Ala
 370 375 380
 Lys Thr Val Lys His Gly Ala Gly Ala Glu Ile Ser Thr Val Asn Pro
 385 390 395 400
 Glu Gln Tyr Ser Lys Arg Phe Asn Glu Phe Met Ser Asn Ile Leu Thr
 405 410 415

<210> 79

<211> 500

<212> PRT

<213> Homo sapiens

<400> 79

Met Arg Gly Glu Leu Trp Leu Leu Val Leu Val Leu Arg Glu Ala Ala
 1 5 10 15
 Arg Ala Leu Ser Pro Gln Pro Gly Ala Gly His Asp Glu Gly Pro Gly
 20 25 30
 Ser Gly Trp Ala Ala Lys Gly Thr Val Arg Gly Trp Asn Arg Arg Ala
 35 40 45
 Arg Glu Ser Pro Gly His Val Ser Glu Pro Asp Arg Thr Gln Leu Ser
 50 55 60
 Gln Asp Leu Gly Gly Gly Thr Leu Ala Met Asp Thr Leu Pro Asp Asn
 65 70 75 80
 Arg Thr Arg Val Val Glu Asp Asn His Ser Tyr Tyr Val Ser Arg Leu
 85 90 95
 Tyr Gly Pro Ser Glu Pro His Ser Arg Glu Leu Trp Val Asp Val Ala
 100 105 110
 Glu Ala Asn Arg Ser Gln Val Lys Ile His Thr Ile Leu Ser Asn Thr
 115 120 125
 His Arg Gln Ala Ser Arg Val Val Leu Ser Phe Asp Phe Pro Phe Tyr
 130 135 140
 Gly His Pro Leu Arg Gln Ile Thr Ile Ala Thr Gly Gly Phe Ile Phe
 145 150 155 160
 Met Gly Asp Val Ile His Arg Met Leu Thr Ala Thr Gln Tyr Val Ala
 165 170 175
 Pro Leu Met Ala Asn Phe Asn Pro Gly Tyr Ser Asp Asn Ser Thr Val
 180 185 190
 Val Tyr Phe Asp Asn Gly Thr Val Phe Val Val Gln Trp Asp His Val
 195 200 205
 Tyr Leu Gln Gly Trp Glu Asp Lys Gly Ser Phe Thr Phe Gln Ala Ala
 210 215 220
 Leu His His Asp Gly Arg Ile Val Phe Ala Tyr Lys Glu Ile Pro Met
 225 230 235 240
 Ser Val Pro Glu Ile Ser Ser Ser Gln His Pro Val Lys Thr Gly Leu
 245 250 255
 Ser Asp Ala Phe Met Ile Leu Asn Pro Ser Pro Asp Val Pro Glu Ser
 260 265 270
 Arg Arg Arg Ser Ile Phe Glu Tyr His Arg Ile Glu Leu Asp Pro Ser
 275 280 285
 Lys Val Thr Ser Met Ser Ala Val Glu Phe Thr Pro Leu Pro Thr Cys
 290 295 300

Leu Gln His Arg Ser Cys Asp Ala Cys Met Ser Ser Asp Leu Thr Phe
 305 310 315 320
 Asn Cys Ser Trp Cys His Val Leu Gln Arg Cys Ser Ser Gly Phe Asp
 325 330 335
 Arg Tyr Arg Gln Glu Trp Met Asp Tyr Gly Cys Ala Gln Glu Ala Glu
 340 345 350
 Gly Arg Met Cys Glu Asp Phe Gln Asp Glu Asp His Asp Ser Ala Ser
 355 360 365
 Pro Asp Thr Ser Phe Ser Pro Tyr Asp Gly Asp Leu Thr Thr Thr Ser
 370 375 380
 Ser Ser Leu Phe Ile Asp Ser Leu Thr Thr Glu Asp Asp Thr Lys Leu
 385 390 395 400
 Asn Pro Tyr Ala Gly Gly Asp Gly Leu Gln Asn Asn Leu Ser Pro Lys
 405 410 415
 Thr Lys Gly Thr Pro Val His Leu Gly Thr Ile Val Gly Ile Val Leu
 420 425 430
 Ala Val Leu Leu Val Ala Ala Ile Ile Leu Ala Gly Ile Tyr Ile Asn
 435 440 445
 Gly His Pro Thr Ser Asn Ala Ala Leu Phe Phe Ile Glu Arg Arg Pro
 450 455 460
 His His Trp Pro Ala Met Lys Phe Arg Ser His Pro Asp His Ser Thr
 465 470 475 480
 Tyr Ala Glu Val Glu Pro Ser Gly His Glu Lys Glu Gly Phe Met Glu
 485 490 495
 Ala Glu Gln Cys
 500

<210> 80

<211> 509

<212> PRT

<213> Homo sapiens

<400> 80

Met Glu Asp Ile Gln Thr Asn Ala Glu Leu Lys Ser Thr Gln Glu Gln
 1 5 10 15
 Ser Val Pro Ala Glu Ser Ala Ala Val Leu Asn Asp Tyr Ser Leu Thr
 20 25 30
 Lys Ser His Glu Met Glu Asn Val Asp Ser Gly Glu Gly Pro Ala Asn
 35 40 45
 Glu Asp Glu Asp Ile Gly Asp Asp Ser Met Lys Val Lys Asp Glu Tyr
 50 55 60
 Ser Glu Arg Asp Glu Asn Val Leu Lys Ser Glu Pro Met Gly Asn Ala
 65 70 75 80
 Glu Glu Pro Glu Ile Pro Tyr Ser Tyr Ser Arg Glu Tyr Asn Glu Tyr
 85 90 95
 Glu Asn Ile Lys Leu Glu Arg His Val Val Ser Phe Asp Ser Ser Arg
 100 105 110
 Pro Thr Ser Gly Lys Met Asn Cys Asp Val Cys Gly Leu Ser Cys Ile
 115 120 125
 Ser Phe Asn Val Leu Met Val His Lys Arg Ser His Thr Gly Glu Arg
 130 135 140
 Pro Phe Gln Cys Asn Gln Cys Gly Ala Ser Phe Thr Gln Lys Gly Asn
 145 150 155 160

Leu Leu Arg His Ile Lys Leu His Thr Gly Glu Lys Pro Phe Lys Cys
 165 170
 His Leu Cys Asn Tyr Ala Cys Gln Arg Arg Asp Ala Leu Thr Gly His
 180 185
 Leu Arg Thr His Ser Val Glu Lys Pro Tyr Lys Cys Glu Phe Cys Gly
 195 200
 Arg Ser Tyr Lys Gln Arg Ser Ser Leu Glu Glu His Lys Glu Arg Cys
 210 215
 Arg Thr Phe Leu Gln Ser Thr Asp Pro Gly Asp Thr Ala Ser Ala Glu
 225 230
 Ala Arg His Ile Lys Ala Glu Met Gly Ser Glu Arg Ala Leu Val Leu
 245 250
 Asp Arg Leu Ala Ser Asn Val Ala Lys Arg Lys Ser Ser Met Pro Gln
 260 265
 Lys Phe Ile Gly Glu Lys Arg His Cys Phe Asp Val Asn Tyr Asn Ser
 275 280
 Ser Tyr Met Tyr Glu Lys Glu Ser Glu Leu Ile Gln Thr Arg Met Met
 290 295
 Asp Gln Ala Ile Asn Asn Ala Ile Ser Tyr Leu Gly Ala Glu Ala Leu
 305 310
 Cys Pro Leu Val Gln Thr Pro Pro Ala Pro Thr Ser Glu Met Val Pro
 325 330
 Val Ile Ser Ser Met Tyr Pro Ile Ala Leu Thr Arg Ala Glu Met Ser
 340 345
 Asn Gly Ala Pro Gln Glu Leu Glu Arg Lys Ser Ile Leu Leu Pro Glu
 355 360
 Lys Ser Val Pro Ser Glu Arg Gly Leu Ser Pro Asn Asn Ser Gly His
 370 375
 Asp Ser Thr Asp Thr Asp Ser Asn His Glu Arg Gln Asn His Ile
 385 390
 Tyr Gln Gln Asn His Met Val Leu Ser Arg Ala Arg Asn Gly Met Pro
 405 410
 Leu Leu Lys Glu Val Pro Arg Ser Tyr Glu Leu Leu Lys Pro Pro
 420 425
 Ile Cys Pro Arg Asp Ser Val Lys Val Ile Asp Lys Glu Gly Glu Val
 435 440
 Met Asp Val Tyr Arg Cys Asp His Cys Arg Val Leu Phe Leu Asp Tyr
 450 455
 Val Met Phe Thr Ile His Met Gly Cys His Gly Phe Arg Asp Pro Phe
 465 470
 Glu Cys Asn Met Cys Gly Asp Arg Ser His Asp Arg Tyr Glu Phe Ser
 485 490
 Ser His Ile Ala Arg Gly Glu His Arg Ser Leu Leu Lys
 500 505

<210> 81

<211> 440

<212> PRT

<213> Homo sapiens

<400> 81
 Met Pro Ile Pro Pro Pro Pro Pro Pro Gly Pro Pro Pro Pro
 1 5 10 15
 Pro Thr Phe His Gln Ala Asn Thr Glu Gln Pro Lys Leu Ser Arg Asp
 20 25 30
 Glu Gln Arg Gly Arg Gly Ala Leu Leu Gln Asp Ile Cys Lys Gly Thr
 35 40 45
 Lys Leu Lys Lys Val Thr Asn Ile Asn Asp Arg Ser Ala Pro Ile Leu
 50 55 60
 Glu Lys Pro Lys Gly Ser Ser Gly Gly Tyr Gly Ser Gly Gly Ala Ala
 65 70 75 80
 Leu Gln Pro Lys Gly Gly Leu Phe Gln Gly Gly Val Leu Lys Leu Arg
 85 90 95
 Pro Val Gly Ala Lys Asp Gly Ser Glu Asn Leu Ala Gly Lys Pro Ala
 100 105 110
 Leu Gln Ile Pro Ser Ser Arg Ala Ala Ala Pro Arg Pro Pro Val Ser
 115 120 125
 Ala Ala Ser Gly Arg Pro Gln Asp Asp Thr Asp Ser Ser Arg Ala Ser
 130 135 140
 Leu Pro Glu Leu Pro Arg Met Gln Arg Pro Ser Leu Pro Asp Leu Ser
 145 150 155 160
 Arg Pro Asn Thr Thr Ser Ser Thr Gly Met Lys His Ser Ser Ser Ala
 165 170 175
 Pro Pro Pro Pro Pro Gly Arg Arg Ala Asn Ala Pro Pro Thr Pro
 180 185 190
 Leu Pro Met His Ser Ser Lys Ala Pro Ala Tyr Asn Arg Glu Lys Pro
 195 200 205
 Leu Pro Pro Thr Pro Gly Gln Arg Leu His Pro Gly Arg Glu Gly Pro
 210 215 220
 Pro Ala Pro Pro Pro Val Lys Pro Pro Pro Ser Pro Val Asn Ile Arg
 225 230 235 240
 Thr Gly Pro Ser Gly Gln Ser Leu Ala Pro Pro Pro Pro Tyr Arg
 245 250 255
 Gln Pro Pro Gly Val Pro Asn Gly Pro Ser Ser Pro Thr Asn Glu Ser
 260 265 270
 Ala Pro Glu Leu Pro Gln Arg His Asn Ser Leu His Arg Lys Thr Pro
 275 280 285
 Gly Pro Val Arg Gly Leu Ala Pro Pro Pro Thr Ser Ala Ser Pro
 290 295 300
 Ser Leu Leu Ser Asn Arg Pro Pro Pro Ala Arg Asp Pro Pro Ser
 305 310 315 320
 Arg Gly Ala Ala Pro Pro Pro Pro Pro Val Ile Arg Asn Gly Ala
 325 330 335
 Arg Asp Ala Pro Pro Pro Pro Pro Tyr Arg Met His Gly Ser Glu
 340 345 350
 Pro Pro Ser Arg Gly Lys Pro Pro Pro Pro Ser Arg Thr Pro Ala
 355 360 365
 Gly Pro Pro Pro Pro Pro Pro Pro Leu Arg Asn Gly His Arg Asp
 370 375 380
 Ser Ile Thr Thr Val Arg Ser Phe Leu Asp Asp Phe Glu Ser Lys Tyr
 385 390 395 400
 Ser Phe His Pro Val Glu Asp Phe Pro Ala Pro Glu Glu Tyr Lys His
 405 410 415
 Phe Gln Arg Ile Tyr Pro Ser Lys Thr Asn Arg Ala Ala Arg Gly Ala
 420 425 430
 Pro Pro Leu Pro Pro Ile Leu Arg
 435 440

<210> 82

<211> 205

<212> PRT

<213> Homo sapiens

<400> 82

```

Met Ser Ile Met Ser Tyr Asn Gly Gly Ala Val Met Ala Met Lys Gly
1      5      10      15
Lys Asn Cys Val Ala Ile Ala Ala Asp Arg Arg Phe Gly Ile Gln Ala
15     20     25
Gln Met Val Thr Thr Asp Phe Gln Lys Ile Phe Pro Met Gly Asp Arg
35     40     45
Leu Tyr Ile Gly Leu Ala Gly Leu Ala Thr Asp Val Gln Thr Val Ala
50     55     60
Gln Arg Leu Lys Phe Arg Leu Asn Leu Tyr Glu Leu Lys Glu Gly Arg
20     65     70     75     80
Gln Ile Lys Pro Tyr Thr Leu Met Ser Met Val Ala Asn Leu Leu Tyr
85     90     95
Glu Lys Arg Phe Gly Pro Tyr Tyr Thr Glu Pro Val Ile Ala Gly Leu
100    105    110
Asp Pro Lys Thr Phe Lys Pro Phe Ile Cys Ser Leu Asp Leu Ile Gly
25     115    120    125
Cys Pro Met Val Thr Asp Asp Phe Val Val Ser Gly Thr Cys Ala Glu
130    135    140
Gln Met Tyr Gly Met Cys Glu Ser Leu Trp Glu Pro Asn Met Asp Pro
30     145    150    155    160
Asp His Leu Phe Glu Thr Ile Ser Gln Ala Met Leu Asn Ala Val Asp
165    170    175
Arg Asp Ala Val Ser Gly Met Gly Val Ile Val His Ile Ile Glu Lys
180    185    190
Asp Lys Ile Thr Thr Arg Thr Leu Lys Ala Arg Met Asp
35     195    200    205

```

<210> 83

<211> 190

<212> PRT

<213> Homo sapiens

<400> 83

```

Leu Thr Arg Ser Cys Ser Thr Cys Cys Pro Ala Val Ala Cys Leu Val
1      5      10      15
Gly Arg Gly Val Val Thr Ser Gly Ala Met His Gln Cys Trp Gly Glu
50     20     25     30
Glu Met Leu Gln Gly Met Leu Leu Trp Gly Trp Ala Thr Cys Pro Leu
35     40     45
Ser Asn Pro Gly Arg Trp Gly Arg Thr Val Gly Leu Gln His Pro Ala
50     55     60
Val Val Ser Ala Phe Arg Ala Leu Leu Leu Met Leu Thr Val His
65     70     75     80

```

Val Ser Tyr Leu Ser Leu Ile Arg Phe Asp Tyr Gly Tyr Asn Leu Val
 85
 Ala Asn Val Ala Ile Gly Leu Val Asn Val Val Trp Trp Leu Ala Trp
 100 105
 Cys Leu Trp Asn Gln Arg Arg Leu Pro His Val Arg Lys Cys Val Val
 115 120 125
 Val Val Leu Leu Leu Gln Gly Leu Ser Leu Leu Glu Leu Leu Asp Phe
 130 135 140
 Pro Pro Leu Phe Trp Val Leu Asp Ala His Ala Ile Trp His Ile Ser
 145 150 155 160
 Thr Ile Pro Val His Val Leu Phe Phe Ser Phe Leu Glu Asp Asp Ser
 165 170 175
 Leu Tyr Leu Leu Lys Glu Ser Glu Asp Lys Phe Lys Leu Asp
 180 185 190

<210> 84
 <211> 368
 <212> PRT
 <213> Homo sapiens

<400> 84
 Ala Pro Pro Pro Ala Ala Ser Gln Gly Glu Arg Met Ala Gly Leu Ala
 1 5 10 15
 Ala Arg Leu Val Leu Leu Ala Gly Ala Ala Leu Ala Ser Gly Ser
 20 25 30
 Gln Gly Asp Arg Glu Pro Val Tyr Arg Asp Cys Val Leu Gln Cys Glu
 35 40 45
 Glu Gln Asn Cys Ser Gly Gly Ala Leu Asn His Phe Arg Ser Arg Gln
 50 55 60
 Pro Ile Tyr Met Ser Leu Ala Gly Trp Thr Cys Arg Asp Asp Cys Lys
 65 70 75 80
 Tyr Glu Cys Met Trp Val Thr Val Gly Leu Tyr Leu Gln Glu Gly His
 85 90 95
 Lys Val Pro Gln Phe His Gly Lys Trp Pro Phe Ser Arg Phe Leu Phe
 100 105 110
 Phe Gln Glu Pro Ala Ser Ala Val Ala Ser Phe Leu Asn Gly Leu Ala
 115 120 125
 Ser Leu Val Met Leu Cys Arg Tyr Arg Thr Phe Val Pro Ala Ser Ser
 130 135 140
 Pro Met Tyr His Thr Cys Val Ala Phe Ala Trp Val Ser Leu Asn Ala
 145 150 155 160
 Trp Phe Trp Ser Thr Val Phe His Thr Arg Asp Thr Asp Leu Thr Glu
 165 170 175
 Lys Met Asp Tyr Phe Cys Ala Ser Thr Val Ile Leu His Ser Ile Tyr
 180 185 190
 Leu Cys Cys Val Arg Thr Val Gly Leu Gln His Pro Ala Val Val Ser
 195 200 205
 Ala Phe Arg Ala Leu Leu Leu Leu Met Leu Thr Val His Val Ser Tyr
 210 215 220
 Leu Ser Leu Ile Arg Phe Asp Tyr Gly Tyr Asn Leu Val Ala Asn Val
 225 230 235 240
 Ala Ile Gly Leu Val Asn Val Val Trp Trp Leu Ala Trp Cys Leu Trp
 245 250 255

Asn Gln Arg Arg Leu Pro His Val Arg Lys Cys Val Val Val Val Leu
 260 265 270
 Leu Leu Gln Gly Leu Ser Leu Leu Glu Leu Leu Asp Phe Pro Pro Leu
 275 280 285
 Phe Trp Val Leu Asp Ala His Ala Ile Trp His Ile Ser Thr Ile Pro
 290 295 300
 Val His Val Leu Phe Phe Ser Phe Leu Glu Asp Asp Ser Leu Tyr Leu
 305 310 315 320
 Leu Lys Glu Ser Glu Asp Lys Phe Lys Leu Val Glu Ala Asp Trp Ile
 325 330 335
 Phe Ala Leu Pro Leu Thr Pro Cys Pro Ser Leu Arg Glu Gly Ser Tyr
 340 345 350
 Ala Arg Thr Pro Thr Ser Gly Thr Arg Val Ala Cys Ala Ser Phe Phe
 355 360 365

<210> 85

<211> 190

<212> PRT

<213> Homo sapiens

<400> 85

Leu Thr Arg Ser Cys Ser Thr Cys Cys Pro Ala Val Ala Cys Leu Val
 1 5 10 15
 Gly Arg Gly Val Val Thr Ser Gly Ala Met His Gln Cys Trp Gly Glu
 20 25 30
 Glu Met Leu Gln Gly Met Leu Leu Trp Gly Trp Ala Thr Cys Pro Leu
 35 40 45
 Ser Asn Pro Gly Arg Trp Gly Arg Thr Val Gly Leu Gln His Pro Ala
 50 55 60
 Val Val Ser Ala Phe Arg Ala Leu Leu Leu Leu Met Leu Thr Val His
 65 70 75 80
 Val Ser Tyr Leu Ser Leu Ile Arg Phe Asp Tyr Gly Tyr Asn Leu Val
 85 90 95
 Ala Asn Val Ala Ile Gly Leu Val Asn Val Val Trp Trp Leu Ala Trp
 100 105 110
 Cys Leu Trp Asn Gln Arg Arg Leu Pro His Val Arg Lys Cys Val Val
 115 120 125
 Val Val Leu Leu Leu Gln Gly Leu Ser Leu Leu Glu Leu Leu Asp Phe
 130 135 140
 Pro Pro Leu Phe Trp Val Leu Asp Ala His Ala Ile Trp His Ile Ser
 145 150 155 160
 Thr Ile Pro Val His Val Leu Phe Phe Ser Phe Leu Glu Asp Asp Ser
 165 170 175
 Leu Tyr Leu Leu Lys Glu Ser Glu Asp Lys Phe Lys Leu Asp
 180 185 190

<210> 86

<211> 318

<212> PRT

<213> Homo sapiens

<400> 86

Met Ala Gly Leu Ala Ala Arg Leu Val Leu Leu Ala Gly Ala Ala Ala
 1 5 10 15
 Leu Ala Ser Gly Ser Gln Gly Asp Arg Glu Pro Val Tyr Arg Asp Cys
 20 25 30
 Val Leu Gln Cys Glu Glu Gln Asn Cys Ser Gly Gly Ala Leu Asn His
 35 40 45
 Phe Arg Ser Arg Gln Pro Ile Tyr Met Ser Leu Ala Gly Trp Thr Cys
 50 55 60
 Arg Asp Asp Cys Lys Tyr Glu Cys Met Trp Val Thr Val Gly Leu Tyr
 65 70 75 80
 Leu Gln Glu Gly His Lys Val Pro Gln Phe His Gly Lys Trp Pro Phe
 85 90 95
 Ser Arg Phe Leu Phe Phe Gln Glu Pro Ala Ser Ala Val Ala Ser Phe
 100 105 110
 Leu Asn Gly Leu Ala Ser Leu Val Met Leu Cys Arg Tyr Arg Thr Phe
 115 120 125
 Val Pro Ala Ser Ser Pro Met Tyr His Thr Cys Val Ala Phe Ala Trp
 130 135 140
 Val Ser Leu Asn Ala Trp Phe Trp Ser Thr Val Phe His Thr Arg Asp
 145 150 155 160
 Thr Asp Leu Gln Arg Lys Trp Thr Thr Ser Val Pro Pro Val Ser Tyr
 165 170 175
 Thr Gln Ser Thr Cys Ala Ala Ser Gly Pro Trp Gly Cys Ser Thr Gln
 180 185 190
 Leu Trp Ser Ser Ala Phe Arg Ala Leu Leu Leu Met Leu Thr Val
 195 200 205
 His Val Ser Tyr Leu Ser Leu Ile Arg Phe Asp Tyr Gly Tyr Asn Leu
 210 215 220
 Val Ala Asn Val Ala Ile Gly Leu Val Asn Val Val Trp Trp Leu Ala
 225 230 235 240
 Trp Cys Leu Trp Asn Gln Arg Arg Leu Pro His Val Arg Lys Cys Val
 245 250 255
 Val Val Val Leu Leu Leu Gln Gly Leu Ser Leu Leu Glu Leu Leu Asp
 260 265 270
 Phe Pro Pro Leu Phe Trp Val Leu Asp Ala His Ala Ile Trp His Ile
 275 280 285
 Ser Thr Ile Pro Val His Val Leu Phe Phe Ser Phe Leu Glu Asp Asp
 290 295 300
 Ser Leu Tyr Leu Leu Lys Glu Ser Glu Asp Lys Phe Lys Leu
 305 310 315

<210> 87

<211> 226

<212> PRT

<213> Homo sapiens

<400> 87

Met Ala Gly Leu Ala Ala Arg Leu Val Leu Leu Ala Gly Ala Ala Ala
 1 5 10 15
 Leu Ala Ser Gly Ser Gln Gly Asp Arg Glu Pro Val Tyr Arg Asp Cys
 20 25 30
 Val Leu Gln Cys Glu Glu Gln Asn Cys Ser Gly Gly Ala Leu Asn His
 35 40 45
 Phe Arg Ser Arg Gln Pro Ile Tyr Met Ser Leu Ala Gly Trp Thr Cys
 50 55 60
 Arg Asp Asp Cys Lys Tyr Glu Cys Met Trp Val Thr Val Gly Leu Tyr
 65 70 75 80
 Leu Gln Glu Gly His Lys Val Pro Gln Phe His Gly Lys Trp Pro Phe
 85 90 95
 Ser Arg Phe Leu Phe Phe Gln Glu Pro Ala Ser Ala Val Ala Ser Phe
 100 105 110
 Leu Asn Gly Leu Ala Ser Leu Val Met Leu Cys Arg Tyr Arg Thr Phe
 115 120 125
 Val Pro Ala Ser Ser Pro Met Tyr His Thr Cys Val Ala Phe Ala Trp
 130 135 140
 Val Ser Leu Asn Ala Trp Phe Trp Ser Thr Val Phe His Thr Arg Asp
 145 150 155 160
 Thr Asp Leu Thr Glu Lys Met Asp Tyr Phe Cys Ala Ser Thr Val Ile
 165 170 175
 Leu His Ser Ile Tyr Leu Cys Cys Val Arg Pro Gly Gln Arg Gly Val
 180 185 190
 Val Ala Gly Leu Val Pro Val Glu Pro Ala Ala Ala Ser Arg Ala
 195 200 205
 Gln Val Arg Gly Gly Gly Leu Ala Ala Ala Gly Ala Val Pro Ala Arg
 210 215 220
 Ala Ala
 225

<210> 88

<211> 320

<212> PRT

<213> Homo sapiens

<400> 88

Met Ala Gly Leu Ala Ala Arg Leu Val Leu Leu Ala Gly Ala Ala Ala
 1 5 10 15
 Leu Ala Ser Gly Ser Gln Gly Asp Arg Glu Pro Val Tyr Arg Asp Cys
 20 25 30
 Val Leu Gln Cys Glu Glu Gln Asn Cys Ser Gly Gly Ala Leu Asn His
 35 40 45
 Phe Arg Ser Arg Gln Pro Ile Tyr Met Ser Leu Ala Gly Trp Thr Cys
 50 55 60
 Arg Asp Asp Cys Lys Tyr Glu Cys Met Trp Val Thr Val Gly Leu Tyr
 65 70 75 80
 Leu Gln Glu Gly His Lys Val Pro Gln Phe His Gly Lys Trp Pro Phe
 85 90 95
 Ser Arg Phe Leu Phe Phe Gln Glu Pro Ala Ser Ala Val Ala Ser Phe
 100 105 110
 Leu Asn Gly Leu Ala Ser Leu Val Met Leu Cys Arg Tyr Arg Thr Phe
 115 120 125

Val Pro Ala Ser Ser Pro Met Tyr His Thr Cys Val Ala Phe Ala Trp
 130 135 140
 Val Ser Leu Asn Ala Trp Phe Trp Ser Thr Val Phe His Thr Arg Asp
 145 150 155 160
 Thr Asp Leu Thr Glu Lys Met Asp Tyr Phe Cys Ala Ser Thr Val Ile
 165 170 175
 Leu His Ser Ile Tyr Leu Cys Cys Val Arg Thr Val Gly Leu Gln His
 180 185 190
 Pro Ala Val Val Ser Ala Phe Arg Ala Leu Leu Leu Leu Mat Leu Thr
 195 200 205
 Val His Val Ser Tyr Leu Ser Leu Ile Arg Phe Asp Tyr Gly Tyr Asn
 210 215 220
 Leu Val Ala Asn Val Ala Ile Gly Leu Val Asn Val Val Trp Trp Leu
 225 230 235 240
 Ala Trp Cys Leu Trp Asn Gln Arg Arg Leu Pro His Val Arg Lys Cys
 245 250 255
 Val Val Val Val Leu Leu Leu Gln Gly Leu Ser Leu Leu Glu Leu Leu
 260 265 270
 Asp Phe Pro Pro Leu Phe Trp Val Leu Asp Ala His Ala Ile Trp His
 275 280 285
 Ile Ser Thr Ile Pro Val His Val Leu Phe Phe Ser Phe Leu Glu Asp
 290 295 300
 Asp Ser Leu Tyr Leu Leu Lys Glu Ser Glu Asp Lys Phe Lys Leu Asp
 305 310 315 320

 <210> 89

 <211> 217

 <212> PRT

 <213> Homo sapiens

 <400> 89
 Ala Pro Pro Pro Ala Ala Ser Gln Gly Glu Arg Met Ala Gly Leu Ala
 1 5 10 15
 Ala Arg Leu Val Leu Leu Ala Gly Ala Ala Leu Ala Ser Gly Ser
 20 25 30
 Gln Gly Asp Arg Glu Pro Val Tyr Arg Asp Cys Val Leu Gln Cys Glu
 35 40 45
 Glu Gln Asn Cys Ser Gly Gly Ala Leu Asn His Phe Arg Ser Arg Gln
 50 55 60
 Pro Ile Tyr Met Ser Leu Ala Gly Trp Thr Cys Arg Asp Asp Cys Lys
 65 70 75 80
 Tyr Glu Cys Met Trp Val Thr Val Gly Leu Tyr Leu Gln Glu Gly His
 85 90 95
 Lys Val Pro Gln Phe His Gly Lys Trp Pro Phe Ser Arg Phe Leu Phe
 100 105 110
 Phe Gln Glu Pro Ala Ser Ala Val Ala Ser Phe Leu Asn Gly Leu Ala
 115 120 125
 Ser Leu Val Met Leu Cys Arg Tyr Arg Thr Phe Val Pro Ala Ser Ser
 130 135 140
 Pro Met Tyr His Thr Cys Val Ala Phe Ala Trp Val Ser Leu Asn Ala
 145 150 155 160
 Trp Phe Trp Ser Thr Val Phe His Thr Arg Asp Thr Asp Leu Thr Glu
 165 170 175

Lys Met Asp Tyr Phe Cys Ala Ser Thr Val Ile Leu His Ser Ile Tyr
 180 185 190
 5 Leu Cys Cys Val Ser Phe Leu Glu Asp Asp Ser Leu Tyr Leu Leu Lys
 195 200 205
 Glu Ser Glu Asp Lys Phe Lys Leu Asp
 210 215
 <210> 90
 <211> 153
 <212> PRT
 <213> Homo sapiens
 <400> 90
 20 Met Asn Val Gly Thr Ala His Ser Glu Val Asn Pro Asn Thr Arg Val
 1 5 10 15
 Met Asn Ser Arg Gly Ile Trp Leu Ser Tyr Val Leu Ala Ile Gly Leu
 20 25 30
 Leu His Ile Val Leu Leu Ser Ile Pro Phe Val Ser Val Pro Val Val
 35 40 45
 25 Trp Thr Leu Thr Asn Leu Ile His Asn Met Gly Met Tyr Ile Phe Leu
 50 55 60
 His Thr Val Lys Gly Thr Pro Phe Glu Thr Pro Asp Gln Gly Lys Ala
 65 70 75 80
 Arg Leu Leu Thr His Trp Glu Gln Met Asp Tyr Gly Val Gln Phe Thr
 85 90 95
 30 Ala Ser Arg Lys Phe Leu Thr Ile Thr Pro Ile Val Leu Tyr Phe Leu
 100 105 110
 Thr Ser Phe Tyr Thr Lys Tyr Asp Gln Ile His Phe Val Leu Asn Thr
 115 120 125
 Val Ser Leu Met Ser Val Leu Ile Pro Lys Leu Pro Gln Leu His Gly
 130 135 140
 35 Val Arg Ile Phe Gly Ile Asn Lys Tyr
 145 150
 <210> 91
 <211> 436
 <212> PRT
 <213> Homo sapiens
 <400> 91
 Met Arg Arg Asp Val Asn Gly Val Thr Lys Ser Arg Phe Glu Met Phe
 1 5 10 15
 50 Ser Asn Ser Asp Glu Ala Val Ile Asn Lys Lys Leu Pro Lys Glu Leu
 20 25 30
 Leu Leu Arg Ile Phe Ser Phe Leu Asp Val Val Thr Leu Cys Arg Cys
 35 40 45
 Ala Gln Val Ser Arg Ala Trp Asn Val Leu Ala Leu Asp Gly Ser Asn
 50 55 60

Trp Gln Arg Ile Asp Leu Phe Asp Phe Gln Arg Asp Ile Glu Gly Arg
 65 70 75 80
 Val Val Glu Asn Ile Ser Lys Arg Cys Gly Gly Phe Leu Arg Lys Leu
 85 90 95
 Ser Leu Arg Gly Cys Leu Gly Val Gly Asp Asn Ala Leu Arg Thr Phe
 100 105 110
 Ala Gln Asn Cys Arg Asn Ile Glu Val Leu Asn Leu Asn Gly Cys Thr
 115 120 125
 Lys Thr Thr Asp Ala Thr Cys Thr Ser Leu Ser Lys Phe Cys Ser Lys
 130 135 140
 Leu Arg His Leu Asp Leu Ala Ser Cys Thr Ser Ile Thr Asn Met Ser
 145 150 155 160
 Leu Lys Ala Leu Ser Glu Gly Cys Pro Leu Leu Glu Gln Leu Asn Ile
 165 170 175
 Ser Trp Cys Asp Gln Val Thr Lys Asp Gly Ile Gln Ala Leu Val Arg
 180 185 190
 Gly Cys Gly Gly Leu Lys Ala Leu Phe Leu Lys Gly Cys Thr Gln Leu
 195 200 205
 Glu Asp Glu Ala Leu Lys Tyr Ile Gly Ala His Cys Pro Glu Leu Val
 210 215 220
 Thr Leu Asn Leu Gln Thr Cys Leu Gln Ile Thr Asp Glu Gly Leu Ile
 225 230 235 240
 Thr Ile Cys Arg Gly Cys His Lys Leu Gln Ser Leu Cys Ala Ser Gly
 245 250 255
 Cys Ser Asn Ile Thr Asp Ala Ile Leu Asn Ala Leu Gly Gln Asn Cys
 260 265 270
 Pro Arg Leu Arg Ile Leu Glu Val Ala Arg Cys Ser Gln Leu Thr Asp
 275 280 285
 Val Gly Phe Thr Thr Leu Ala Arg Asn Cys His Glu Leu Lys Met
 290 295 300
 Asp Leu Glu Glu Cys Val Gln Ile Thr Asp Ser Thr Leu Ile Gln Leu
 305 310 315 320
 Ser Ile His Cys Pro Arg Leu Gln Val Leu Ser Leu Ser His Cys Glu
 325 330 335
 Leu Ile Thr Asp Asp Gly Ile Arg His Leu Gly Asn Gly Ala Cys Ala
 340 345 350
 His Asp Gln Leu Glu Val Ile Glu Leu Asp Asn Cys Pro Leu Ile Thr
 355 360 365
 Asp Ala Ser Leu Glu His Leu Lys Ser Cys His Ser Leu Glu Arg Ile
 370 375 380
 Glu Leu Tyr Asp Cys Gln Gln Ile Thr Arg Ala Gly Ile Lys Arg Leu
 385 390 395 400
 Arg Thr His Leu Pro Asn Ile Lys Val His Ala Tyr Phe Ala Pro Val
 405 410 415
 Thr Pro Pro Pro Ser Val Gly Gly Ser Arg Gln Arg Phe Cys Arg Cys
 420 425 430
 Cys Ile Ile Leu
 435

<210> 92

<211> 204

<212> PRT

<213> Homo sapiens

<400> 92

Met Asp Pro Lys Asp Arg Lys Lys Ile Gln Phe Ser Val Pro Ala Pro
 1 5 10 15
 Pro Ser Gln Leu Asp Pro Arg Gln Val Glu Met Ile Arg Arg Arg
 20 25 30
 Pro Thr Pro Ala Met Leu Phe Arg Leu Ser Glu His Ser Ser Pro Glu
 35 40 45
 Glu Glu Ala Ser Pro His Gln Arg Ala Ser Gly Glu Gly His His Leu
 50 55 60
 Lys Ser Lys Arg Pro Asn Pro Cys Ala Tyr Thr Pro Pro Ser Leu Lys
 65 70 75 80
 Ala Val Gln Arg Ile Ala Glu Ser His Leu Gln Ser Ile Ser Asn Leu
 85 90 95
 Asn Glu Asn Gln Ala Ser Glu Glu Glu Asp Glu Leu Gly Glu Leu Arg
 100 105 110
 Glu Leu Gly Tyr Pro Arg Glu Glu Asp Glu Glu Glu Glu Asp Asp
 115 120 125
 Glu Glu Glu Glu Glu Glu Asp Ser Gln Ala Glu Val Leu Lys Val
 130 135 140
 Ile Arg Gln Ser Ala Gly Gln Lys Thr Thr Cys Gly Gln Gly Leu Glu
 145 150 155 160
 Gly Pro Trp Glu Arg Pro Pro Leu Asp Glu Ser Glu Arg Asp Gly
 165 170 175
 Gly Ser Glu Asp Gln Val Glu Asp Pro Ala Leu Ser Glu Pro Gly Glu
 180 185 190
 Glu Pro Gln Arg Pro Ser Pro Ser Glu Pro Gly Thr
 195 200

<210> 93

<211> 115

<212> PRT

<213> Homo sapiens

<400> 93

Met Ser Gly Glu Pro Gly Gln Thr Ser Val Ala Pro Pro Pro Glu Glu
 1 5 10 15
 Val Glu Pro Gly Ser Gly Val Arg Ile Val Val Glu Tyr Cys Glu Pro
 20 25 30
 Cys Gly Phe Glu Ala Thr Tyr Leu Glu Leu Ala Ser Ala Val Lys Glu
 35 40 45
 Gln Tyr Pro Gly Ile Glu Ile Glu Ser Arg Leu Gly Gly Thr Gly Ala
 50 55 60
 Phe Glu Ile Glu Ile Asn Gly Gln Leu Val Phe Ser Lys Leu Glu Asn
 65 70 75 80
 Gly Gly Phe Pro Tyr Glu Lys Asp Leu Ile Glu Ala Ile Arg Arg Ala
 85 90 95
 Ser Asn Gly Glu Thr Leu Glu Lys Ile Thr Asn Ser Arg Pro Pro Cys
 100 105 110
 Val Ile Leu
 115

<210> 94

<211> 144

<212> PRT

<213> Homo sapiens

<400> 94

```

Met Gly Ala Val Val Leu Cys Arg Pro Ser Pro Leu Asn Phe Leu Ile
1      5      10      15
Gln Thr Gly Thr Gly Gln Gly Leu Ser Cys Gly Ser His Met Trp Arg
      20      25      30
Cys Glu Ala Thr Pro Cys Gly Val Cys Gly Glu Ser Pro Val Gly Ser
15     35     40     45
Leu Leu Lys Gln His Arg Gly Arg Gly Lys Thr Trp Pro Val Gly Thr
      50     55     60
Val Ser Ala Cys Arg Glu Glu Ser Glu Ala Gly Ser Leu Ser Leu Gly
65     70     75     80
Trp Ser Leu Leu Pro Ser Pro Val Gly Leu Gly Ala Val Leu Ile Leu
20     85     90     95
Lys Arg Cys Gly Ser Leu Cys Pro Leu Pro Gly Val Gln Gly Asn Arg
      100    105    110
Arg Gly His Trp Ala Cys Phe Leu Pro Pro Asp Pro Ala Ser Pro Thr
115    120    125
25 Pro Cys Ile Ile Gly Asn Phe His Leu Lys Ile Phe Leu Ser Lys Val
      130    135    140

```

<210> 95

<211> 425

<212> PRT

<213> Homo sapiens

<400> 95

```

Met Gly Gly Gly Asp Leu Asn Leu Lys Lys Ser Trp His Pro Gln Thr
1      5      10      15
Leu Arg Asn Val Glu Lys Val Trp Lys Ala Glu Gln Lys His Glu Ala
40     20     25     30
Glu Arg Lys Lys Ile Glu Glu Leu Gln Arg Glu Leu Arg Glu Glu Arg
      35     40     45
Ala Arg Glu Glu Met Gln Arg Tyr Ala Glu Asp Val Gly Ala Val Lys
45     50     55     60
Lys Lys Glu Glu Lys Leu Asp Trp Met Tyr Gln Gly Pro Gly Gly Met
65     70     75     80
Val Asn Arg Asp Glu Tyr Leu Leu Gly Arg Pro Ile Asp Lys Tyr Val
      85     90     95
Phe Glu Lys Met Glu Glu Lys Glu Ala Gly Cys Ser Ser Glu Thr Gly
50     100    105    110
Leu Leu Pro Gly Ser Ile Phe Ala Pro Ser Gly Ala Asn Ser Leu Leu
115    120    125
Asp Met Ala Ser Lys Ile Arg Glu Asp Pro Leu Phe Ile Ile Arg Lys
130    135    140
55 Lys Glu Glu Glu Lys Lys Arg Glu Val Leu Asn Asn Pro Val Lys Met
      145    150    155    160

```


Lys Lys Ile Lys Glu Leu Leu Gln Met Ser Leu Glu Lys Lys Glu Lys
 165
 Lys Lys Lys Lys Glu Lys Lys Lys His Lys Lys His Lys His Arg
 180
 Ser Ser Ser Ser Asp Arg Ser Ser Ser Ser Glu Asp Glu His Ser Ala Gly
 195
 Arg Ser Gln Lys Lys Met Ala Asn Ser Ser Pro Val Leu Ser Lys Val
 210
 Pro Gly Tyr Gly Leu Gln Val Arg Asn Ser Asp Arg Asn Gln Gly Leu
 225
 Gln Gly Pro Leu Thr Ala Glu Gln Lys Arg Gly His Gly Met Lys Asn
 245
 His Ser Arg Ser Arg Ser Ser Ser His Ser Pro Pro Arg His Ala Ser
 260
 Lys Lys Ser Thr Arg Glu Ala Gly Ser Arg Asp Arg Arg Ser Arg Ser
 275
 Leu Gly Arg Arg Ser Arg Ser Pro Arg Pro Ser Lys Leu His Asn Ser
 290
 Lys Val Asn Arg Arg Glu Thr Gly Gln Thr Arg Ser Pro Ser Pro Lys
 305
 Lys Glu Val Tyr Gln Arg Arg His Ala Pro Gly Tyr Thr Arg Lys Leu
 325
 Ser Ala Glu Glu Leu Glu Arg Lys Arg Gln Glu Met Met Glu Asn Ala
 340
 Lys Trp Arg Glu Glu Glu Arg Leu Asn Ile Leu Lys Arg His Ala Lys
 355
 Asp Glu Glu Arg Glu Gln Arg Leu Glu Lys Leu Asp Ser Arg Asp Gly
 370
 Lys Phe Ile His Arg Met Lys Leu Glu Ser Ala Ser Thr Ser Ser Leu
 385
 Glu Asp Arg Val Lys Arg Asn Ile Tyr Ser Leu Gln Arg Thr Ser Val
 405
 Ala Leu Glu Lys Asn Phe Met Lys Arg
 420
 425
 <210> 96
 <211> 394
 <212> PRT
 <213> Homo sapiens
 <400> 96
 Met Phe Ser Val Phe Glu Glu Ile Thr Arg Ile Val Val Lys Glu Met
 1
 Asp Ala Gly Gly Asp Met Ile Ala Val Arg Ser Leu Val Asp Ala Asp
 20
 Arg Phe Arg Cys Phe His Leu Val Gly Glu Lys Arg Thr Phe Phe Gly
 35
 Cys Arg His Tyr Thr Thr Gly Leu Thr Leu Met Asp Ile Leu Asp Thr
 50
 His Gly Asp Lys Trp Leu Asp Glu Leu Asp Ser Gly Leu Gln Gly Gln
 65
 Lys Ala Glu Phe Gln Ile Leu Asp Asn Val Asp Ser Thr Gly Glu Leu
 85
 90
 95

Ile Val Arg Leu Pro Lys Glu Ile Thr Ile Ser Gly Ser Phe Gln Gly
 100 105 110
 Phe His His Gln Lys Ile Lys Ile Ser Glu Asn Arg Ile Ser Gln Gln
 115 120
 Tyr Leu Ala Thr Leu Glu Asn Arg Lys Leu Lys Arg Glu Leu Pro Phe
 130 135 140
 Ser Phe Arg Ser Ile Asn Thr Arg Glu Asn Leu Tyr Leu Val Thr Glu
 145 150 155 160
 Thr Leu Glu Thr Val Lys Glu Glu Thr Leu Lys Ser Asp Arg Gln Tyr
 165 170 175
 Lys Phe Trp Ser Gln Ile Ser Gln Gly His Leu Ser Tyr Lys His Lys
 180 185 190
 Gly Gln Arg Glu Val Thr Ile Pro Pro Asn Arg Val Leu Ser Tyr Arg
 195 200 205
 Val Lys Gln Leu Val Phe Pro Asn Lys Glu Thr Met Arg Lys Ser Leu
 210 215 220
 Gly Ser Glu Asp Ser Arg Asn Met Lys Glu Lys Leu Glu Asp Met Glu
 225 230 235 240
 Ser Val Leu Lys Asp Leu Thr Glu Glu Lys Arg Lys Asp Val Leu Asn
 245 250 255
 Ser Leu Ala Lys Cys Leu Gly Lys Glu Asp Ile Arg Gln Asp Leu Glu
 260 265 270
 Gln Arg Val Ser Glu Val Leu Ile Ser Gly Glu Leu His Met Glu Asp
 275 280 285
 Pro Asp Lys Pro Leu Leu Ser Ser Leu Phe Asn Ala Ala Gly Val Leu
 290 295 300
 Val Glu Ala Arg Ala Lys Ala Ile Leu Asp Phe Leu Asp Ala Leu Leu
 305 310 315 320
 Glu Leu Ser Glu Glu Gln Gln Phe Val Ala Glu Ala Leu Glu Lys Gly
 325 330 335
 Thr Leu Pro Leu Leu Lys Asp Gln Val Lys Ser Val Met Glu Gln Asn
 340 345 350
 Trp Asp Glu Leu Ala Ser Ser Pro Pro Asp Met Asp Tyr Asp Pro Glu
 355 360 365
 Ala Arg Ile Leu Cys Ala Leu Tyr Val Val Val Ser Ile Leu Leu Glu
 370 375 380
 Leu Ala Glu Gly Pro Thr Ser Val Ser Ser
 385 390

 <210> 97

 <211> 456

 <212> PRT

 <213> Homo sapiens

 <400> 97
 Met Glu Gly Pro Glu Gly Leu Gly Arg Lys Gln Ala Cys Leu Ala Met
 1 5 10 15
 Leu Leu His Phe Leu Asp Thr Tyr Gln Gly Leu Leu Gln Glu Glu
 20 25 30
 Gly Ala Gly His Ile Ile Lys Asp Leu Tyr Leu Leu Ile Met Lys Asp
 35 40 45
 Glu Ser Leu Tyr Gln Gly Leu Arg Glu Asp Thr Leu Arg Leu His Gln
 50 55 60

5 Leu Val Glu Thr Val Glu Leu Lys Ile Pro Glu Glu Asn Gln Pro Pro
 65 70 75 80
 Ser Lys Gln Val Lys Pro Leu Phe Arg His Phe Arg Arg Ile Asp Ser
 85 90 95
 Cys Leu Gln Thr Arg Val Ala Phe Arg Gly Ser Asp Glu Ile Phe Cys
 100 105 110
 Arg Val Tyr Met Pro Asp His Ser Tyr Val Thr Ile Arg Ser Arg Leu
 115 120 125
 10 Ser Ala Ser Val Gln Asp Ile Leu Gly Ser Val Thr Glu Lys Leu Gln
 130 135 140
 Tyr Ser Glu Glu Pro Ala Gly Arg Glu Asp Ser Leu Ile Leu Val Ala
 145 150 155 160
 Val Ser Ser Ser Gly Glu Lys Val Leu Leu Gln Pro Thr Glu Asp Cys
 165 170 175
 15 Val Phe Thr Ala Leu Gly Ile Asn Ser His Leu Phe Ala Cys Thr Arg
 180 185 190
 Asp Ser Tyr Glu Ala Leu Val Pro Leu Pro Glu Glu Ile Gln Val Ser
 195 200 205
 Pro Gly Asp Thr Glu Ile His Arg Val Glu Pro Glu Asp Val Ala Asn
 210 215 220
 20 His Leu Thr Ala Phe His Trp Glu Leu Phe Arg Cys Val His Glu Leu
 225 230 235 240
 Glu Phe Val Asp Tyr Val Phe His Gly Glu Arg Gly Arg Arg Glu Thr
 245 250 255
 25 Ala Asn Leu Glu Leu Leu Gln Arg Cys Ser Glu Val Thr His Trp
 260 265 270
 Val Ala Thr Glu Val Leu Leu Cys Glu Ala Pro Gly Lys Arg Ala Gln
 275 280 285
 Leu Leu Lys Lys Phe Ile Lys Ile Ala Ala Leu Cys Lys Gln Asn Gln
 290 295 300
 30 Asp Leu Leu Ser Phe Tyr Ala Val Val Met Gly Leu Asp Asn Ala Ala
 305 310 315 320
 Val Ser Arg Leu Arg Leu Thr Trp Glu Lys Leu Pro Gly Lys Phe Lys
 325 330 335
 Asn Leu Phe Arg Lys Phe Glu Asn Leu Thr Asp Pro Cys Arg Asn His
 340 345 350
 35 Lys Ser Tyr Arg Glu Val Ile Ser Lys Met Lys Pro Pro Val Ile Pro
 355 360 365
 Phe Val Pro Leu Ile Leu Lys Asp Leu Thr Phe Leu His Glu Gly Ser
 370 375 380
 Lys Thr Leu Val Asp Gly Leu Val Asn Ile Glu Lys Leu His Ser Val
 385 390 395 400
 40 Ala Glu Lys Val Arg Thr Ile Arg Lys Tyr Arg Ser Arg Pro Leu Cys
 405 410 415
 Leu Asp Met Glu Ala Ser Pro Asn His Leu Gln Thr Lys Ala Tyr Val
 420 425 430
 45 Arg Gln Phe Gln Val Ile Asp Asn Gln Asn Leu Leu Phe Glu Leu Ser
 435 440 445
 Tyr Lys Leu Glu Ala Asn Ser Gln
 450 455

<210> 98
 <211> 715
 <212> PRT
 <213> Homo sapiens

<400> 98
 5 Met Ser Gln Val Met Ser Ser Pro Leu Leu Ala Gly Gly His Ala Val
 1 5 10 15
 Ser Leu Ala Pro Cys Asp Glu Pro Arg Thr Leu His Pro Ala Pro
 20 25 30
 Ser Pro Ser Leu Pro Pro Gln Cys Ser Tyr Tyr Thr Thr Gly Trp
 35 40 45
 10 Gly Ala Gln Ala Leu Met Ala Pro Val Pro Cys Met Gly Pro Pro Gly
 50 55 60
 Arg Leu Gln Gln Ala Pro Gln Val Glu Ala Lys Ala Thr Cys Phe Leu
 65 70 75 80
 15 Pro Ser Pro Gly Glu Lys Ala Leu Gly Thr Pro Glu Asp Leu Asp Ser
 85 90 95
 Tyr Ile Asp Phe Ser Leu Glu Ser Leu Asn Gln Met Ile Leu Glu Leu
 100 105 110
 Asp Pro Thr Phe Gln Leu Leu Pro Pro Gly Thr Gly Gly Ser Gln Ala
 115 120 125
 20 Glu Leu Ala Gln Ser Thr Met Ser Met Arg Lys Lys Glu Glu Ser Glu
 130 135 140
 Ala Leu Asp Ile Lys Tyr Ile Glu Val Thr Ser Ala Arg Ser Arg Cys
 145 150 155 160
 His Asp Trp Pro Gln His Cys Ser Ser Pro Ser Val Thr Pro Pro Phe
 165 170 175
 25 Gly Ser Pro Arg Ser Gly Gly Leu Leu Leu Ser Arg Asp Val Pro Arg
 180 185 190
 Glu Thr Arg Ser Ser Ser Glu Ser Leu Ile Phe Ser Gly Asn Gln Gly
 195 200 205
 Arg Gly His Gln Arg Pro Leu Pro Pro Ser Glu Gly Leu Ser Pro Arg
 210 215 220
 30 Pro Pro Asn Ser Pro Ser Ile Ser Ile Pro Cys Met Gly Ser Lys Ala
 225 230 235 240
 Ser Ser Pro His Gly Leu Gly Ser Pro Leu Val Ala Ser Pro Arg Leu
 245 250 255
 Glu Lys Arg Leu Gly Gly Leu Ala Pro Gln Arg Gly Ser Arg Ile Ser
 260 265 270
 35 Val Leu Ser Ala Ser Pro Val Ser Asp Val Ser Tyr Met Phe Gly Ser
 275 280 285
 Ser Gln Ser Leu Leu His Ser Ser Asn Ser Ser His Gln Ser Ser Ser
 290 295 300
 Arg Ser Leu Glu Ser Pro Ala Asn Ser Ser Ser Ser Leu His Ser Leu
 305 310 315 320
 40 Gly Ser Val Ser Leu Cys Thr Arg Pro Ser Asp Phe Gln Ala Pro Arg
 325 330 335
 Asn Pro Thr Leu Thr Met Gly Gln Pro Arg Thr Pro His Ser Pro Pro
 340 345 350
 45 Leu Ala Lys Glu His Ala Ser Ile Cys Pro Pro Ser Ile Thr Asn Ser
 355 360 365
 Met Val Asp Ile Pro Ile Val Leu Ile Asn Gly Cys Pro Glu Pro Gly
 370 375 380
 Ser Ser Pro Pro Gln Arg Thr Pro Gly His Gln Asn Ser Val Gln Pro
 385 390 395 400
 50 Gly Ala Ala Ser Pro Ser Asn Pro Cys Pro Ala Thr Arg Ser Asn Ser
 405 410 415
 Gln Thr Leu Ser Asp Ala Pro Phe Thr Thr Cys Pro Glu Gly Pro Ala
 420 425 430
 Arg Asp Met Gln Pro Thr Met Lys Phe Val Met Asp Thr Ser Lys Tyr
 435 440 445

Trp Phe Lys Pro Asn Ile Thr Arg Glu Gln Ala Ile Glu Leu Leu Arg
 450 455 460
 Lys Glu Glu Pro Gly Ala Phe Val Ile Arg Asp Ser Ser Ser Tyr Arg
 465 470 475
 Gly Ser Phe Gly Leu Ala Leu Lys Val Gln Glu Val Pro Ala Ser Ala
 485 490 495
 Gln Asn Arg Pro Gly Glu Asp Ser Asn Asp Leu Ile Arg His Phe Leu
 500 505 510
 Ile Glu Ser Ser Ala Lys Gly Val His Leu Lys Gly Ala Asp Glu Glu
 515 520 525
 Pro Tyr Phe Gly Ser Leu Ser Ala Phe Val Cys Gln His Ser Ile Met
 530 535 540
 Ala Leu Ala Leu Pro Cys Lys Leu Thr Ile Pro Gln Arg Glu Leu Gly
 545 550 555
 Gly Ala Asp Gly Ala Ser Asp Ser Thr Asp Ser Pro Ala Ser Cys Gln
 565 570 575
 Lys Lys Ser Ala Gly Cys His Thr Leu Tyr Leu Ser Ser Val Ser Val
 580 585 590
 Glu Thr Leu Thr Gly Ala Leu Ala Val Gln Lys Ala Ile Ser Thr Thr
 595 600 605
 Phe Glu Arg Asp Ile Leu Pro Thr Thr Val Val His Phe Glu Val
 610 615 620
 Thr Glu Gln Gly Ile Thr Leu Thr Asp Val Gln Arg Lys Val Phe Phe
 625 630 635
 Arg Arg His Tyr Pro Leu Thr Thr Leu Arg Phe Cys Gly Met Asp Pro
 645 650 655
 Glu Gln Arg Lys Trp Gln Lys Tyr Cys Lys Pro Ser Trp Ile Phe Gly
 660 665 670
 Phe Val Ala Lys Ser Gln Thr Glu Pro Gln Glu Asn Val Cys His Leu
 675 680 685
 Phe Ala Glu Tyr Asp Met Val Gln Pro Ala Ser Gln Val Ile Gly Leu
 690 695 700
 Val Thr Ala Leu Leu Gln Asp Ala Glu Arg Met
 705 710 715

35 <210> 99
 <211> 35
 <212> DNA
 40 <213> Artificial sequence

45 <220>
 <223> PCR primer
 <400> 99
 ccataataa aaccactgtc ctgtcctttg tggct

50 <210> 100
 <211> 26
 <212> DNA
 55 <213> Artificial sequence

35

5	<220>	
	<223> PCR primer	
	<400> 100	
	cccccatctg totgtctata ttgttc	26
10	<210> 101	
	<211> 22	
15	<212> DNA	
	<213> Artificial sequence	
20	<220>	
	<223> PCR primer	
	<400> 101	
	tgccctaogct gaogactatg tg	22
25	<210> 102	
	<211> 25	
30	<212> DNA	
	<213> Artificial sequence	
35	<220>	
	<223> PCR primer	
	<400> 102	
	tttgggttttc tacaactgtt gotat	25
40	<210> 103	
	<211> 19	
45	<212> DNA	
	<213> Artificial sequence	
50	<220>	
	<223> PCR primer	
	<400> 103	
	gggctccaca caccagatg	19
55		

<210> 104
 <211> 21
 <212> DNA
 <213> Artificial sequence
 <220>
 <223> PCR primer
 <400> 104
 acgctctgag caccctctac a 21
 <210> 105
 <211> 31
 <212> DNA
 <213> Artificial sequence
 <220>
 <223> PCR primer
 <400> 105
 tgtcacaggy actgaaaacc tctctcatg t 31
 <210> 106
 <211> 17
 <212> DNA
 <213> Artificial sequence
 <220>
 <223> PCR primer
 <400> 106
 cccaaggcca cgagctt 17
 <210> 107
 <211> 24
 <212> DNA
 <213> Artificial sequence

	<220>	
5	<223> PCR primer	
	<400> 107	24
	tggtgctctc ttaacgaatc gaaa	
	<210> 108	
10	<211> 29	
	<212> DNA	
	<213> Artificial sequence	
15	<220>	
	<223> PCR primer	
20	<400> 108	29
	ctgggtcaaac aaactctctg aacccctcc	
	<210> 109	
25	<211> 20	
	<212> DNA	
	<213> Artificial sequence	
30	<220>	
	<223> PCR primer	
35	<400> 109	20
	tggtgaggaa aagcggacat	
	<210> 110	
	<211> 21	
40	<212> DNA	
	<213> Artificial sequence	
45	<220>	
	<223> PCR primer	
50	<400> 110	21
	ctggccttgga ggacagtga g	
	<210> 111	
55	<211> 24	

<212> DNA
 <213> Artificial sequence
 5
 <220>
 <223> PCR primer
 10 <400> 111
 ccaagccctc cccatcccat gtat 24
 <210> 112
 15 <211> 21
 <212> DNA
 <213> Artificial sequence
 20
 <220>
 <223> PCR primer
 25 <400> 112
 gaggtgtcgt acccggttct a 21
 <210> 113
 30 <211> 21
 <212> DNA
 <213> Artificial sequence
 35
 <220>
 <223> PCR primer
 40 <400> 113
 ccgtttcgtc attccctgc t 21
 <210> 114
 45 <211> 23
 <212> DNA
 <213> Artificial sequence
 50
 <220>
 <223> PCR primer
 55 <400> 114
 ccagaccgc ttcactgacc tgc 23

<210> 115
 5 <211> 20
 <212> DNA
 <213> Artificial sequence
 10
 <220>
 <223> PCR primer
 15 <400> 115 20
 ogcctgtact tcagcatgga
 <210> 116
 20 <211> 18
 <212> DNA
 <213> Artificial sequence
 25
 <220>
 <223> PCR primer
 30 <400> 116 18
 ggggttcagc tgggtgga
 <210> 117
 35 <211> 25
 <212> DNA
 <213> Artificial sequence
 40
 <220>
 <223> PCR primer
 45 <400> 117 25
 accccgaggc atcaccacaa atcat
 <210> 118
 50 <211> 23
 <212> DNA
 <213> Artificial sequence
 55

	<220>	
5	<223> PCR primer	
	<400> 118	
	agttctgct ctctgacaac cat	23
	<210> 119	
10	<211> 23	
	<212> DNA	
	<213> Artificial sequence	
15	<220>	
	<223> PCR primer	
20	<400> 119	
	taggtctcaga gtcagaccca aac	23
	<210> 120	
25	<211> 21	
	<212> DNA	
	<213> Artificial sequence	
30	<220>	
	<223> PCR primer	
35	<400> 120	
	cctctgtggg cttgtctctg g	21
	<210> 121	
	<211> 21	
40	<212> DNA	
	<213> Artificial sequence	
45	<220>	
	<223> PCR primer	
50	<400> 121	
	aagcgcgcag ttoattttt t	21
	<210> 122	
55	<211> 25	

<212> DNA
 <213> Artificial sequence
 5
 <220>
 <223> PCR primer
 <400> 122
 10 cttgtggttc aagtcaaatg ttcag 25
 <210> 123
 <211> 21
 15 <212> DNA
 <213> Artificial sequence
 20
 <220>
 <223> PCR primer
 <400> 123
 25 tctgctgag ctctgtogg t 21
 <210> 124
 <211> 18
 30 <212> DNA
 <213> Artificial sequence
 35
 <220>
 <223> PCR primer
 <400> 124
 40 gggctgggca cctgactt 18
 <210> 125
 <211> 20
 45 <212> DNA
 <213> Artificial sequence
 50
 <220>
 <223> PCR primer
 <400> 125
 55 cccaacaagg gtccagact 20

<210> 126
 5 <211> 17
 <212> DNA
 <213> Artificial sequence
 10
 <220>
 <223> PCR primer
 15 <400> 126 17
 cgggcattg agcggcg
 <210> 127
 20 <211> 20
 <212> DNA
 <213> Artificial sequence
 25
 <220>
 <223> PCR primer
 30 <400> 127 20
 cccaaggac ttogtgaatg
 <210> 128
 35 <211> 21
 <212> DNA
 <213> Artificial sequence
 40
 <220>
 <223> PCR primer
 45 <400> 128 21
 ggcgatcct gatgacaagt a
 <210> 129
 50 <211> 29
 <212> DNA
 <213> Artificial sequence
 55

<220>
 <223> PCR primer#
 5 <400> 129 agcaccaact gtgaaccagg tacaatggc 29
 <210> 130
 <211> 19
 <212> DNA
 <213> Artificial sequence
 15
 <220>
 <223> PCR primer
 20 <400> 130 gagggaggct ctgctttgg 19
 <210> 131
 <211> 21
 25 <212> DNA
 <213> Artificial sequence
 30
 <220>
 <223> PCR primer
 35 <400> 131 tcacaactag cgggtgagga g 21
 <210> 132
 <211> 21
 40 <212> DNA
 <213> Artificial sequence
 45
 <220>
 <223> PCR primer
 50 <400> 132 tgcagaggaa cggcgtgagc g 21
 <210> 133
 <211> 22
 55

	<212> DNA	
5	<213> Artificial sequence	
	<220>	
10	<223> PCR primer	
	<400> 133	
	tgagggtttcc tcccaaatcg ta	22
	<210> 134	
15	<211> 22	
	<212> DNA	
20	<213> Artificial sequence	
	<220>	
25	<223> PCR primer	
	<400> 134	
	cagctcaagg gaagctgtca to	22
	<210> 135	
30	<211> 24	
	<212> DNA	
35	<213> Artificial sequence	
	<220>	
40	<223> PCR primer	
	<400> 135	
	ccccacatg ttcccaaga tgct	24
	<210> 136	
45	<211> 21	
	<212> DNA	
50	<213> Artificial sequence	
	<220>	
55	<223> PCR primer	
	<400> 136	
	ggaggcgcta aaggtctacg t	21

<210> 137
 5 <211> 21
 <212> DNA
 <213> Artificial sequence
 10
 <220>
 <223> PCR primer
 15 <400> 137
 tgatgcttcg caggtcagta a 21
 <210> 138
 20 <211> 26
 <212> DNA
 <213> Artificial sequence
 25
 <220>
 <223> PCR primer
 30 <400> 138
 otctgccc tctaaagct gaagcc 26
 <210> 139
 35 <211> 17
 <212> DNA
 <213> Artificial sequence
 40
 <220>
 <223> PCR primer
 45 <400> 139
 ggacgctgg gcttttc 17
 <210> 140
 50 <211> 20
 <212> DNA
 <213> Artificial sequence
 55

<220>
 <223> PCR primer
 5 <400> 140 20
 tgtgggtgtg gacaccttc
 <210> 141
 <211> 25
 <212> DNA
 <213> Artificial sequence
 15
 <220>
 <223> PCR primer
 20 <400> 141 25
 ccacagctg aaggcagaca aggc
 <210> 142
 <211> 20
 <212> DNA
 <213> Artificial sequence
 25
 <220>
 <223> PCR primer
 30 <400> 142 20
 ggggattctc atggaacaca
 <210> 143
 <211> 20
 <212> DNA
 <213> Artificial sequence
 35
 <220>
 <223> PCR primer
 40 <400> 143 20
 ggtcagccag gagattttg
 <210> 144
 <211> 23
 45
 50
 55

<212> DNA
 <213> Artificial sequence
 5

<220>
 <223> PCR primer
 10 <400> 144
 accacottgc gcaggttgto cag 23
 <210> 145
 15 <211> 18
 <212> DNA
 <213> Artificial sequence
 20

<220>
 <223> PCR primer
 25 <400> 145
 ogcattgcacg acctgaac 18
 <210> 146
 30 <211> 23
 <212> DNA
 <213> Artificial sequence
 35

<220>
 <223> PCR primer
 40 <400> 146
 gtctogattct tggacagott ctg 23
 <210> 147
 45 <211> 22
 <212> DNA
 <213> Artificial sequence
 50

<220>
 <223> PCR primer
 55 <400> 147
 acactgtcca cactgcccga gg 22

<210> 148
 5 <211> 21
 <212> DNA
 <213> Artificial sequence
 10
 <220>
 <223> PCR primer
 15 <400> 148
 ctgggcagaa tggaaggatc t 21
 <210> 149
 20 <211> 22
 <212> DNA
 <213> Artificial sequence
 25
 <220>
 <223> PCR primer
 30 <400> 149
 gggactotag cagaccaca ct 22
 <210> 150
 35 <211> 22
 <212> DNA
 <213> Artificial sequence
 40
 <220>
 <223> PCR primer
 45 <400> 150
 caccacactg gattccctgt tc 22
 <210> 151
 50 <211> 23
 <212> DNA
 <213> Artificial sequence
 55

	<220>	
5	<223> PCR primer	
	<400> 151	23
	ccttcagaca ggctagatg atg	
	<210> 152	
10	<211> 29	
	<212> DNA	
	<213> Artificial sequence	
15	<220>	
	<223> PCR primer	
20	<400> 152	29
	gggtattatt tctttattag gtgocactt	
	<210> 153	
25	<211> 30	
	<212> DNA	
	<213> Artificial sequence	
30	<220>	
	<223> PCR primer	
35	<400> 153	30
	tccoctaagg ctttcagtac ccaggatctg	
	<210> 154	
	<211> 18	
40	<212> DNA	
	<213> Artificial sequence	
45	<220>	
	<223> PCR primer	
50	<400> 154	18
	ccagcttggc cctttcct	
	<210> 155	
55	<211> 23	

<212> DNA
 <213> Artificial sequence
 5
 <220>
 <223> PCR primer
 <400> 155
 gaatgggtcg cttttgttct tag 23
 <210> 156
 <211> 22
 <212> DNA
 <213> Artificial sequence
 20
 <220>
 <223> PCR primer
 <400> 156
 tcaoggacct cagcctgcc ct 22
 <210> 157
 <211> 21
 <212> DNA
 <213> Artificial sequence
 35
 <220>
 <223> PCR primer
 <400> 157
 tggatgaaggt gtcagccatg t 21
 <210> 158
 <211> 21
 <212> DNA
 <213> Artificial sequence
 50
 55

<220>
 5 <223> PCR primer
 <400> 158
 tcagagtgc gcaatggctt t 21
 <210> 159
 10 <211> 20
 <212> DNA
 <213> Artificial sequence
 15
 <220>
 <223> PCR primer
 20 <400> 159
 acctccttc ccagctccc 20
 <210> 160
 25 <211> 24
 <212> DNA
 <213> Artificial sequence
 30
 <220>
 <223> PCR primer
 35 <400> 160
 ggcaacatct tacttgtct ttga 24
 <210> 161
 <211> 25
 40 <212> DNA
 <213> Artificial sequence
 45
 <220>
 <223> PCR primer
 50 <400> 161
 ccaaggaagc acagacaact atttc 25
 <210> 162
 <211> 30
 55

<212> DNA
 <213> Artificial sequence
 5
 <220>
 <223> PCR primer
 <400> 162
 tctctccctat ccatggcact aaaccacttc 30
 <210> 163
 15 <211> 19
 <212> DNA
 <213> Artificial sequence
 20
 <220>
 <223> PCR primer
 <400> 163
 25 tgggcaaggg ctctctatct 19
 <210> 164
 30 <211> 21
 <212> DNA
 <213> Artificial sequence
 35
 <220>
 <223> PCR primer
 <400> 164
 40 gttaccctcg gcagacgtat g 21
 <210> 165
 45 <211> 31
 <212> DNA
 <213> Artificial sequence
 50
 <220>
 <223> PCR primer
 <400> 165
 55 tgcctctgag tctgaatctc ccaagagag a 31

5 <210> 166
 <211> 31
 <212> DNA
 <213> Artificial sequence
 10
 <220>
 <223> PCR primer
 15 <400> 166
 gagtagttat gtgattatatt cagctcttga c 31
 <210> 167
 20 <211> 21
 <212> DNA
 <213> Artificial sequence
 25
 <220>
 <223> PCR primer
 30 <400> 167
 tcaaatgttg tccccgagtc t 21
 <210> 168
 35 <211> 34
 <212> DNA
 <213> Artificial sequence
 40
 <220>
 <223> PCR primer
 45 <400> 168
 cagaattcg gaagacagaa ctattgtcat gcct 34
 <210> 169
 <211> 27
 50 <212> DNA
 <213> Artificial sequence
 55

	<220>		
5	<223> PCR primer		
	<400> 169		
	gattagtaac ccatagcagt tgaaggt	27	
	<210> 170		
10	<211> 26		
	<212> DNA		
	<213> Artificial sequence		
15	<220>		
	<223> PCR primer		
20	<400> 170		
	atttactgac ggtggtotga acatac	26	
	<210> 171		
25	<211> 31		
	<212> DNA		
	<213> Artificial sequence		
30	<220>		
	<223> PCR primer		
35	<400> 171		
	tgacagactc caaatcacaa gcacagtoaa c	31	
	<210> 172		
	<211> 25		
40	<212> DNA		
	<213> Artificial sequence		
45	<220>		
	<223> PCR primer		
50	<400> 172		
	tgatggtttg gaggaaagtt tatatt	25	
	<210> 173		
55	<211> 24		

	<212> DNA	
5	<213> Artificial sequence	
	<220>	
10	<223> PCR primer	
	<400> 173	
	tttggttggg tottttagagg aatc	24
	<210> 174	
15	<211> 24	
	<212> DNA	
20	<213> Artificial sequence	
	<220>	
25	<223> PCR primer	
	<400> 174	
	tgccaacccat gcatcaggta gcc	24
	<210> 175	
30	<211> 20	
	<212> DNA	
35	<213> Artificial sequence	
	<220>	
40	<223> PCR primer	
	<400> 175	
	cagctcacct ggcaacttca	20
	<210> 176	
45	<211> 20	
	<212> DNA	
50	<213> Artificial sequence	
	<220>	
55	<223> PCR primer	
	<400> 176	
	cctgattttc ccagcgatgt	20

<210> 177
 5 <211> 19
 <212> DNA
 <213> Artificial sequence
 10
 <220>
 <223> PCR primer
 15 <400> 177
 agccgctccc ggttatgct 19
 <210> 178
 20 <211> 20
 <212> DNA
 <213> Artificial sequence
 25
 <220>
 <223> PCR primer
 30 <400> 178
 tggccaagcg taagctgatt 20
 <210> 179
 35 <211> 21
 <212> DNA
 <213> Artificial sequence
 40
 <220>
 <223> PCR primer
 45 <400> 179
 gctgcagtga tcggatcato t 21
 <210> 180
 50 <211> 22
 <212> DNA
 <213> Artificial Sequence
 55

<220>
 5 <223> MLLT6
 <400> 180
 caccatggag cccatcgtgc tg 22
 <210> 181
 10 <211> 19
 <212> DNA
 <213> Artificial Sequence
 15
 <220>
 <223> MLLT6 for
 20 <400> 181
 atccccgagg tgcaatttg 19
 <210> 182
 25 <211> 21
 <212> DNA
 <213> Artificial Sequence
 30
 <220>
 <223> MLLT6 rev
 35 <400> 182
 agcgatcatg aggcacgtac t 21
 <210> 183
 40 <211> 29
 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 <223> ZNF144
 50 <400> 183
 cctgccagag ataggagacc cagacagct 29
 <210> 184
 <211> 19
 55

<212> DNA
 <213> Artificial Sequence
 5
 <220>
 <223> ZNF144 for
 <400> 184
 atccccctga gccttttoa 19
 <210> 185
 15 <211> 19
 <212> DNA
 <213> Artificial Sequence
 20
 <220>
 <223> ZNF144 rev
 <400> 185
 cagcctctgg tcccaccat 19
 <210> 186
 30 <211> 28
 <212> DNA
 <213> Artificial Sequence
 35
 <220>
 <223> PIP5K2B
 <400> 186
 tgatcatcaa ttccaaacct ctccgaa 28
 <210> 187
 45 <211> 19
 <212> DNA
 <213> Artificial Sequence
 50
 <220>
 <223> PIP5K2B for
 <400> 187
 ccccatggtg ttccgaaac 19
 55

```

5      <210> 188
      <211> 19
      <212> DNA
      <213> Artificial Sequence

10

      <220>

      <223> PIP5K2B rev
      <400> 188
      tgccaggagc ctccataacc
15
      <210> 189
      <211> 29
      <212> DNA
      <213> Artificial Sequence

20

      <220>

      <223> TEM7
      <400> 189
      cagcctttcta aaacacaatg tattcatgt
25
      <210> 190
      <211> 29
      <212> DNA
      <213> Artificial Sequence

30

      <220>

      <223> TEM7 for
      <400> 190
      cctgaactta atggtagaat tcaaagatc
35
      <210> 191
      <211> 27
      <212> DNA
      <213> Artificial Sequence

40

      <220>

      <223> TEM7 for
      <400> 190
      cctgaactta atggtagaat tcaaagatc
45
      <210> 191
      <211> 27
      <212> DNA
      <213> Artificial Sequence

50

      <220>

      <223> TEM7 for
      <400> 190
      cctgaactta atggtagaat tcaaagatc
55
      <210> 191
      <211> 27
      <212> DNA
      <213> Artificial Sequence

```

	<220>	
5	<223> TEM7 rev <400> 191 tattaaacct gagaatccat gcagaga	27
	<210> 192	
10	<211> 35	
	<212> DNA	
	<213> Artificial Sequence	
15	<220>	
20	<223> ZNFN1A3 <400> 192 tatctggtct cagggattgc toctatgtat tcagc	35
	<210> 193	
25	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
30	<220>	
35	<223> ZNFN1A3 for <400> 193 cacagagccc tgctgaagtg	20
	<210> 194	
40	<211> 23	
	<212> DNA	
	<213> Artificial Sequence	
45	<220>	
50	<223> ZNFN1A3 rev <400> 194 gcgaggtcat tggtttttag aaa	23
	<210> 195	
55	<211> 22	

	<212> DNA	
5	<213> Artificial Sequence	
	<220>	
10	<223> WIRE	
	<400> 195	
	ctgtgatccg aaatgggtgcc ag	22
	<210> 196	
15	<211> 20	
	<212> DNA	
20	<213> Artificial Sequence	
	<220>	
25	<223> WIRE for	
	<400> 196	
	cogtctccac atccaaacct	20
	<210> 197	
30	<211> 20	
	<212> DNA	
35	<213> Artificial Sequence	
	<220>	
40	<223> WIRE rev	
	<400> 197	
	acctatgcac ttggtatggt	20
	<210> 198	
45	<211> 21	
	<212> DNA	
50	<213> Artificial Sequence	
	<220>	
55	<223> PSMB3	
	<400> 198	
	agtggcaact ggcgcgaaca a	21


```

5      <210> 199
        <211> 21
        <212> DNA
        <213> Artificial Sequence

10     <220>

        <223> PSMB3 for
        <400> 199
        ccccatggtg actgatgact t                21

        <210> 200

        <211> 21
        <212> DNA
        <213> Artificial Sequence

20     <220>

        <223> PSMB3 rev
        <400> 200
        ccagaggggac tcacacattc c            21

        <210> 201

        <211> 29
        <212> DNA
        <213> Artificial Sequence

35     <220>

        <223> MGC9753
        <400> 201
        ccagaaactt tccatccc aa aggcagtct    29

        <210> 202

        <211> 21
        <212> DNA
        <213> Artificial Sequence

50     <220>

        <223> MGC9753
        <400> 201
        ccagaaactt tccatccc aa aggcagtct    29

        <210> 202

        <211> 21
        <212> DNA
        <213> Artificial Sequence

55     <220>

        <223> MGC9753
        <400> 201
        ccagaaactt tccatccc aa aggcagtct    29

        <210> 202

        <211> 21
        <212> DNA
        <213> Artificial Sequence

```

	<220>	
5	<223> MGC9753 for	
	<400> 202	
	ctgcccaca ggaatagaat g	21
	<210> 203	
10	<211> 23	
	<212> DNA	
	<213> ARTIFICIAL SEQUENCE	
15	<220>	
	<223> MGC9753 rev	
20	<400> 203	
	aaaaatccag tctgcttcaa cca	23
	<210> 204	
25	<211> 20	
	<212> DNA	
	<213> ARTIFICIAL SEQUENCE	
30	<220>	
	<223> ORMDL3	
35	<400> 204	
	agctgccca gctccaogga	20
	<210> 205	
	<211> 21	
40	<212> DNA	
	<213> ARTIFICIAL SEQUENCE	
45	<220>	
	<223> ORMDL3 for	
50	<400> 205	
	tccctgatga gctgcttat c	21
	<210> 206	
55	<211> 28	

<212> DNA
 5 <213> ARTIFICIAL SEQUENCE

 <220>
 10 <223> ORMDL3 rev
 <400> 206
 tctcagtact tattgattcc aaaaatcc 28

 <210> 207
 15 <211> 25
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 20

 <220>
 25 <223> MGC15482
 <400> 207
 tccagtggaa gcaaccccag tgttc 25

 <210> 208
 30 <211> 25
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 35

 <220>
 40 <223> MGC15482 for
 <400> 208
 cacttctaga gctaccgtgg agtct 25

 <210> 209
 45 <211> 22
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 50

 <220>
 55 <223> MGC15482 rev
 <400> 209
 cactcacttt gtaacccttg ct 22

<210> 210
 5 <211> 20
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 10
 <220>
 <223> PPP1R1B
 15 <400> 210
 cagcgtggcg caacaaccca 20
 <210> 211
 20 <211> 21
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 25
 <220>
 <223> PPP1R1B for
 30 <400> 211
 gggattgttt cgccacacat a 21
 <210> 212
 35 <211> 20
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 40
 <220>
 <223> PPP1R1B rev
 45 <400> 212
 ccgatgttaa ggcccatagc 20
 <210> 213
 50 <211> 27
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 55

	<220>	
5	<223> MGC14832	
	<400> 213	
	taaaaatgtcc ggccaacatg agttccc	27
	<210> 214	
10	<211> 17	
	<212> DNA	
	<213> ARTIFICIAL SEQUENCE	
15	<220>	
	<223> MGC14832 for	
20	<400> 214	
	cgccagtgcct ggcacat	17
	<210> 215	
25	<211> 20	
	<212> DNA	
	<213> ARTIFICIAL SEQUENCE	
30	<220>	
	<223> MGC14832 rev	
35	<400> 215	
	gacacccccct gacctatgga	20
	<210> 216	
	<211> 25	
40	<212> DNA	
	<213> ARTIFICIAL SEQUENCE	
45	<220>	
	<223> LOC51242	
50	<400> 216	
	cagtgacctc tccggttccc ttgga	25
	<210> 217	
55	<211> 20	

	<212> DNA	
5	<213> ARTIFICIAL SEQUENCE	
	<220>	
10	<223> LOC51242 for <400> 217 tgggtcaatg tgcctctctc	20
	<210> 218	
15	<211> 20	
	<212> DNA	
20	<213> ARTIFICIAL SEQUENCE	
	<220>	
25	<223> LOC51242 for <400> 218 agggtcagga gggagaaaac	20
	<210> 219	
30	<211> 26	
	<212> DNA	
35	<213> ARTIFICIAL SEQUENCE	
	<220>	
40	<223> FLJ20291 <400> 219 ccagtgcaca ccggttaaag agtcaa	26
	<210> 220	
45	<211> 24	
	<212> DNA	
50	<213> ARTIFICIAL SEQUENCE	
	<220>	
55	<223> FLJ20291 for <400> 220 ttgtgggaca ctcagtaact ttgg	24

<210> 221
 5 <211> 20
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 10
 <220>
 <223> FLJ20291 rev
 15 <400> 221
 acaagcactc ccacgagat 20
 <210> 222
 20 <211> 24
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 25
 <220>
 <223> PRO2521
 30 <400> 222
 agttctgtcct cactgccatc gcc 24
 <210> 223
 35 <211> 21
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 40
 <220>
 <223> PRO2521 for
 45 <400> 223
 aagcctctgg gttttccatt t 21
 <210> 224
 50 <211> 20
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 55

	<220>	
5	<223> PRO2521 rev	
	<400> 224	
	cccactggtg acaggatggt	20
	<210> 225	
10	<211> 23	
	<212> DNA	
	<213> ARTIFICIAL SEQUENCE	
15	<220>	
	<223> LINK-GEFII	
20	<400> 225	
	catctgacat cttcccggtg gag	23
	<210> 226	
25	<211> 21	
	<212> DNA	
	<213> ARTIFICIAL SEQUENCE	
30	<220>	
	<223> LINK-GEFII for	
35	<400> 226	
	ctttgcaaga tgtctcaacc a	21
	<210> 227	
40	<211> 18	
	<212> DNA	
	<213> ARTIFICIAL SEQUENCE	
45	<220>	
	<223> LINK-GEFII rev	
50	<400> 227	
	tttcccggtg agcaggaa	18
	<210> 228	
55	<211> 26	

<212> DNA
 <213> ARTIFICIAL SEQUENCE
 5
 <220>
 <223> CTEN
 <400> 228
 10 ccgcgccta atatcaaca ttaggg 26
 <210> 229
 <211> 23
 15 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 20
 <220>
 <223> CTEN for
 <400> 229
 25 cgagtattcc aaagctggta tcg 23
 <210> 230
 <211> 24
 30 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 35
 <220>
 <223> CTEN rev
 <400> 230
 40 atcacagaga gatggccott atct 24
 <210> 231
 <211> 25
 45 <212> DNA
 <213> Artificial Sequence
 50
 <220>
 <223> D178946 forward primer
 <400> 231
 55 acagtctatc aagcagaaaa atcct 25

<210> 232
 5 <211> 16
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> D17S946 reverse primer
 15 <400> 232
 tgccgtgccg gagaga 16
 <210> 233
 20 <211> 20
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> D17S1181 forward primer
 30 <400> 233
 gacaacagag cgagactccc 20
 <210> 234
 35 <211> 20
 <212> DNA
 <213> Artificial Sequence
 40
 <220>
 <223> D17S1181 reverse primer
 45 <400> 234
 gccagcctg tcacttattc 20
 <210> 235
 50 <211> 18
 <212> DNA
 <213> Artificial Sequence
 55

	<220>		
5	<223> D17S2026 forward primer		
	<400> 235		
	tggtcattcg acaacgaa	18	
	<210> 236		
10	<211> 18		
	<212> DNA		
	<213> Artificial Sequence		
15	<220>		
	<223> D17S2026 reverse primer		
20	<400> 236		
	cagcattgga tgcaatcc	18	
	<210> 237		
25	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
30	<220>		
	<223> D17S838 forward primer		
35	<400> 237		
	ctccagaatc cagaccatga	20	
	<210> 238		
40	<211> 20		
	<212> DNA		
	<213> Artificial Sequence		
45	<220>		
	<223> D17S838 reverse primer		
50	<400> 238		
	aggacagtgt gtagcccttc	20	
	<210> 239		
55	<211> 20		

<212> DNA
 <213> Artificial Sequence
 5

<220>
 <223> D178250 forward primer
 10 <400> 239
 ggaagaatca aatagacaat 20
 <210> 240
 15 <211> 24
 <212> DNA
 <213> Artificial Sequence
 20

<220>
 <223> D178250 reverse primer
 25 <400> 240
 gctggccata tatatatatta aacc 24
 <210> 241
 30 <211> 23
 <212> DNA
 <213> Artificial Sequence
 35

<220>
 <223> D1781818 forward primer
 40 <400> 241
 cataggtatg ttcagaaatg tga 23
 <210> 242
 45 <211> 18
 <212> DNA
 <213> Artificial Sequence
 50

<220>
 <223> D1781818 reverse primer
 55 <400> 242
 tgcctactg aaaccaga 18

<210> 243
 5 <211> 23
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> D17S614 forward primer
 15 <400> 243 23
 aaggggaagg ggctttcaaa gct
 <210> 244
 20 <211> 23
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> D17S614 reverse primer
 30 <220>
 <221> misc_feature
 <222> (1)..(1)
 35 <223> n=a, c, g or t
 <400> 244
 40 nggaggttgc agtgagccaa gat 23
 <210> 245
 <211> 23
 <212> DNA
 45 <213> Artificial Sequence
 <220>
 50 <223> D17S2019 forward primer
 <400> 245 23
 caaaagctta tgatgctcaa acc
 55

<210> 246
 5 <211> 22
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> D17S2019 reverse primer
 15 <400> 246
 ttgtttccct ttgactttct ga 22
 <210> 247
 <211> 25
 20 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> D17S608 forward primer
 30 <400> 247
 taggttcacc tctcattttc ttcag 25
 <210> 248
 <211> 24
 35 <212> DNA
 <213> Artificial Sequence
 40
 <220>
 <223> D17S608 reverse primer
 45 <220>
 <221> misc_feature
 <222> (17)..(17)
 50 <223> n=a, c, g or t
 <400> 248
 atctaatgat ttataaact tata 24
 55

<211> 20
 5 <212> DNA
 <213> Artificial Sequence

 10 <220>
 <223> D17S1655 forward primer
 <400> 249
 cggaaccagag tgttccatgg 20
 15 <210> 250
 <211> 20
 <212> DNA
 20 <213> Artificial Sequence

 25 <220>
 <223> D17S1655 reverse primer
 <400> 250
 gcatacagca ccctctacct 20
 30 <210> 251
 <211> 25
 <212> DNA
 35 <213> Artificial Sequence

 40 <220>
 <223> D17S2147 forward primer
 <400> 251
 aggggagaat aaataaaatc tgttg 25
 45 <210> 252
 <211> 22
 <212> DNA
 50 <213> Artificial Sequence

 55

<220>
 5 <223> D1782147 reverse primer
 <400> 252
 caggagtgg acactctcca tg 22
 <210> 253
 10 <211> 22
 <212> DNA
 <213> Artificial Sequence
 15
 <220>
 20 <223> D178754 forward primer
 <400> 253
 tggattacct gactcagcct gc 22
 <210> 254
 25 <211> 22
 <212> DNA
 <213> Artificial Sequence
 30
 <220>
 35 <223> D178754 reverse primer
 <400> 254
 gogtgtctgt ctccatgtgt gc 22
 <210> 255
 40 <211> 18
 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 50 <223> D1781814 forward primer
 <400> 255
 tccccaatga cgggtgatg 18
 <210> 256
 55 <211> 20

<212> DNA
 <213> Artificial Sequence
 5
 <220>
 <223> D17S1814 reverse primer
 <400> 256
 ctggaggttg gcttgtggat 20
 <210> 257
 15 <211> 18
 <212> DNA
 <213> Artificial Sequence
 20
 <220>
 <223> D17S2007 forward primer
 <400> 257
 ggtccaaga atttgotg 18
 <210> 258
 30 <211> 20
 <212> DNA
 <213> Artificial Sequence
 35
 <220>
 <223> D17S2007 reverse primer
 <400> 258
 ccaccagaa aaacaggaga 20
 <210> 259
 40 <211> 20
 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 <223> D17S1246 forward primer
 <400> 259
 togatctcct gaccttgtga 20
 50
 55

<210> 260
 5 <211> 20
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> D17S1246 reverse primer
 15 <400> 260
 ttgtcaccac attgccttcc 20
 <210> 261
 20 <211> 21
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> D17S1979 forward primer
 30 <400> 261
 ccttgatag attcagctcc c 21
 <210> 262
 35 <211> 21
 <212> DNA
 <213> Artificial Sequence
 40
 <220>
 <223> D17S1979 reverse primer
 45 <400> 262
 cttgtccctt ctcaatctcc c 21
 <210> 263
 50 <211> 25
 <212> DNA
 <213> Artificial Sequence
 55

<220>
 <223> D17S1984 forward primer
 5 <400> 263
 ttaagcaagg tttaattaa gctgc 25
 <210> 264
 <211> 21
 <212> DNA
 <213> Artificial Sequence
 15
 <220>
 <223> D17S1984 reverse primer
 20 <400> 264
 gattacagtg ctccctctcc c 21
 <210> 265
 <211> 22
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> G11580 forward primer
 30 <400> 265
 gggtttaatt aagatgcatg gc 22
 <210> 266
 <211> 21
 <212> DNA
 <213> Artificial Sequence
 35
 <220>
 <223> G11580 reverse primer
 40 <400> 266
 gattacagtg ctccctctcc c 21
 <210> 267
 <211> 20
 45
 50
 55

	<212> DNA	
	<213> Artificial Sequence	
5		
	<220>	
10	<223> D17S1867 forward primer	
	<400> 267	
	agtttgacac tgaggctttg	20
	<210> 268	
15	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
20		
	<220>	
25	<223> D17S1867 reverse primer	
	<400> 268	
	tttagacttg gtaactgcgc	20
	<210> 269	
30	<211> 24	
	<212> DNA	
	<213> Artificial Sequence	
35		
	<220>	
40	<223> D17S1788 forward primer	
	<400> 269	
	tgcaatgcc taagaacttt tcag	24
	<210> 270	
45	<211> 19	
	<212> DNA	
	<213> Artificial Sequence	
50		
	<220>	
55	<223> D17S1788 reverse primer	
	<400> 270	
	gccatgatct cccaaagcc	19

<210> 271
 5 <211> 18
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> D17S1836 forward primer
 15 <400> 271 18
 togaggttat ggtgagcc
 <210> 272
 20 <211> 24
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> D17S1836 reverse primer
 30 <400> 272 24
 aaactgtgtg tgtcaaagga tact
 <210> 273
 35 <211> 19
 <212> DNA
 <213> Artificial Sequence
 40
 <220>
 <223> D17S1787 forward primer
 45 <400> 273 19
 gctgatctga agccaatga
 <210> 274
 50 <211> 19
 <212> DNA
 <213> Artificial Sequence
 55

	<220>		
5	<223> D17S1787 reverse primer		
	<400> 274		
	tacatgaagg catggtctg	19	
	<210> 275		
10	<211> 23		
	<212> DNA		
	<213> Artificial Sequence		
15	<220>		
	<223> D17S1660 forward primer		
20	<400> 275		
	ctaataataat cctgggcaca tgg	23	
	<210> 276		
25	<211> 18		
	<212> DNA		
	<213> Artificial Sequence		
30	<220>		
	<223> D17S1660 reverse primer		
35	<400> 276		
	gctgoggacc agacagat	18	
	<210> 277		
40	<211> 22		
	<212> DNA		
	<213> Artificial Sequence		
45	<220>		
	<223> D17S2154 forward primer		
50	<400> 277		
	gataaaaaca agcactggct cc	22	
	<210> 278		
55	<211> 20		

<212> DNA
 <213> Artificial Sequence
 5
 <220>
 <223> D17S2154 reverse primer
 <400> 278
 cccaagggtt tcttgatcta 20
 <210> 279
 15 <211> 21
 <212> DNA
 <213> Artificial Sequence
 20
 <220>
 <223> D17S1955 forward primer
 <400> 279
 tgtaatgtaa gcccatgag g 21
 <210> 280
 30 <211> 25
 <212> DNA
 <213> Artificial Sequence
 35
 <220>
 <223> D17S1955 reverse primer
 <400> 280
 caactaactc aacagtctaa aggtg 25
 <210> 281
 40 <211> 25
 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 <223> D17S2098 forward primer
 <400> 281
 gtgagttcaa gcatagtaat tatoc 25
 50
 55

<210> 282
 5 <211> 23
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> D17S2098 reverse primer
 15 <400> 282 23
 attcagcctc agttcactgc ttc
 <210> 283
 20 <211> 20
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> D17S518 forward primer
 30 <400> 283 20
 gatccagtgg agactcagag
 <210> 284
 35 <211> 20
 <212> DNA
 <213> Artificial Sequence
 40
 <220>
 <223> D17S518 reverse primer
 45 <400> 284 20
 tagtctctgg gacaccaga
 <210> 285
 50 <211> 25
 <212> DNA
 <213> Artificial Sequence
 55

<220>
 5 <223> D17S518 forward primer
 <400> 285
 atttctgagt gtctacctg ttgag 25
 <210> 286
 10 <211> 17
 <212> DNA
 <213> Artificial Sequence
 15
 <220>
 20 <223> D17S518 reverse primer
 <400> 286
 actgactgag ccaactgc 17
 <210> 287
 25 <211> 20
 <212> DNA
 <213> Artificial Sequence
 30
 <220>
 35 <223> D11S4358 forward primer
 <400> 287
 togagaagga caaaatcacc 20
 <210> 288
 40 <211> 20
 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 50 <223> D11S4358 reverse primer
 <400> 288
 gaacagggtt agtccattcg 20
 <210> 289
 55 <211> 19

	<212> DNA	
5	<213> Artificial Sequence	
	<220>	
10	<223> D17S964 forward primer	
	<400> 289	19
	gtttcttctc cttgtggg	
	<210> 290	
15	<211> 19	
	<212> DNA	
20	<213> Artificial Sequence	
	<220>	
25	<223> D17S964 reverse primer	
	<400> 290	19
	agtcagctga gattgtgcc	
	<210> 291	
30	<211> 20	
	<212> DNA	
35	<213> Artificial Sequence	
	<220>	
40	<223> D19S1091 forward primer	
	<400> 291	20
	caagccaaga catccagtt	
	<210> 292	
45	<211> 20	
	<212> DNA	
50	<213> Artificial Sequence	
	<220>	
55	<223> D19S1091 reverse primer	
	<400> 292	20
	ccccacac agctcatatg	

<210> 293
 5 <211> 22
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> D17S1179 forward primer
 15 <400> 293 22
 tttttctctct cattccattg gg
 <210> 294
 20 <211> 20
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> D17S1179 reverse primer
 30 <400> 294 20
 gcaacagagg gagactccaa
 <210> 295
 35 <211> 19
 <212> DNA
 <213> Artificial Sequence
 40
 <220>
 <223> D10S2160 forward primer
 45 <400> 295 19
 tccatcccg taagacctc
 <210> 296
 50 <211> 25
 <212> DNA
 <213> Artificial Sequence
 55

<220>
 5 <223> D1082160 reverse primer
 <400> 296
 tatggagtac ctactctatg ccagg 25
 <210> 297
 10 <211> 20
 <212> DNA
 <213> Artificial Sequence
 15
 <220>
 <223> D17S1230 forward primer
 20 <400> 297
 attcaaaagct ggatoccttt 20
 <210> 298
 25 <211> 20
 <212> DNA
 <213> Artificial Sequence
 30
 <220>
 <223> D17S1230 reverse primer
 35 <400> 298
 agctgtgaca aatgcctgta 20
 <210> 299
 40 <211> 20
 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 <223> D17S1338 forward primer
 50 <400> 299
 tcacctgaga ttgggagacc 20
 <210> 300
 55 <211> 18

	<212> DNA	
5	<213> Artificial Sequence	
	<220>	
10	<223> D17S1338 reverse primer	
	<400> 300	
	aagatggggc aggaatgg	18
	<210> 301	
15	<211> 19	
	<212> DNA	
20	<213> Artificial Sequence	
	<220>	
25	<223> D17S2011 forward primer	
	<400> 301	
	tcaatgtcct ccaagccag	19
	<210> 302	
30	<211> 20	
	<212> DNA	
35	<213> Artificial Sequence	
	<220>	
40	<223> D17S2011 reverse primer	
	<400> 302	
	aaacaccaca ctctcccctg	20
	<210> 303	
45	<211> 20	
	<212> DNA	
50	<213> Artificial Sequence	
	<220>	
55	<223> D17S2011 forward primer	
	<400> 303	
	ttcttgggct tcccgtagcc	20

<210> 304
 5 <211> 20
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> D17S2011 reverse primer
 15 <400> 304
 ggggcagacg actctcctt 20
 <210> 305
 20 <211> 23
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> D17S2038 forward primer
 30 <400> 305
 ggggatacaa cotttaaagt tcc 23
 <210> 306
 35 <211> 25
 <212> DNA
 <213> Artificial Sequence
 40
 <220>
 <223> D17S2038 reverse primer
 45 <400> 306
 attcacctaa tgaggattct tcttt 25
 <210> 307
 50 <211> 24
 <212> DNA
 <213> Artificial Sequence
 55

<220>
 <223> D17S2091 forward primer
 5 <400> 307
 gctgaatag ccatottgag ctac 24
 <210> 308
 <211> 23
 10 <212> DNA
 <213> Artificial Sequence
 15
 <220>
 <223> D17S2091 reverse primer
 20 <400> 308
 tcgcactct ttttaagag cac 23
 <210> 309
 <211> 24
 25 <212> DNA
 <213> Artificial Sequence
 30
 <220>
 <223> D17S649 forward primer
 35 <400> 309
 otttcaactot ttcagctgaa gagg 24
 <210> 310
 <211> 25
 40 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 <223> D17S649 reverse primer
 50 <400> 310
 tgaogtgota tttcctgttt tgtot 25
 <210> 311
 <211> 18
 55

<212> DNA

<213> Artificial Sequence

<220>

No.	Doccode	Number of pages
1	1001	6
2	1002	1
3	DESC	4
4	CLMS	1
5	DRAW	1
6	ABST	1
7	PRIODOC	8

<220>

<223> D17S1190 reverse primer

<400> 312

caacacacta ccccaqga

Claims

1. A method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least 2 markers **characterized in that** the markers are genes and fragments thereof or genomic nucleic acid sequences that are located on one chromosomal region which is altered in malignant neoplasia.
2. A method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least 2 markers **characterized in that** the markers are:
 - a) genes that are located on one or more chromosomal region(s) which is/are altered in malignant neoplasia; and
 - b)
 - i) receptor and ligand; or
 - ii) members of the same signal transduction pathway; or
 - iii) members of synergistic signal transduction pathways; or
 - iv) members of antagonistic signal transduction pathways; or
 - v) transcription factor and transcription factor binding site.
3. The method of claim 1 or 2 wherein the malignant neoplasia is breast cancer, ovarian cancer, gastric cancer, colon cancer, esophageal cancer, mesenchymal cancer, bladder cancer or non-small cell lung cancer.
4. The method of claim 1 or 2 wherein at least one chromosomal region is defined as the cytogenetic region: 1p13, 1q32, 3p21-p24, 5p13-p14, 8q23-q²⁴, 1 1q13, 12q13, 17q12-q24 or 20q13.

5. The method of claim 1 or 2 wherein at least chromosomal region is defined as the cytogenetic region 17q1 1.2-21.3 and the malignant neoplasia is breast cancer, ovarian cancer, gastric cancer, colon cancer, esophageal cancer, mesenchymal cancer, bladder cancer or non-small cell lung cancer.

6. The method of claim 1 or 2 wherein at least one chromosomal region is defined as the cytogenetic region 3p21-24 and the malignant neoplasia is breast cancer, ovarian cancer, gastric cancer, colon cancer, esophageal cancer, mesenchymal cancer, bladder cancer or non-small cell lung cancer.

7. The method of claim 1 or 2 wherein at least one chromosomal region is defined as the cytogenetic region 12q13 and the malignant neoplasia is breast cancer, ovarian cancer, gastric cancer, colon cancer, esophageal cancer, mesenchymal cancer, bladder cancer or non-small cell lung cancer.

8. A method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least one marker whereby the marker is a VNTR, SNP, RFLP or STS **characterized in that** the marker is located on one chromosomal region which is altered in malignant neoplasia due to amplification and the marker is detected in a cancerous and a non-cancerous tissue or biological sample of the same individual.

9. The method of claim 8 wherein the marker is selected from the group consisting of the VNTRs:

D17S946, D17S1181, D17S2026, D17S838, D17S250, D17S1818, D17S614, D17S2019, D17S608, D17S1655, D17S2147, D17S754, D17S1814, D17S2007, D17S1246, D17S1979, D17S1984, D17S1984, D17S1867, D17S1788, D17S1836, D17S1787, D17S1660, D17S2154, D17S1955, D17S2098, D17S518, D17S1851, D11S4358, D17S964, D19S1091, D17S1179, D10S2160, D17S1230, D17S1338, D17S2011, D17S1237, D17S2038, D17S2091, D17S649, D17S1190 and M87506.

10. The method of claim 8 wherein the marker is selected from the group consisting of the SNPs:

rs2230698, rs2230700, rs1058808, rs1801200, rs903506, rs2313170, rs1136201, rs2934968, rs2172826, rs1810132, rs1801201, rs2230702, rs2230701, rs1126503, rs3471, rs13695, rs471692, rs558068, rs1064288, rs1061692, rs520630, rs782774, rs565121, rs2586112, rs532299, rs2732786, rs1804539, rs1804538, rs1804537, rs1141364 rs12231, rs1132259 rs1132257, rs113225 rs1132254, rs113225 rs1132268 and rs11322

11. A method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least one marker **characterized in that** the marker is selected from:

a) a polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75 ;

b) a polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) and encodes a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

c) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (c) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (d)

e) a purified polypeptide encoded by a polynucleotide or polynucleotide analog sequence specified in (a) to (e)

f) a purified polypeptide comprising at least one of the sequences of SEQ ID NO: 28 to 32, 34, 35, 37 to 42, 44, 45, 47 to 52 or 76 to 98;

are detected.

12. A method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least 2 markers

characterized in that at least 2 markers are selected from:

a) a polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 1 to 26 or 53 to 75;

b) a polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) and encodes a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

c) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c)

e) a purified polypeptide encoded by a polynucleotide sequence or polynucleotide analog specified in (a) to (d)

f) a purified polypeptide comprising at least one of the sequences of SEQ ID NO: 27 to 52 or 76 to 98

are detected.

13. The method of any of the claims 1 or 12 wherein the detection method comprises the use of PCR, arrays or beads.

14. A diagnostic kit comprising instructions for conducting the method of any of claims 1 to 13.

15. A composition for the prediction, diagnosis or prognosis of malignant neoplasia comprising:

a) a detection agent for:

i) any polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

ii) any polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

iii) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

iv) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c)

v) a polypeptide encoded by a polynucleotide or polynucleotide analog sequence specified in (a) to (d);

vi) a polypeptide comprising at least one of the sequences of SEQ ID NO: 28 to 32, 34, 35, 37 to 42, 44, 45, 47 to 52 or 76 to 98.

or

b) at least 2 detection agents for at least 2 markers selected from:

i) any polynucleotide comprising at least one of the sequences of SEQ ID NO: 1 to 26 or 53 to 75;

ii) any polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

iii) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

iv) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c)

v) a polypeptide encoded by a polynucleotide sequence specified in (a) to (d);

vi) a polypeptide comprising at least one of the sequences of SEQ ID NO: 27 to 52 or 76 to 98.

16. An array comprising a plurality of polynucleotides or polynucleotide analogs wherein each of the polynucleotides is selected from:

a) a polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 1 to 26 or 53 to 75;

b) a polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

c) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c)

attached to a solid support.

17. A method of screening for agents which regulate the activity of a polypeptide encoded by a polynucleotide or polynucleotide analog selected from the group consisting of:

a) a polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

b) a polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

c) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);

comprising the steps of:

i) contacting a test compound with at least one polypeptide encoded by a polynucleotide specified in (a) to (d); and

ii) detecting binding of the test compound to the polypeptide, wherein a test compound which binds to the polypeptide is identified as a potential therapeutic agent for modulating the activity of the polypeptide in order to prevent or treat malignant neoplasia.

18. A method of screening for agents which regulate the activity of a polypeptide encoded by a polynucleotide or polynucleotide analog selected from the group consisting of:

a) a polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

b) a polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

c) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c)

comprising the steps of:

i) contacting a test compound with at least one polypeptide encoded by a polynucleotide specified in (a) to (d); and

ii) detecting the activity of the polypeptide as specified for the respective sequence in Table 2 or 3, wherein a test compound which increases the activity is identified as a potential preventive or therapeutic agent for increasing the polypeptide activity in malignant neoplasia, and wherein a test compound which decreases the activity of the polypeptide is identified as a potential therapeutic agent for decreasing the polypeptide activity in malignant neoplasia.

19. A method of screening for agents which regulate the activity of a polynucleotide or polynucleotide analog selected from group consisting of:

a) a polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

b) a polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

c) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c)

comprising the steps of:

i) contacting a test compound with at least one polynucleotide or polynucleotide analog specified in (a) to (d), and

ii) detecting binding of the test compound to the polynucleotide, wherein a test compound which binds to the polynucleotide is identified as a potential preventive or therapeutic agent for regulating the activity of the polynucleotide in malignant neoplasia.

20. Use of

a) a polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide or polynucleotide analog specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective

sequence in Table 2 or 3;

c) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3;

d) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);

e) an antisense molecule targeting specifically one of the polynucleotide sequences specified in (a) to (d);

f) a purified polypeptide encoded by a polynucleotide or polynucleotide analog sequence specified in (a) to (d)

g) a purified polypeptide comprising at least one of the sequences of SEQ ID NO: 28 to 32, 34, 35, 37 to 42, 44, 45, 47 to 52 or 76 to 98;

h) an antibody capable of binding to one of the polynucleotide specified in (a) to (d) or a polypeptide specified in (f) and (g);

i) a reagent identified by any of the methods of claim 17 to 19 that modulates the amount or activity of a polynucleotide sequence specified in (a) to (d) or a polypeptide specified in (f) and (g);

in the preparation of a composition for the prevention, prediction, diagnosis, prognosis or a medicament for the treatment of malignant neoplasia.

21. Use of claim 20 wherein the disease is breast cancer.

22. A reagent that regulates the activity of a polypeptide selected from the group consisting of:

a) a polypeptide encoded by any polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

b) a polypeptide encoded by any polynucleotide or polynucleotide analog which hybridizes under stringent conditions to any polynucleotide comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75 encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

c) a polypeptide encoded by any polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polypeptide encoded by any polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

e) or a polypeptide comprising at least one of the sequences of SEQ ID NO: 28 to 32, 34, 35, 37 to 42, 44, 45, 47 to 52 or 76 to 98;

wherein said reagent is identified by the method of any of the claims 17 to 19.

23. A reagent that regulates the activity of a polynucleotide or polynucleotide analog selected from the group consisting of:

a) a polynucleotide or polynucleotide analog comprising at least one of the sequences SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

b) a polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective

sequence in Table 2 or 3

c) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

wherein said reagent is identified by the method of any of the claims 17 to 19.

24. A pharmaceutical composition, comprising:

a) an expression vector containing at least one polynucleotide or polynucleotide analog selected from the group consisting of:

i) a polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

ii) a polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

iii) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

iv) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3;

or the reagent of claim 22 or 23 and a pharmaceutically acceptable carrier.

25. A computer-readable medium comprising:

a) at least one digitally encoded value representing a level of expression of at least one polynucleotide sequence of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75

b) at least 2 digitally encoded values representing the levels of expression of at least 2 polynucleotide sequences selected from SEQ ID NO: 1 to 26 or 53 to 75

in a cell from the a subject at risk for or having malignant neoplasia.

26. A method for the detection of chromosomal alterations characterized in that the relative abundance of individual mRNAs, encoded by genes, located in altered chromosomal regions is detected.

27. A method for the detection of chromosomal alterations characterized in that the copy number of one or more chromosomal region(s) is detected by quantitative PCR.

Figure J





Figure 1




Figure 3




Figure 4

